

Basic Linear Algebra, Metric Spaces, Differential Calculus
and Nonlinear Programming¹
Fall 2017

antonio villanacci

October 13, 2017

¹I would like to thank the following friends for helpful comments and discussions: Laura Carosi, Michele Gori, Marina Pireddu and all the students and teaching assistants of the courses I used these notes for in the past several years.

Contents

I	Basic Linear Algebra	7
1	Systems of linear equations	9
1.1	Linear equations and solutions	9
1.2	Systems of linear equations, equivalent systems and elementary operations	10
1.3	Systems in triangular and echelon form	11
1.4	Reduction algorithm	12
1.5	Matrices	14
1.6	Systems of linear equations and matrices	18
1.7	Exercises	19
2	The Euclidean Space \mathbb{R}^n	21
2.1	Sum and scalar multiplication	21
2.2	Scalar product	22
2.3	Norms and Distances	23
2.4	Exercises	25
3	Matrices	27
3.1	Matrix operations	28
3.2	Inverse matrices	33
3.3	Elementary matrices	34
3.4	Elementary column operations	41
3.5	Exercises	44
4	Vector spaces	45
4.1	Definition	45
4.2	Examples	47
4.3	Vector subspaces	48
4.4	Linear combinations	49
4.5	Row and column space of a matrix	50
4.6	Linear dependence and independence	54
4.7	Basis and dimension	60
4.8	Coordinates	63
4.9	Row and column span	64
4.10	Exercises	66
5	Determinant and rank of a matrix	67
5.1	Definition and properties of the determinant of a matrix	67
5.2	Rank of a matrix	71
5.3	Inverse matrices (continued)	72
5.4	Span of a matrix, linearly independent rows and columns, rank	74
5.5	Exercises	76

6	Linear functions	77
6.1	Definition	77
6.2	Kernel and Image of a linear function	78
6.3	Nonsingular functions and isomorphisms	81
6.4	Exercises	84
7	Linear functions and matrices	85
7.1	From a linear function to the associated matrix	85
7.2	From a matrix to the associated linear function	87
7.3	$\mathbb{M}(m, n)$ and $\mathcal{L}(V, U)$ are isomorphic	88
7.4	Some related properties of a linear function and associated matrix	90
7.5	Some facts on $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$	93
7.6	Examples of computation of $[l]_{\mathbf{v}}^{\mathbf{u}}$	95
7.7	Exercises	96
7.8	Appendices.	96
7.8.1	The dual and double dual space of a vector space	96
7.8.2	Vector spaces as Images or Kernels of well chosen linear functions	99
8	Solutions to systems of linear equations	103
8.1	Some preliminary basic facts	103
8.2	A solution method: Rouchè-Capelli's and Cramer's theorems	104
8.3	Exercises	113
II	Some topology in metric spaces	115
9	Metric spaces	117
9.1	Definitions and examples	117
9.2	Open and closed sets	121
9.2.1	Sets which are open or closed in metric subspaces.	125
9.3	Sequences	127
9.4	Sequential characterization of closed sets	130
9.5	Compactness	130
9.5.1	Compactness and bounded, closed sets	131
9.5.2	Sequential compactness	133
9.6	Completeness	139
9.6.1	Cauchy sequences	139
9.6.2	Complete metric spaces	140
9.6.3	Completeness and closedness	141
9.7	Fixed point theorem: contractions	142
9.8	Appendices.	144
9.8.1	Some characterizations of open and closed sets	144
9.8.2	Norms and metrics	148
9.9	Exercises	151
10	Functions	153
10.1	Limits of functions	153
10.2	Continuous Functions	154
10.3	Continuous functions on compact sets	158
10.4	Exercises	160
11	Correspondence, maximum theorem and a fixed point theorem	161
11.1	Continuous Correspondences	161
11.2	The Maximum Theorem	168
11.3	Fixed point theorems	171
11.4	Application of the maximum theorem to the consumer problem	172

III	Differential calculus in Euclidean spaces	175
12	Partial derivatives and directional derivatives	177
12.1	Partial Derivatives	177
12.2	Directional Derivatives	178
13	Differentiability	185
13.1	Total Derivative and Differentiability	185
13.2	Total Derivatives in terms of Partial Derivatives.	187
14	Some Theorems	189
14.1	The chain rule	190
14.2	Mean value theorem	192
14.3	A sufficient condition for differentiability	195
14.4	A sufficient condition for equality of mixed partial derivatives	195
14.5	Taylor's theorem for real valued functions	195
15	Implicit function theorem	197
15.1	Some intuition	197
15.2	Functions with full rank square Jacobian	199
15.3	The inverse function theorem	202
15.4	The implicit function theorem	204
15.5	Some geometrical remarks on the gradient	206
15.6	Extremum problems with equality constraints.	206
15.7	Exercises on part III	208
IV	Nonlinear programming	209
16	Convex sets	211
16.1	Definition	211
16.2	Separation of convex sets	211
16.3	Farkas' Lemma	213
17	Concave functions	217
17.1	Different Kinds of Concave Functions	217
17.1.1	Concave Functions.	218
17.1.2	Strictly Concave Functions.	220
17.1.3	Quasi-Concave Functions.	222
17.1.4	Strictly Quasi-concave Functions.	226
17.1.5	Pseudo-concave Functions.	228
17.2	Relationships among Different Kinds of Concavity	229
17.2.1	Hessians and Concavity.	232
18	Maximization Problems	235
18.1	The case of inequality constraints: Kuhn-Tucker theorems	235
18.1.1	On uniqueness of the solution	239
18.2	The Case of Equality Constraints: Lagrange Theorem.	240
18.3	The Case of Both Equality and Inequality Constraints.	242
18.4	Main Steps to Solve a (Nice) Maximization Problem	244
18.4.1	Some problems and some solutions	249
18.5	The Implicit Function Theorem and Comparative Statics Analysis	251
18.5.1	Maximization problem without constraint	251
18.5.2	Maximization problem with equality constraints	252
18.5.3	Maximization problem with Inequality Constraints	252
18.6	The Envelope Theorem and the meaning of multipliers	254
18.6.1	The Envelope Theorem	254
18.6.2	On the meaning of the multipliers	255

19 Applications to Economics	257
19.1 The Walrasian Consumer Problem	257
19.2 Production	259
19.3 The demand for insurance	261
19.4 Exercises on part IV	262
 V Problem Sets	 263
20 Exercises	265
20.1 Linear Algebra	265
20.2 Some topology in metric spaces	270
20.2.1 Basic topology in metric spaces	270
20.2.2 Correspondences	273
20.3 Differential Calculus in Euclidean Spaces	274
20.4 Nonlinear Programming	277
 21 Solutions	 279
21.1 Linear Algebra	279
21.2 Some topology in metric spaces	288
21.2.1 Basic topology in metric spaces	288
21.2.2 Correspondences	294
21.3 Differential Calculus in Euclidean Spaces	296
21.4 Nonlinear Programming	302

Part I

Basic Linear Algebra

Chapter 1

Systems of linear equations

1.1 Linear equations and solutions

Definition 1 ¹ A linear equation in the unknowns x_1, x_2, \dots, x_n is an equation of the form

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = b, \quad (1.1)$$

where $b \in \mathbb{R}$ and $\forall j \in \{1, \dots, n\}$, $a_j \in \mathbb{R}$. The real number a_j is called the coefficient of x_j and b is called the constant of the equation. a_j for $j \in \{1, \dots, n\}$ and b are also called parameters of system (1.1).

Definition 2 A solution to the linear equation (1.1) is an ordered n -tuple $(\bar{x}_1, \dots, \bar{x}_n) := (\bar{x}_j)_{j=1}^n$ such² that the following statement (obtained by substituting \bar{x}_j in the place of x_j for any j) is true:

$$a_1\bar{x}_1 + a_2\bar{x}_2 + \dots + a_n\bar{x}_n = b,$$

The set of all such solutions is called the solution set or the general solution or, simply, the solution of equation (1.1).

The following fact is well known.

Proposition 3 Let the linear equation

$$ax = b \quad (1.2)$$

in the unknown (variable) $x \in \mathbb{R}$ and parameters $a, b \in \mathbb{R}$ be given. Then,

1. if $a \neq 0$, then $x = \frac{b}{a}$ is the unique solution to (1.2);
2. if $a = 0$ and $b \neq 0$, then (1.2) has no solutions;
3. if $a = 0$ and $b = 0$, then any real number is a solution to (1.2).

Definition 4 A linear equation (1.1) is said to be degenerate if $\forall j \in \{1, \dots, n\}$, $a_j = 0$, i.e., it has the form

$$0x_1 + 0x_2 + \dots + 0x_n = b, \quad (1.3)$$

Clearly,

1. if $b \neq 0$, then equation (1.3) has no solution,
2. if $b = 0$, any n -tuple $(\bar{x}_j)_{j=1}^n$ is a solution to (1.3).

Definition 5 Let a nondegenerate equation of the form (1.1) be given. The leading unknown of the linear equation (1.1) is the first unknown with a nonzero coefficient, i.e., x_p is the leading unknown if

$$\forall j \in \{1, \dots, p-1\}, a_j = 0 \quad \text{and} \quad a_p \neq 0.$$

For any $j \in \{1, \dots, n\} \setminus \{p\}$, x_j is called a free variable - consistently with the following obvious result.

¹In this part, I often follow Lipschutz (1991).

²“ $:=$ ” means “equal by definition”.

Proposition 6 Consider a nondegenerate linear equation $a_1x_1 + a_2x_2 + \dots + a_nx_n = b$ with leading unknown x_p . Then the set of solutions to that equation is

$$\left\{ (x_k)_{k=1}^n : \forall j \in \{1, \dots, n\} \setminus \{p\}, x_j \in \mathbb{R} \text{ and } x_p = \frac{b - \sum_{j \in \{1, \dots, n\} \setminus \{p\}} a_j x_j}{a_p} \right\}$$

1.2 Systems of linear equations, equivalent systems and elementary operations

Definition 7 A system of m linear equations in the n unknowns x_1, x_2, \dots, x_n is a system of the form

$$\begin{cases} a_{11}x_1 + \dots + a_{1j}x_j + \dots + a_{1n}x_n = b_1 \\ \dots \\ a_{i1}x_1 + \dots + a_{ij}x_j + \dots + a_{in}x_n = b_i \\ \dots \\ a_{m1}x_1 + \dots + a_{mj}x_j + \dots + a_{mn}x_n = b_m \end{cases} \quad (1.4)$$

where $\forall i \in \{1, \dots, m\}$ and $\forall j \in \{1, \dots, n\}$, $a_{ij} \in \mathbb{R}$ and $\forall i \in \{1, \dots, m\}$, $b_i \in \mathbb{R}$. We call L_i the i -th linear equation of system (1.4).

A solution to the above system is an ordered n -tuple $(\bar{x}_j)_{j=1}^n$ which is a solution of each equation of the system. The set of all such solutions is called the solution set of the system.

Definition 8 Systems of linear equations are equivalent if their solutions set is the same.

The following fact is obvious.

Proposition 9 Assume that a system of linear equations contains the degenerate equation

$$L : \quad 0x_1 + 0x_2 + \dots + 0x_n = b.$$

1. If $b = 0$, then L may be deleted from the system without changing the solution set;
2. if $b \neq 0$, then the system has no solutions.

A way to solve a system of linear equations is to transform it in an equivalent system whose solution set is “easy” to be found. In what follows we make precise the above sentence.

Definition 10 An elementary operation on a system of linear equations (1.4) is one of the following operations:

- [E₁] Interchange L_i with L_j , an operation denoted by $L_i \leftrightarrow L_j$ (which we can read “put L_i in the place of L_j and L_j in the place of L_i ”);
- [E₂] Multiply L_i by $k \in \mathbb{R} \setminus \{0\}$, denoted by $kL_i \rightarrow L_i$, $k \neq 0$ (which we can read “put kL_i in the place of L_i , with $k \neq 0$ ”);
- [E₃] Replace L_i by (k times L_j plus L_i), denoted by $(L_i + kL_j) \rightarrow L_i$ (which we can read “put $L_i + kL_j$ in the place of L_i ”).

Sometimes we apply [E₂] and [E₃] in one step, i.e., we perform the following operation

- [E] Replace L_i by (k' times L_j and $k \in \mathbb{R} \setminus \{0\}$ times L_i), denoted by $(k'L_j + kL_i) \rightarrow L_i$, $k \neq 0$.

Elementary operations are important because of the following obvious result.

Proposition 11 If S_1 is a system of linear equations obtained from a system S_2 of linear equations using a finite number of elementary operations, then system S_1 and S_2 are equivalent.

In what follows, first we define two types of “simple” systems (triangular and echelon form systems), and we see why those systems are in fact “easy” to solve. Then, we show how to transform any system in one of those “simple” systems.

Proposition 17 *Let a system in echelon form with r equations and s variables be given. Then, the following results hold true.*

1. If $s = r$, i.e., the number of unknowns is equal to the number of equations, then the system has a unique solution;
2. if $s > r$, i.e., the number of unknowns is greater than the number of equations, then we can arbitrarily assign values to the $n - r > 0$ free variables and obtain solutions of the system.

Proof. We prove the theorem by induction on the number r of equations of the system.

Step 1. $r = 1$.

In this case, we have a single, nondegenerate linear equation, to which Proposition 6 applies if $s > r = 1$, and Proposition 3 applies if $s = r = 1$.

Step 2.

Assume that $r > 1$ and the desired conclusion is true for a system with $r - 1$ equations. Consider the given system in the form (1.6) and erase the first equation, so obtaining the following system:

$$\left\{ \begin{array}{ccccccc} a_{2j_2}x_{j_2} & + \dots & + a_{2,j_3}x_{j_3} & + \dots & & & = b_2 \\ & & a_{3,j_3}x_{j_3} & + \dots & & & \\ & & & \dots & & & \\ & & & a_{r,j_r}x_{j_r} & + a_{r,j_r+1} & + \dots + a_{rs}x_s & = b_r \end{array} \right. \quad (1.7)$$

in the unknowns x_{j_2}, \dots, x_s . First of all observe that the above system is in echelon form and has $r - 1$ equation; therefore we can apply the induction argument distinguishing the two case $s > r$ and $s = r$.

If $s > r$, then we can assign arbitrary values to the free variables, whose number is (the “old” number minus the erased ones)

$$s - r - (j_2 - j_1 - 1) = s - r - j_2 + 2$$

and obtain a solution of system (1.7). Consider the first equation of the original system

$$a_{11}x_1 + a_{12}x_2 + \dots + a_{1,j_2-1}x_{j_2-1} + a_{1j_2}x_{j_2} + \dots = b_1. \quad (1.8)$$

We immediately see that the above found values together with arbitrary values for the additional

$$j_2 - 2$$

free variable of equation (1.8) yield a solution of that equation, as desired. Observe also that the values given to the variables x_1, \dots, x_{j_2-1} from the first equation do satisfy the other equations simply because their coefficients are zero there.

If $s = r$, the system in echelon form, in fact, becomes a system in triangular form and then the solution exists and it is unique. ■

Remark 18 From the proof of the previous Proposition, if the echelon system (1.6) contains more unknowns than equations, i.e., $s > r$, then the system has an infinite number of solutions since each of the $s - r \geq 1$ free variables may be assigned an arbitrary real number.

1.4 Reduction algorithm

The following algorithm (sometimes called row reduction) reduces system (1.4) of m equation and n unknowns to either echelon form, or triangular form, or shows that the system has no solution. The algorithm then gives a proof of the following result.

Proposition 19 Any system of linear equations has either

1. infinite solutions, or
2. a unique solution, or
3. no solutions.

Reduction algorithm.

Consider a system of the form (1.4) such that

$$\forall j \in \{1, \dots, n\}, \quad \exists i \in \{1, \dots, m\} \text{ such that } a_{ij} \neq 0, \quad (1.9)$$

i.e., a system in which each variable has a nonzero coefficient in at least one equation. If that is not the case, the remaining variables can be renamed in order to have (1.9) satisfied.

Step 1. Interchange equations so that the first unknown, x_1 , appears with a nonzero coefficient in the first equation; i.e., rearrange the equations in the system in order to have $a_{11} \neq 0$.

Step 2. Use a_{11} as a “pivot” to eliminate x_1 from all equations but the first equation. That is, for each $i > 1$, apply the elementary operation

$$[E_3] : -\left(\frac{a_{i1}}{a_{11}}\right) L_1 + L_i \rightarrow L_i$$

or

$$[E] : -a_{i1}L_1 + a_{11}L_i \rightarrow L_i.$$

Step 3. Examine each new equation L :

1. If L has the form

$$0x_1 + 0x_2 + \dots + 0x_n = 0,$$

or if L is a multiple of another equation, then delete L from the system.³

2. If L has the form

$$0x_1 + 0x_2 + \dots + 0x_n = b,$$

with $b \neq 0$, then exit the algorithm. The system has no solutions.

Step 4. Repeat Steps 1, 2 and 3 with the subsystem formed by all the equations, excluding the first equation.

Step 5. Continue the above process until the system is in echelon form or a degenerate equation is obtained in Step 3.2.

Summarizing, our method for solving system (1.4) consists of two steps:

Step A. Use the above reduction algorithm to reduce system (1.4) to an equivalent simpler system (in triangular form, system (1.5) or echelon form (1.6)).

Step B. If the system is in triangular form, use back-substitution to find the solution; if the system is in echelon form, bring the free variables on the right hand side of each equation, give them arbitrary values (say, the name of the free variable with an upper bar), and then use back-substitution.

Example 20

$$\begin{cases} x_1 & + & 2x_2 & + & (-3)x_3 & = & -1 \\ 3x_1 & + & (-1)x_2 & + & 2x_3 & = & 7 \\ 5x_1 & + & 3x_2 & + & (-4)x_3 & = & 2 \end{cases}$$

Step A.

Step 1. Nothing to do.

³The justification of Step 3 is Proposition 9 and the fact that if $L = kL'$ for some other equation L' in the system, then the operation $-kL' + L \rightarrow L$ replace L by $0x_1 + 0x_2 + \dots + 0x_n = 0$, which again may be deleted by Proposition 9.

Step 2. Apply the operations

$$-3L_1 + L_2 \rightarrow L_2$$

and

$$-5L_1 + L_3 \rightarrow L_3,$$

to get

$$\begin{cases} x_1 + 2x_2 + (-3)x_3 = -1 \\ (-7)x_2 + 11x_3 = 10 \\ (-7)x_2 + 11x_3 = 7 \end{cases}$$

Step 3. Examine each new equations L_2 and L_3 :

1. L_2 and L_3 do not have the form

$$0x_1 + 0x_2 + \dots + 0x_n = 0;$$

L_2 is not a multiple L_3 ;

2. L_2 and L_3 do not have the form

$$0x_1 + 0x_2 + \dots + 0x_n = b,$$

Step 4.

Step 1.1 Nothing to do.

Step 2.1 Apply the operation

$$-L_2 + L_3 \rightarrow L_3$$

to get

$$\begin{cases} x_1 + 2x_2 + (-3)x_3 = -1 \\ (-7)x_2 + 11x_3 = 10 \\ 0x_1 + 0x_2 + 0x_3 = -3 \end{cases}$$

Step 3.1 L_3 has the form

$$0x_1 + 0x_2 + \dots + 0x_n = b,$$

1. with $b = -3 \neq 0$, then exit the algorithm. The system has no solutions.

1.5 Matrices

Definition 21 Given $m, n \in \mathbb{N}$, a matrix (of real numbers) of order $m \times n$ is a table of real numbers with m rows and n columns as displayed below.

$$\begin{bmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \dots & & & & & \\ a_{i1} & a_{i2} & \dots & a_{ij} & \dots & a_{in} \\ \dots & & & & & \\ a_{m1} & a_{m2} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix}$$

For any $i \in \{1, \dots, m\}$ and any $j \in \{1, \dots, n\}$ the real numbers a_{ij} are called entries of the matrix; the first subscript i denotes the row the entries belongs to, the second subscript j denotes the column the entries belongs to. We will usually denote matrices with capital letters and we will write $A_{m \times n}$ to denote a matrix of order $m \times n$. Sometimes it is useful to denote a matrix by its "typical" element and we write $[a_{ij}]_{\substack{i \in \{1, \dots, m\} \\ j \in \{1, \dots, n\}}}$, or simply $[a_{ij}]$ if no ambiguity arises about the number of rows and columns. For $i \in \{1, \dots, m\}$,

$$[a_{i1} \quad a_{i2} \quad \dots \quad a_{ij} \quad \dots \quad a_{in}]$$

is called the i – th row of A and it denoted by $R^i(A)$. For $j \in \{1, \dots, n\}$,

$$\begin{bmatrix} a_{1j} \\ a_{2j} \\ \dots \\ a_{ij} \\ \dots \\ a_{mj} \end{bmatrix}$$

is called the j – th column of A and it denoted by $C^j(A)$.

We denote the set of $m \times n$ matrices by $\mathcal{M}_{m,n}$, and we write, in an equivalent manner, $A_{m \times n}$ or $A \in \mathcal{M}_{m,n}$.

Definition 22 The matrix

$$A_{m \times 1} = \begin{bmatrix} a_1 \\ \dots \\ a_m \end{bmatrix}$$

is called column vector and the matrix

$$A_{1 \times n} = [a_1, \dots, a_n]$$

is called row vector. We usually denote row or column vectors by small Latin letters.

Definition 23 The first nonzero entry in a row R of a matrix $A_{m \times n}$ is called the leading nonzero entry of R . If R has no leading nonzero entries, i.e., if every entry in R is zero, then R is called a zero row. If all the rows of A are zero, i.e., each entry of A is zero, then A is called a zero matrix, denoted by $0_{m \times n}$ or simply 0 , if no confusion arises.

In the previous sections, we defined triangular and echelon systems of linear equations. Below, we define triangular, echelon matrices and a special kind of echelon matrices. In Section (1.6), we will see that there is a simple relationship between systems and matrices.

Definition 24 A matrix $A_{m \times n}$ is square if $m = n$. A square matrix A belonging to $\mathcal{M}_{m,m}$ is called square matrix of order m .

Definition 25 Given $A = [a_{ij}] \in \mathcal{M}_{m,m}$, the main diagonal of A is made up by the entries a_{ii} with $i \in \{1, \dots, m\}$.

Definition 26 A square matrix $A = [a_{ij}] \in \mathcal{M}_{m,m}$ is an upper triangular matrix or simply a triangular matrix if all entries below the main diagonal are equal to zero, i.e., $\forall i, j \in \{1, \dots, m\}$, if $i > j$, then $a_{ij} = 0$.

Definition 27 $A \in \mathcal{M}_{m,m}$ is called diagonal matrix of order m if any element outside the principal diagonal is equal to zero, i.e., $\forall i, j \in \{1, \dots, m\}$ such that $i \neq j$, $a_{ij} = 0$.

Definition 28 A matrix $A \in \mathcal{M}_{m,n}$ is called an echelon (form) matrix, or it is said to be in echelon form, if the following two conditions hold:

1. All zero rows, if any, are on the bottom of the matrix.
2. The leading nonzero entry of each row is to the right of the leading nonzero entry in the preceding row.

Definition 29 If a matrix A is in echelon form, then its leading nonzero entries are called pivot entries, or simply, pivots

Remark 30 If a matrix $A \in \mathcal{M}_{m,n}$ is in echelon form and r is the number of its pivot entries, then $r \leq \min\{m, n\}$. In fact, $r \leq m$, because the matrix may have zero rows and $r \leq n$, because the leading nonzero entries of the first row maybe not in the first column, and the other leading nonzero entries may be “strictly to the right” of previous leading nonzero entry.

Definition 31 A matrix $A \in \mathcal{M}_{m,n}$ is called in row canonical form if

1. it is in echelon form,
2. each pivot is 1, and
3. each pivot is the only nonzero entry in its column.

Example 32 1. All the matrices below are echelon matrices; only the fourth one is in row canonical form.

$$\begin{bmatrix} 0 & 7 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & -3 & 3 \\ 0 & 0 & 0 & 0 & 0 & 7 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 2 & 3 & 2 & 0 & 1 & 2 & 4 \\ 0 & 0 & 1 & 1 & -3 & 3 & 0 \\ 0 & 0 & 0 & 0 & 0 & 7 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 & 3 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 3 & 0 & 0 & 4 \\ 0 & 0 & 0 & 1 & 0 & -3 \\ 0 & 0 & 0 & 0 & 1 & 2 \end{bmatrix}.$$

2. Any zero matrix is in row canonical form.

Remark 33 Let a matrix $A_{m \times n}$ in row canonical form be given. As a consequence of the definition, we have what follows.

1. If some rows from A are erased, the resulting matrix is still in row canonical form.
2. If some columns of zeros are added, the resulting matrix is still in row canonical form.

Definition 34 Denote by R^i the i -th row of a matrix A . An elementary row operation is one of the following operations on the rows of A :

- $[E_1]$ (Row interchange) Interchange R^i with R^j , an operation denoted by $R^i \leftrightarrow R^j$ (which we can read “put R^i in the place of R^j and R^j in the place of R^i ”);
- $[E_2]$ (Row scaling) Multiply R^i by $k \in \mathbb{R} \setminus \{0\}$, denoted by $kR^i \rightarrow R^i$, $k \neq 0$ (which we can read “put kR^i in the place of R^i , with $k \neq 0$ ”);
- $[E_3]$ (Row addition) Replace R^i by (k times R^j plus R^i), denoted by $(R^i + kR^j) \rightarrow R^i$ (which we can read “put $R^i + kR^j$ in the place of R^i ”).

Sometimes we apply $[E_2]$ and $[E_3]$ in one step, i.e., we perform the following operation

- $[E]$ Replace R^i by (k' times R^j and $k \in \mathbb{R} \setminus \{0\}$ times R^i), denoted by $(k'R^j + kR^i) \rightarrow R^i$, $k \neq 0$.

Definition 35 A matrix $A \in \mathcal{M}_{m,n}$ is said to be row equivalent to a matrix $B \in \mathcal{M}_{m,n}$ if B can be obtained from A by a finite number of elementary row operations.

It is hard not to recognize the similarity of the above operations and those used in solving systems of linear equations.

We use the expression “row reduce” as having the meaning of “transform a given matrix into another matrix using row operations”. The following algorithm “row reduces” a matrix A into a matrix in echelon form.

Row reduction algorithm to echelon form.

Consider a matrix $A = [a_{ij}] \in \mathcal{M}_{m,n}$.

Step 1. Find the first column with a nonzero entry. Suppose it is column j_1 .

Step 2. Interchange the rows so that a nonzero entry appears in the first row of column j_1 , i.e., so that $a_{1j_1} \neq 0$.

Step 3. Use a_{1j_1} as a “pivot” to obtain zeros below a_{1j_1} , i.e., for each $i > 1$, apply the row operation

$$[E_3] : -\left(\frac{a_{ij_1}}{a_{1j_1}}\right) R^1 + R^i \rightarrow R^i$$

or

$$[E] : -a_{ij_1} R^1 + a_{11} R^i \rightarrow R^i.$$

Step 4. Repeat Steps 1, 2 and 3 with the submatrix formed by all the rows, excluding the first row.

Step 5. Continue the above process until the matrix is in echelon form.

Example 36 *Let's apply the above algorithm to the following matrix*

$$\begin{bmatrix} 1 & 2 & -3 & -1 \\ 3 & -1 & 2 & 7 \\ 5 & 3 & -4 & 2 \end{bmatrix}$$

Step 1. Find the first column with a nonzero entry: that is C^1 , and therefore $j_1 = 1$.

Step 2. Interchange the rows so that a nonzero entry appears in the first row of column j_1 , i.e., so that $a_{1j_1} \neq 0$: $a_{1j_1} = a_{11} = 1 \neq 0$.

Step 3. Use a_{11} as a “pivot” to obtain zeros below a_{11} . Apply the row operations

$$-3R^1 + R^2 \rightarrow R^2$$

and

$$-5R^1 + R^3 \rightarrow R^3,$$

to get

$$\begin{bmatrix} 1 & 2 & -3 & -1 \\ 0 & -7 & 11 & 10 \\ 0 & -7 & 11 & 7 \end{bmatrix}$$

Step 4. Apply the operation

$$-R^2 + R^3 \rightarrow R^3$$

to get

$$\begin{bmatrix} 1 & 2 & -3 & -1 \\ 0 & -7 & 11 & 10 \\ 0 & 0 & 0 & -3 \end{bmatrix}$$

which is in echelon form.

Row reduction algorithm from echelon form to row canonical form.

Consider a matrix $A = [a_{ij}] \in \mathcal{M}_{m,n}$ in echelon form, say with pivots

$$a_{1j_1}, a_{2j_2}, \dots, a_{rj_r}.$$

Step 1. Multiply the last nonzero row R^r by $\frac{1}{a_{rj_r}}$, so that the leading nonzero entry of that row becomes 1.

Step 2. Use a_{rj_r} as a “pivot” to obtain zeros above the pivot, i.e., for each $i \in \{r-1, r-2, \dots, 1\}$, apply the row operation

$$[E_3] : -a_{i,j_r}R^r + R^i \rightarrow R^i.$$

Step 3. Repeat Steps 1 and 2 for rows $R^{r-1}, R^{r-2}, \dots, R^2$.

Step 4. Multiply R^1 by $\frac{1}{a_{1j_1}}$.

Example 37 *Consider the matrix*

$$\begin{bmatrix} 1 & 2 & -3 & -1 \\ 0 & -7 & 11 & 10 \\ 0 & 0 & 0 & -3 \end{bmatrix}$$

in echelon form, with leading nonzero entries

$$a_{11} = 1, a_{22} = -7, a_{34} = -3.$$

Step 1. Multiply the last nonzero row R^3 by $\frac{1}{-3}$, so that the leading nonzero entry becomes 1:

$$\begin{bmatrix} 1 & 2 & -3 & -1 \\ 0 & -7 & 11 & 10 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Step 2. Use $a_{rj_r} = a_{34}$ as a “pivot” to obtain zeros above the pivot, i.e., for each $i \in \{r-1, r-2, \dots, 1\} = \{2, 1\}$, apply the row operation

$$[E_3] : -a_{i,j_r}R^r + R^i \rightarrow R^i,$$

which in our case are

$$\begin{aligned} -a_{2,4}R^3 + R^2 &\rightarrow R^2 \quad \text{i.e.,} \quad -10R^3 + R^2 \rightarrow R^2, \\ -a_{1,4}R^3 + R^1 &\rightarrow R^1 \quad \text{i.e.,} \quad R^3 + R^1 \rightarrow R^1. \end{aligned}$$

Then, we get

$$\begin{bmatrix} 1 & 2 & -3 & 0 \\ 0 & -7 & 11 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Step 3. Multiply R^2 by $\frac{1}{-7}$, and get

$$\begin{bmatrix} 1 & 2 & -3 & 0 \\ 0 & 1 & -\frac{11}{7} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Use a_{23} as a “pivot” to obtain zeros above the pivot, applying the operation:

$$-2R^2 + R^1 \rightarrow R^1,$$

to get

$$\begin{bmatrix} 1 & 0 & \frac{1}{7} & 0 \\ 0 & 1 & -\frac{11}{7} & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

which is in row reduced form.

Proposition 38 Any matrix $A \in \mathcal{M}_{m,n}$ is row equivalent to a matrix in row canonical form.

Proof. The two above algorithms show that any matrix is row equivalent to at least one matrix in row canonical form. ■

Remark 39 In fact, in Proposition 159, we will show that: Any matrix $A \in \mathcal{M}_{m,n}$ is row equivalent to a **unique** matrix in row canonical form.

1.6 Systems of linear equations and matrices

Definition 40 Given system (1.4), i.e., a system of m linear equation in the n unknowns x_1, x_2, \dots, x_n

$$\begin{cases} a_{11}x_1 + \dots + a_{1j}x_j + \dots + a_{1n}x_n = b_1 \\ \dots \\ a_{i1}x_1 + \dots + a_{ij}x_j + \dots + a_{in}x_n = b_i \\ \dots \\ a_{m1}x_1 + \dots + a_{mj}x_j + \dots + a_{mn}x_n = b_m, \end{cases}$$

the matrix

$$\begin{bmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} & b_1 \\ \dots & & & & & \\ a_{i1} & \dots & a_{ij} & \dots & a_{in} & b_i \\ \dots & & & & & \\ a_{m1} & \dots & a_{mj} & \dots & a_{mn} & b_m \end{bmatrix}$$

is called the augmented matrix M of system (1.4).

Each row of M corresponds to an equation of the system, and each column of M corresponds to the coefficients of an unknown, except the last column which corresponds to the constant of the system.

In an obvious way, given an arbitrary matrix M , we can find a unique system whose associated matrix is M ; moreover, given a system of linear equations, there is only one matrix M associated with it. We can therefore identify system of linear equations with (augmented) matrices.

The coefficient matrix of the system is

$$A = \begin{bmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ \dots & & & & \\ a_{i1} & \dots & a_{ij} & \dots & a_{in} \\ \dots & & & & \\ a_{m1} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix}$$

One way to solve a system of linear equations is as follows:

1. Reduce its augmented matrix M to echelon form, which tells if the system has solution; if M has a row of the form $(0, 0, \dots, 0, b)$ with $b \neq 0$, then the system has no solution and you can stop. If the system admits solutions go to the step below.

2. Reduce the matrix in echelon form obtained in the above step to its row canonical form. Write the corresponding system. In each equation, bring the free variables on the right hand side, obtaining a triangular system. Solve by back-substitution.

The simple justification of this process comes from the following facts:

1. Any elementary row operation of the augmented matrix M of the system is equivalent to applying the corresponding operation on the system itself.
2. The system has a solution if and only if the echelon form of the augmented matrix M does not have a row of the form $(0, 0, \dots, 0, b)$ with $b \neq 0$ - simply because that row corresponds to a degenerate equation.
3. In the row canonical form of the augmented matrix M (excluding zero rows) the coefficient of each non-free variable is a leading nonzero entry which is equal to one and is the only nonzero entry in its respective column; hence the free variable form of the solution is obtained by simply transferring the free variable terms to the other side of each equation.

Example 41 Consider the system presented in Example 20:

$$\begin{cases} x_1 + 2x_2 + (-3)x_3 = -1 \\ 3x_1 + (-1)x_2 + 2x_3 = 7 \\ 5x_1 + 3x_2 + (-4)x_3 = 2 \end{cases}$$

The associated augmented matrix is:

$$\begin{bmatrix} 1 & 2 & -3 & -1 \\ 3 & -1 & 2 & 7 \\ 5 & 3 & -4 & 2 \end{bmatrix}$$

In example 36, we have seen that the echelon form of the above matrix is

$$\begin{bmatrix} 1 & 2 & -3 & -1 \\ 0 & -7 & 11 & 10 \\ 0 & 0 & 0 & -3 \end{bmatrix}$$

which has its last row of the form $(0, 0, \dots, 0, b)$ with $b = -3 \neq 0$, and therefore the system has no solution.

1.7 Exercises

Chapter 1 in Lipschutz:

1,2,3,4,6,7,8,10, 11,12,14,15,16.

Chapter 2

The Euclidean Space \mathbb{R}^n

2.1 Sum and scalar multiplication

It is well known that the real line is a representation of the set \mathbb{R} of real numbers. Similarly, a ordered pair (x, y) of real numbers can be used to represent a point in the plane and a triple (x, y, z) or (x_1, x_2, x_3) a point in the space. In general, if $n \in \mathbb{N} := \{1, 2, \dots\}$, we can define (x_1, x_2, \dots, x_n) or $(x_i)_{i=1}^n$ as a point in the n - space.

Definition 42 $\mathbb{R}^n := \mathbb{R} \times \dots \times \mathbb{R}$.

In other words, \mathbb{R}^n is the Cartesian product of \mathbb{R} multiplied n times by itself.

Definition 43 The elements of \mathbb{R}^n are ordered n -tuple of real numbers and are denoted by

$$x = (x_1, x_2, \dots, x_n) \text{ or } x = (x_i)_{i=1}^n.$$

x_i is called i - th component of $x \in \mathbb{R}^n$.

Definition 44 $x = (x_i)_{i=1}^n \in \mathbb{R}^n$ and $y = (y_i)_{i=1}^n$ are equal if

$$\forall i \in \{1, \dots, n\} , \quad x_i = y_i.$$

In that case we write $x = y$.

Let us introduce two operations on \mathbb{R}^n and analyze some properties they satisfy.

Definition 45 Given $x \in \mathbb{R}^n, y \in \mathbb{R}^n$, we call addition or sum of x and y the element denoted by $x + y \in \mathbb{R}^n$ obtained as follows




$$x + y := (x_i + y_i)_{i=1}^n.$$

Definition 46 An element $\lambda \in \mathbb{R}$ is called scalar.

Definition 47 Given $x \in \mathbb{R}^n$ and $\lambda \in \mathbb{R}$, we call scalar multiplication of x by λ the element $\lambda x \in \mathbb{R}^n$ obtained as follows

$$\lambda x := (\lambda x_i)_{i=1}^n.$$

Geometrical interpretation of the two operations in the case $n = 2$.



From the well known properties of the sum and product of real numbers it is possible to verify that the following properties of the above operations do hold true.

Properties of addition.

- A1. (Associative) $\forall x, y \in \mathbb{R}^n, (x + y) + z = x + (y + z)$;
- A2. (existence of null element) there exists an element e in \mathbb{R}^n such that for any $x \in \mathbb{R}^n$, $x + e = x$; in fact such element is unique and it is denoted by 0;
- A3. (existence of inverse element) $\forall x \in \mathbb{R}^n \exists y \in \mathbb{R}^n$ such that $x + y = 0$; in fact, that element is unique and denoted by $-x$;
- A4. (Commutative) $\forall x, y \in \mathbb{R}^n, x + y = y + x$.

Properties of multiplication.

- M1. (distributive) $\forall \alpha \in \mathbb{R}, x \in \mathbb{R}^n, y \in \mathbb{R}^n \quad \alpha(x + y) = \alpha x + \alpha y$;
- M2. (distributive) $\forall \alpha \in \mathbb{R}, \beta \in \mathbb{R}, x \in \mathbb{R}^n, (\alpha + \beta)x = \alpha x + \beta x$
- M3. $\forall \alpha \in \mathbb{R}, \beta \in \mathbb{R}, x \in \mathbb{R}^n, (\alpha\beta)x = \alpha(\beta x)$;
- M4. $\forall x \in \mathbb{R}^n, 1x = x$.

2.2 Scalar product

Definition 48 Given $x = (x_i)_{i=1}^n, y = (y_i)_{i=1}^n \in \mathbb{R}^n$, we call dot, scalar or inner product of x and y , denoted by xy or $x \cdot y$, the scalar

$$\sum_{i=1}^n x_i \cdot y_i \in \mathbb{R}.$$

Remark 49 The scalar product of elements of \mathbb{R}^n satisfies the following properties.

- 1. $\forall x, y \in \mathbb{R}^n \quad x \cdot y = y \cdot x$;
- 2. $\forall \alpha, \beta \in \mathbb{R}, \forall x, y, z \in \mathbb{R}^n \quad (\alpha x + \beta y) \cdot z = \alpha(x \cdot z) + \beta(y \cdot z)$;
- 3. $\forall x \in \mathbb{R}^n, \quad x \cdot x \geq 0$;
- 4. $\forall x \in \mathbb{R}^n, \quad x \cdot x = 0 \iff x = 0$.

Definition 50 The set \mathbb{R}^n with above described three operations (addition, scalar multiplication and dot product) is usually called Euclidean space of dimension n .

Definition 51 Given $x = (x_i)_{i=1}^n \in \mathbb{R}^n$, we denote the (Euclidean) norm or length of x by


$$\|x\| := (x \cdot x)^{\frac{1}{2}} = \sqrt{\sum_{i=1}^n (x_i)^2}.$$

Geometrical Interpretation of scalar products in \mathbb{R}^2 .

Given $x = (x_1, x_2) \in \mathbb{R}^2 \setminus \{0\}$, from elementary trigonometry we know that

$$x = (\|x\| \cos \alpha, \|x\| \sin \alpha) \tag{2.1}$$

where α is the measure of the angle between the positive part of the horizontal axes and x itself.



Using the above observation we can verify that given $x = (x_1, x_2)$ and $y = (y_1, y_2)$ in $\mathbb{R}^2 \setminus \{0\}$,

$$xy = \|x\| \cdot \|y\| \cdot \cos(\gamma)$$

where γ is an¹ angle between x and y .

scan and insert picture (Marcellini-Sbordone page 179)

From the picture and (2.1), we have

$$x = (\|x\| \cos(\alpha_1), \|x\| \sin(\alpha_1))$$

and

$$y = (\|y\| \cos(\alpha_2), \|y\| \sin(\alpha_2)).$$

Then²

$$xy = \|x\| \|y\| (\cos(\alpha_1) \cdot \cos(\alpha_2) + \sin(\alpha_1) \cdot \sin(\alpha_2)) = \|x\| \|y\| \cos(\alpha_2 - \alpha_1).$$

Taken x and y not belonging to the same line, define $\theta^* :=$ (angle between x and y with minimum measure). From the above equality, it follows that

$$\begin{aligned} \theta^* = \frac{\pi}{2} &\Leftrightarrow x \cdot y = 0 \\ \theta^* < \frac{\pi}{2} &\Leftrightarrow x \cdot y > 0 \\ \theta^* > \frac{\pi}{2} &\Leftrightarrow x \cdot y < 0. \end{aligned}$$

Definition 52 $x, y \in \mathbb{R}^n \setminus \{0\}$ are orthogonal if $xy = 0$.

2.3 Norms and Distances

Proposition 53 (Properties of the norm). Let $\alpha \in \mathbb{R}$ and $x, y \in \mathbb{R}^n$.

1. $\|x\| \geq 0$, and $\|x\| = 0 \Leftrightarrow x = 0$,
2. $\|\alpha x\| = |\alpha| \cdot \|x\|$,
3. $\|x + y\| \leq \|x\| + \|y\|$ (Triangle inequality),
4. $|xy| \leq \|x\| \cdot \|y\|$ (Cauchy-Schwarz inequality).

Proof. 1. By definition $\|x\| = \sqrt{\sum_{i=1}^n (x_i)^2} \geq 0$. Moreover, $\|x\| = 0 \Leftrightarrow \|x\|^2 = 0 \Leftrightarrow \sum_{i=1}^n (x_i)^2 = 0 \Leftrightarrow x = 0$.

$$2. \|\alpha x\| = \sqrt{\sum_{i=1}^n \alpha^2 (x_i)^2} = |\alpha| \sqrt{\sum_{i=1}^n (x_i)^2} = |\alpha| \cdot \|x\|.$$

4. (3 is proved using 4)

We want to show that $|xy| \leq \|x\| \cdot \|y\|$ or $|xy|^2 \leq \|x\|^2 \cdot \|y\|^2$, i.e.,

$$\left(\sum_{i=1}^n x_i y_i \right)^2 \leq \left(\sum_{i=1}^n x_i^2 \right) \cdot \left(\sum_{i=1}^n y_i^2 \right)$$

Defined $X := \sum_{i=1}^n x_i^2$, $Y := \sum_{i=1}^n y_i^2$ and $Z := \sum_{i=1}^n x_i y_i$, we have to prove that

$$Z^2 \leq XY. \tag{2.2}$$

¹ Recall that $\forall x \in \mathbb{R}$, $\cos x = \cos(-x) = \cos(2\pi - x)$.

² Recall that for any $x_1, x_2 \in \mathbb{R}$

$$\cos(x_1 \pm x_2) = \cos(x_1) \cdot \cos(x_2) \mp \sin(x_1) \cdot \sin(x_2),$$

and

$$\cos(x_1) = \cos(-x_1)$$

Observe that

$$\forall a \in \mathbb{R}, \quad \begin{array}{l} 1. \sum_{i=1}^n (ax_i + y_i)^2 \geq 0, \quad \text{and} \\ 2. \sum_{i=1}^n (ax_i + y_i)^2 = 0 \quad \Leftrightarrow \quad \forall i \in \{1, \dots, n\}, ax_i + y_i = 0 \end{array}$$

Moreover,

$$\sum_{i=1}^n (ax_i + y_i)^2 = a^2 \sum_{i=1}^n x_i^2 + 2a \sum_{i=1}^n x_i y_i + \sum_{i=1}^n y_i^2 = a^2 X + 2aZ + Y \geq 0 \quad (2.3)$$

If $X > 0$, we can take $a = -\frac{Z}{X}$, and from (2.3), we get

$$0 \leq \frac{Z^2}{X^2} X - 2 \frac{Z^2}{X} + Y$$

or

$$Z^2 \leq XY,$$

as desired.

If $X = 0$, then $x = 0$ and $Z = 0$, and (2.2) is true simply because $0 \leq 0$.

3. It suffices to show that $\|x + y\|^2 \leq (\|x\| + \|y\|)^2$.

$$\begin{aligned} \|x + y\|^2 &= \sum_{i=1}^n (x_i + y_i)^2 = \sum_{i=1}^n \left((x_i)^2 + 2x_i \cdot y_i + (y_i)^2 \right) = \\ &= \|x\|^2 + 2xy + \|y\|^2 \leq \|x\|^2 + 2|xy| + \|y\|^2 \stackrel{(4 \text{ above})}{\leq} \|x\|^2 + 2\|x\| \cdot \|y\| + \|y\|^2 = (\|x\| + \|y\|)^2. \end{aligned}$$

■

Proposition 54 For any $x, y \in \mathbb{R}^n$ and any $\lambda, \mu \in \mathbb{R}$, we have

1. $\|x\| - \|y\| \leq \|x - y\|$, and
2. $\|\lambda x + \mu y\|^2 = \lambda^2 \|x\|^2 + 2\lambda\mu x \cdot y + \mu^2 \|y\|^2$.

Proof. 1. Recall that $\forall a, b \in \mathbb{R}$

$$-b \leq a \leq b \Leftrightarrow |a| \leq b.$$

From Proposition 53.3, identifying x with $x - y$ and y with y , we get $\|x - y + y\| \leq \|x - y\| + \|y\|$, i.e.,

$$\|x\| - \|y\| \leq \|x - y\|$$

From Proposition 53.3, identifying x with $y - x$ and y with x , we get $\|y - x + x\| \leq \|y - x\| + \|x\|$, i.e.,

$$\|y\| - \|x\| \leq \|y - x\| = \|x - y\|$$

and

$$-\|x - y\| \leq \|x\| - \|y\|,$$

as desired.

2.

$$\begin{aligned} \|\lambda x + \mu y\|^2 &= \sum_{i=1}^n (\lambda x_i + \mu y_i)^2 = \sum_{i=1}^n \left(\lambda^2 (x_i)^2 + 2\lambda\mu (x_i)(y_i) + \mu^2 (y_i)^2 \right) = \\ &= \lambda^2 \sum_{i=1}^n (x_i)^2 + 2\lambda\mu \sum_{i=1}^n (x_i)(y_i) + \mu^2 \sum_{i=1}^n (y_i)^2 = \lambda^2 \|x\|^2 + 2\lambda\mu x \cdot y + \mu^2 \|y\|^2. \end{aligned}$$

■

Definition 55 For any $n \in \mathbb{N} \setminus \{0\}$ and for any $i \in \{1, \dots, n\}$, $e_n^i := (e_{j,n}^i)_{j=1}^n \in \mathbb{R}^n$ with

$$e_{n,j}^i = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}$$

In other words, e_n^i is an element of \mathbb{R}^n whose components are all zero, but the i -th component which is equal to 1. The vector e_n^i is called the i -th canonical vector in \mathbb{R}^n .

Remark 56 $\forall x \in \mathbb{R}^n$,

$$\|x\| \leq \sum_{i=1}^n |x_i|,$$

as verified below.

$$\|x\| = \left\| \sum_{i=1}^n x_i e^i \right\| \stackrel{(1)}{\leq} \sum_{i=1}^n \|x_i e^i\| \stackrel{(2)}{=} \sum_{i=1}^n |x_i| \cdot \|e^i\| = \sum_{i=1}^n |x_i|,$$

where (1) follows from the triangle inequality, i.e., Proposition 53.3, and (2) from Proposition 53.2.

Definition 57 Given $x, y \in \mathbb{R}^n$, we denote the (Euclidean) distance between x and y by

$$d(x, y) := \|x - y\|$$

Proposition 58 (Properties of the distance). Let $x, y, z \in \mathbb{R}^n$.

1. $d(x, y) \geq 0$, and $d(x, y) = 0 \Leftrightarrow x = y$,
2. $d(x, y) = d(y, x)$,
3. $d(x, z) \leq d(x, y) + d(y, z)$ (Triangle inequality).

Proof. 1. It follows from property 1 of the norm.

2. It follows from the definition of the distance as a norm.

3. Identifying x with $x - y$ and y with $y - z$ in property 3 of the norm, we get

$$\|(x - y) + (y - z)\| \leq \|x - y\| + \|y - z\|, \text{ i.e., the desired result. } \blacksquare$$

2.4 Exercises

From Lipschutz (1991), starting from page 53: 2.1 \rightarrow 2.4, 2.12 \rightarrow 2.19, 2.26, 2.27.

Chapter 3

Matrices

We presented the concept of matrix in Definition 21. In this chapter, we study further properties of matrices.

Definition 59 *The transpose of a matrix $A \in \mathcal{M}_{m,n}$, denoted by A^T belongs to $\mathcal{M}_{n,m}$ and it is the matrix obtained by writing the rows of A , in order, as columns:*

$$A^T = \begin{bmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ \dots & & & & \\ a_{i1} & \dots & a_{ij} & \dots & a_{in} \\ \dots & & & & \\ a_{m1} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix}^T = \begin{bmatrix} a_{11} & \dots & a_{i1} & \dots & a_{m1} \\ \dots & & & & \\ a_{1j} & \dots & a_{ij} & \dots & a_{mj} \\ \dots & & & & \\ a_{1n} & \dots & a_{in} & \dots & a_{mn} \end{bmatrix}.$$

In other words, row 1 of the matrix A becomes column 1 of A^T , row 2 of A becomes column 2 of A^T , and so on, up to row m which becomes column m of A^T . Same results is obtained proceeding as follows: column 1 of A becomes row 1 of A^T , column 2 of A becomes row 2 of A^T , and so on, up to column n which becomes row n of A^T . More formally, given $A = [a_{ij}]_{\substack{i \in \{1, \dots, m\} \\ j \in \{1, \dots, n\}}} \in \mathcal{M}_{m,n}$, then

$$A^T = [a_{ji}]_{\substack{j \in \{1, \dots, n\} \\ i \in \{1, \dots, m\}}} \in \mathcal{M}_{n,m}.$$

Definition 60 *A matrix $A \in \mathcal{M}_{n,n}$ is said to be symmetric if $A = A^T$, i.e., $\forall i, j \in \{1, \dots, n\}$, $a_{ij} = a_{ji}$.*

Remark 61 *We can write a matrix $A_{m \times n} = [a_{ij}]$ as*

$$A = \begin{bmatrix} R^1(A) \\ \dots \\ R^i(A) \\ \dots \\ R^m(A) \end{bmatrix} = [C^1(A), \dots, C^j(A), \dots, C^n(A)]$$

where

$$R^i(A) = [a_{i1}, \dots, a_{ij}, \dots, a_{in}] := [R^{i1}(A), \dots, R^{ij}(A), \dots, R^{in}(A)] \in \mathbb{R}^n \quad \text{for } i \in \{1, \dots, m\} \quad \text{and}$$

$$C^j(A) = \begin{bmatrix} a_{1j} \\ a_{ij} \\ a_{mj} \end{bmatrix} := \begin{bmatrix} C^{j1}(A) \\ C^{ji}(A) \\ C^{jm}(A) \end{bmatrix} \in \mathbb{R}^m \quad \text{for } j \in \{1, \dots, n\}.$$

In other words, $R^i(A)$ denotes row i of the matrix A and $C^j(A)$ denotes column j of matrix A .

3.1 Matrix operations

Definition 62 Two matrices $A_{m \times n} := [a_{ij}]$ and $B_{m \times n} := [b_{ij}]$ are equal if for $i \in \{1, \dots, m\}$, $j \in \{1, \dots, n\}$

$$\forall i \in \{1, \dots, m\}, j \in \{1, \dots, n\}, \quad a_{ij} = b_{ij}.$$

Definition 63 Given the matrices $A_{m \times n} := [a_{ij}]$ and $B_{m \times n} := [b_{ij}]$, the sum of A and B , denoted by $A + B$ is the matrix $C_{m \times n} = [c_{ij}]$ such that

$$\forall i \in \{1, \dots, m\}, j \in \{1, \dots, n\}, \quad c_{ij} = a_{ij} + b_{ij}$$

Definition 64 Given the matrices $A_{m \times n} := [a_{ij}]$ and the scalar α , the product of the matrix A by the scalar α , denoted by $\alpha \cdot A$ or αA , is the matrix obtained by multiplying each entry A by α :

$$\alpha A := [\alpha a_{ij}]$$

Remark 65 It is easy to verify that the set of matrices $\mathcal{M}_{m,n}$ with the above defined sum and scalar multiplication satisfies all the properties listed for elements of \mathbb{R}^n in Section 2.1.

Definition 66 Given $A = [a_{ij}] \in \mathcal{M}_{m,n}$, $B = [b_{jk}] \in \mathcal{M}_{n,p}$, the product $A \cdot B$ is a matrix $C = [c_{ik}] \in \mathcal{M}_{m,p}$ such that

$$\forall i \in \{1, \dots, m\}, \forall k \in \{1, \dots, p\}, \quad c_{ik} := \sum_{j=1}^n a_{ij} b_{jk} = R^i(A) \cdot C^k(B)$$

i.e., since

$$A = \begin{bmatrix} R^1(A) \\ \vdots \\ R^i(A) \\ \vdots \\ R^m(A) \end{bmatrix}, B = [C^1(B), \dots, C^k(B), \dots, C^p(B)] \quad (3.1)$$

$$AB = \begin{bmatrix} R^1(A) \cdot C^1(B) & \dots & R^1(A) \cdot C^k(B) & \dots & R^1(A) \cdot C^p(B) \\ \vdots & & \vdots & & \vdots \\ R^i(A) \cdot C^1(B) & \dots & R^i(A) \cdot C^k(B) & \dots & R^i(A) \cdot C^p(B) \\ \vdots & & \vdots & & \vdots \\ R^m(A) \cdot C^1(B) & \dots & R^m(A) \cdot C^k(B) & \dots & R^m(A) \cdot C^p(B) \end{bmatrix} \quad (3.2)$$

Remark 67 If $A \in \mathcal{M}_{1,n}$, $B \in \mathcal{M}_{n,1}$, the above definition coincides with the definition of scalar product between elements of \mathbb{R}^n . In what follows, we often identify an element of \mathbb{R}^n with a row or a column vectors (- see Definition 22) consistently with what we write. In other words $A_{m \times n} x = y$ means that x and y are column vector with n entries, and $w A_{m \times n} = z$ means that w and z are row vectors with m entries.

Definition 68 If two matrices are such that a given operation between them is well defined, we say that they are conformable with respect to that operation.

Remark 69 If $A, B \in \mathcal{M}_{m,n}$, they are conformable with respect to matrix addition. If $A \in \mathcal{M}_{m,n}$ and $B \in \mathcal{M}_{n,p}$, they are conformable with respect to multiplying A on the left of B . We often say the two matrices are conformable and let the context define precisely the sense in which conformability is to be understood.

Remark 70 (For future use) $\forall k \in \{1, \dots, p\}$,

$$A \cdot C^k(B) = \begin{bmatrix} R^1(A) \\ \vdots \\ R^i(A) \\ \vdots \\ R^m(A) \end{bmatrix} \cdot C^k(B) = \begin{bmatrix} R^1(A) \cdot C^k(B) \\ \vdots \\ R^i(A) \cdot C^k(B) \\ \vdots \\ R^m(A) \cdot C^k(B) \end{bmatrix} \quad (3.3)$$

Then, just comparing (3.2) and (3.3), we get

$$AB = \begin{bmatrix} A \cdot C^1(B) & \dots & A \cdot C^k(B) & \dots & A \cdot C^p(B) \end{bmatrix}, \quad (3.4)$$

i.e.,

$$C^k(AB) = A \cdot C^k(B).$$

Similarly, $\forall i \in \{1, \dots, m\}$,

$$\begin{aligned} R^i(A) \cdot B &= R^i(A) \cdot \begin{bmatrix} C^1(B) & \dots & C^k(B) & \dots & C^p(B) \end{bmatrix} = \\ &= \begin{bmatrix} R^i(A) \cdot C^1(B) & \dots & R^i(A) \cdot C^k(B) & \dots & R^i(A) \cdot C^p(B) \end{bmatrix}. \end{aligned} \quad (3.5)$$

Then, just comparing (3.2) and (3.5), we get

$$AB = \begin{bmatrix} R^1(A)B \\ \dots \\ R^i(A)B \\ \dots \\ R^m(A)B \end{bmatrix}, \quad (3.6)$$

i.e.,

$$R^i(AB) = R^i(A) \cdot B.$$

Definition 71 A submatrix of a matrix $A \in M_{m,n}$ is a matrix obtained from A erasing some rows and columns.

Definition 72 A matrix $A \in M_{m,n}$ is partitioned in blocks if it is written as submatrices using a system of horizontal and vertical lines.

Example 73 The matrix

$$\begin{bmatrix} 1 & 2 & 3 & 4 & 5 \\ 6 & 7 & 8 & 9 & 0 \\ 1 & 2 & 3 & 4 & 5 \\ 6 & 7 & 8 & 9 & 0 \end{bmatrix}$$

can be partitioned in block submatrices in several ways. For example as follows

$$\left[\begin{array}{cc|ccc} 1 & 2 & 3 & 4 & 5 \\ 6 & 7 & 8 & 9 & 0 \\ \hline 1 & 2 & 3 & 4 & 5 \\ 6 & 7 & 8 & 9 & 0 \end{array} \right],$$

whose blocks are

$$\begin{bmatrix} 1 & 2 \\ 6 & 7 \end{bmatrix}, \begin{bmatrix} 3 & 4 & 5 \\ 8 & 9 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 \\ 6 & 7 \end{bmatrix}, \begin{bmatrix} 3 & 4 & 5 \\ 8 & 9 & 0 \end{bmatrix}.$$

The reason of the partition into blocks is that the result of operations on block matrices can be obtained by carrying out the computation with blocks, just as if they were actual scalar entries of the matrices, as described below.

Remark 74 We verify below that for matrix multiplication, we do not commit an error if, upon conformably partitioning two matrices, we proceed to regard the partitioned blocks as real numbers and apply the usual rules.

1. Take $a := (a_i)_{i=1}^{n_1} \in \mathbb{R}^{n_1}$, $b := (b_j)_{j=1}^{n_2} \in \mathbb{R}^{n_2}$, $c := (c_i)_{i=1}^{n_1} \in \mathbb{R}^{n_1}$, $d := (d_j)_{j=1}^{n_2} \in \mathbb{R}^{n_2}$,

$$\begin{bmatrix} a & | & b \end{bmatrix}_{1 \times (n_1+n_2)} \begin{bmatrix} c \\ - \\ d \end{bmatrix}_{(n_1+n_2) \times 1} = \sum_{i=1}^{n_1} a_i c_i + \sum_{j=1}^{n_2} b_j d_j = a \cdot c + b \cdot d. \quad (3.7)$$

2.

Take $A \in M_{m,n_1}$, $B \in M_{m,n_2}$, $C \in M_{n_1,p}$, $D \in M_{n_2,p}$, with

$$A = \begin{bmatrix} R^1(A) \\ \dots \\ R^m(A) \end{bmatrix}, B = \begin{bmatrix} R^1(B) \\ \dots \\ R^m(B) \end{bmatrix},$$

$$C = [C^1(C), \dots, C^p(C)],$$

$$D = [C^1(D), \dots, C^p(D)]$$

Then,

$$\begin{aligned} \begin{bmatrix} A & B \end{bmatrix}_{m \times (n_1+n_2)} \begin{bmatrix} C \\ D \end{bmatrix}_{(n_1+n_2) \times p} &= \begin{bmatrix} R^1(A) & R^1(B) \\ \dots & \dots \\ R^m(A) & R^m(B) \end{bmatrix} \begin{bmatrix} C^1(C), \dots, C^p(C) \\ C^1(D), \dots, C^p(D) \end{bmatrix} = \\ &= \begin{bmatrix} R^1(A) \cdot C^1(C) + R^1(B) \cdot C^1(D) & \dots & R^1(A) \cdot C^p(C) + R^1(B) \cdot C^p(D) \\ \dots & \dots & \dots \\ R^m(A) \cdot C^1(C) + R^m(B) \cdot C^1(D) & \dots & R^m(A) \cdot C^p(C) + R^m(B) \cdot C^p(D) \end{bmatrix} = \\ &= \begin{bmatrix} R^1(A) \cdot C^1(C) & \dots & R^1(A) \cdot C^p(C) \\ \dots & \dots & \dots \\ R^m(A) \cdot C^1(C) & \dots & R^m(A) \cdot C^p(C) \end{bmatrix} + \begin{bmatrix} R^1(B) \cdot C^1(D) & \dots & R^1(B) \cdot C^p(D) \\ \dots & \dots & \dots \\ R^m(B) \cdot C^1(D) & \dots & R^m(B) \cdot C^p(D) \end{bmatrix} = \\ &= AC + BD. \end{aligned}$$

Definition 75 Let the matrices $A_i \in \mathcal{M}(n_i, n_i)$ for $i \in \{1, \dots, K\}$, then the matrix

$$A = \begin{bmatrix} A_1 & & & & \\ & A_2 & & & \\ & & \ddots & & \\ & & & A_i & \\ & & & & \ddots \\ & & & & & A_K \end{bmatrix} \in \mathbb{M} \left(\sum_{i=1}^K n_i, \sum_{i=1}^K n_i \right)$$

is called block diagonal matrix.

Very often having information on the matrices A_i gives information on A .

Remark 76 It is easy, but cumbersome, to verify the following properties.

1. (associative) $\forall A \in \mathcal{M}_{m,n}, \forall B \in \mathcal{M}_{n,p}, \forall C \in \mathcal{M}_{p,q}, A(BC) = (AB)C$;
2. (distributive) $\forall A \in \mathcal{M}_{m,n}, \forall B \in \mathcal{M}_{m,n}, \forall C \in \mathcal{M}_{n,p}, (A+B)C = AC + BC$.
3. $\forall x, y \in \mathbb{R}^n$ and $\forall \alpha, \beta \in \mathbb{R}$,

$$A(\alpha x + \beta y) = A(\alpha x) + B(\beta y) = \alpha Ax + \beta Ay$$

It is false that:

1. (commutative) $\forall A \in \mathcal{M}_{m,n}, \forall B \in \mathcal{M}_{n,p}, AB = BA$;
2. $\forall A \in \mathcal{M}_{m,n}, \forall B, C \in \mathcal{M}_{n,p}, \langle A \neq 0, AB = AC \rangle \implies \langle B = C \rangle$;
3. $\forall A \in \mathcal{M}_{m,n}, \forall B \in \mathcal{M}_{n,p}, \langle A \neq 0, AB = 0 \rangle \implies \langle B = 0 \rangle$.

Let's show why the above statements are false.

1.

$$A = \begin{bmatrix} 1 & 2 & 1 \\ -1 & 1 & 3 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 0 & 1 \end{bmatrix}$$

$$AB = \begin{bmatrix} 1 & 2 & 1 \\ -1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 5 & 3 \\ 1 & 4 \end{bmatrix}$$

$$\begin{aligned}
BA &= \begin{bmatrix} 1 & 0 \\ 2 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 1 \\ -1 & 1 & 3 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 5 & 5 \\ -1 & 1 & 3 \end{bmatrix} \\
C &= \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix} \quad D = \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \\
CD &= \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} = \begin{bmatrix} 7 & 4 \\ 2 & 2 \end{bmatrix} \\
DC &= \begin{bmatrix} 1 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 2 \\ 1 & 8 \end{bmatrix}
\end{aligned}$$

Observe that since the commutative property does not hold true, we have to distinguish between “left factor out” and “right factor out” and also between “left multiplication or pre-multiplication” and “right multiplication or post-multiplication”:

$$\begin{aligned}
AB + AC &= A(B + C) \\
EF + GF &= (E + G)F \\
AB + CA &\neq A(B + C) \\
AB + CA &\neq (B + C)A
\end{aligned}$$

2.

Given

$$A = \begin{bmatrix} 3 & 1 \\ 6 & 2 \end{bmatrix} \quad B = \begin{bmatrix} 4 & 1 \\ -5 & 6 \end{bmatrix} \quad C = \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix},$$

we have

$$\begin{aligned}
AB &= \begin{bmatrix} 3 & 1 \\ 6 & 2 \end{bmatrix} \begin{bmatrix} 4 & 1 \\ -5 & 6 \end{bmatrix} = \begin{bmatrix} 7 & 9 \\ 14 & 18 \end{bmatrix} \\
AC &= \begin{bmatrix} 3 & 1 \\ 6 & 2 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} = \begin{bmatrix} 7 & 9 \\ 14 & 18 \end{bmatrix}
\end{aligned}$$

3.

Observe that $3. \Rightarrow 2.$ and therefore $\neg 2. \Rightarrow \neg 3.$ Otherwise, you can simply observe that 3. follows from 2., choosing A in 3. equal to A in 2., and B in 3. equal to $B - C$ in 2.:

$$A(B - C) = \begin{bmatrix} 3 & 1 \\ 6 & 2 \end{bmatrix} \cdot \left(\begin{bmatrix} 4 & 1 \\ -5 & 6 \end{bmatrix} - \begin{bmatrix} 1 & 2 \\ 4 & 3 \end{bmatrix} \right) = \begin{bmatrix} 3 & 1 \\ 6 & 2 \end{bmatrix} \cdot \begin{bmatrix} 3 & -1 \\ -9 & 3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$$

Since the associative property of the product between matrices does hold true we can give the following definition.

Definition 77 Given $A \in \mathcal{M}_{m,m}$,

$$A^k := \underset{1}{A} \cdot \underset{2}{A} \cdot \dots \cdot \underset{k \text{ times}}{A}.$$

Observe that if $A \in \mathcal{M}_{m,m}$ and $k, l \in \mathbb{N} \setminus \{0\}$, then

$$A^k \cdot A^l = A^{k+l}.$$

Remark 78 *Properties of transpose matrices.*

For any $m, n, p \in \mathbb{N}$,

1. $\forall A \in \mathcal{M}_{m,n}$ $(A^T)^T = A$
2. $\forall A, B \in \mathcal{M}_{m,n}$ $(A + B)^T = A^T + B^T$
3. $\forall \alpha \in \mathbb{R}, \forall A \in \mathcal{M}_{m,n}$ $(\alpha A)^T = \alpha A^T$
4. $\forall A \in \mathcal{M}_{m,n}, \forall B \in \mathcal{M}_{n,p}$ $(AB)^T = B^T A^T$

Matrices and linear systems.

In Section 1.6, we have seen that a system of m linear equation in the n unknowns x_1, x_2, \dots, x_n and parameters a_{ij} , for $i \in \{1, \dots, m\}$, $j \in \{1, \dots, n\}$, $(b_i)_{i=1}^n \in \mathbb{R}^n$ is displayed below:

$$\begin{cases} a_{11}x_1 + \dots + a_{1j}x_j + \dots + a_{1n}x_n = b_1 \\ \dots \\ a_{i1}x_1 + \dots + a_{ij}x_j + \dots + a_{in}x_n = b_i \\ \dots \\ a_{m1}x_1 + \dots + a_{mj}x_j + \dots + a_{mn}x_n = b_m \end{cases} \quad (3.8)$$

Moreover, the matrix

$$\begin{bmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} & b_1 \\ \dots & & & & & \\ a_{i1} & \dots & a_{ij} & \dots & a_{in} & b_i \\ \dots & & & & & \\ a_{m1} & \dots & a_{mj} & \dots & a_{mn} & b_m \end{bmatrix}$$

is called the augmented matrix M of system (1.4). The coefficient matrix A of the system is

$$A = \begin{bmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ \dots & & & & \\ a_{i1} & \dots & a_{ij} & \dots & a_{in} \\ \dots & & & & \\ a_{m1} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix}$$

Using the notations we described in the present section, we can rewrite linear equations and systems of linear equations in a convenient and short manner, as described below.

The linear equation in the unknowns x_1, \dots, x_n and parameters $a_1, \dots, a_i, \dots, a_n$, $b \in \mathbb{R}$

$$a_1x_1 + \dots + a_ix_i + \dots + a_nx_n = b$$

can be rewritten as

$$\sum_{i=1}^n a_ix_i = b$$

or

$$a \cdot x = b$$

where $a = [a_1, \dots, a_n]$ and $x = \begin{bmatrix} x_1 \\ \dots \\ x_n \end{bmatrix}$.

The linear system (3.8) can be rewritten as

$$\begin{cases} \sum_{j=1}^n a_{1j}x_j = b_1 \\ \dots \\ \sum_{j=1}^n a_{mj}x_j = b_m \end{cases}$$

or

$$\begin{cases} R^1(A)x = b_1 \\ \dots \\ R^m(A)x = b_m \end{cases}$$

or

$$Ax = b$$

where $A = [a_{ij}]$.

Definition 79 The trace of $A \in \mathcal{M}_{mm}$, written $\text{tr } A$, is the sum of the diagonal entries, i.e.,

$$\text{tr } A = \sum_{i=1}^m a_{ii}.$$

Definition 80 The identity matrix I_m is a diagonal matrix of order m with each element on the principal diagonal equal to 1. If no confusion arises, we simply write I in the place of I_m .

Remark 81 1. $\forall n \in \mathbb{N} \setminus \{0\}, (I_m)^n = I_m$;
2. $\forall A \in \mathcal{M}_{m,n}, I_m A = A I_n = A$.

Proposition 82 Let $A, B \in \mathcal{M}(m, m)$ and $k \in \mathbb{R}$. Then

1. $\text{tr}(A + B) = \text{tr } A + \text{tr } B$;
2. $\text{tr } kA = k \cdot \text{tr } A$;
3. $\text{tr } AB = \text{tr } BA$.

Proof. Exercise. ■

3.2 Inverse matrices

Definition 83 Given a matrix $A_{n \times n}$, a matrix $B_{n \times n}$ is called an inverse of A if

$$AB = BA = I_n.$$

We then say that A is invertible, or that A admits an inverse.

Proposition 84 If A admits an inverse, then the inverse is unique.

Proof. Let the inverse matrices B and C of A be given. Then

$$AB = BA = I_n \tag{3.9}$$

and

$$AC = CA = I_n \tag{3.10}$$

Left multiplying the first two terms in the equality (3.9) by C , we get

$$(CA)B = C(BA)$$

and from (3.10) and (3.9) we get $B = C$, as desired. ■

Thanks to the above Proposition, we can present the following definition.

Definition 85 If the inverse of A does exist, then it is denoted by A^{-1} .

Example 86 Assume that for $i \in \{1, \dots, n\}$, $\lambda_i \neq 0$. The diagonal matrix

$$\begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}$$

is invertible and its inverse is

$$\begin{bmatrix} \frac{1}{\lambda_1} & & \\ & \ddots & \\ & & \frac{1}{\lambda_n} \end{bmatrix}.$$

Remark 87 If a row or a column of A is zero, then A is not invertible, as verified below.

Without loss of generality, assume the first row of A is equal to zero. Assume that B is the inverse of A . But then, since $I = AB$, we would have $1 = R^1(A) \cdot C^1(B) = 0$, a contradiction.

Proposition 88 If $A \in \mathcal{M}_{m,m}$ and $B \in \mathcal{M}_{m,m}$ are invertible matrices, then AB is invertible and

$$(AB)^{-1} = B^{-1}A^{-1}$$

Proof.

$$(AB)B^{-1}A^{-1} = A(BB^{-1})A^{-1} = AIA^{-1} = AA^{-1} = I.$$

$$B^{-1}A^{-1}(AB) = B^{-1}(A^{-1}A)B = B^{-1}IB = B^{-1}B = I.$$

■

Remark 89 The existence of the inverse matrix gives an obvious way of solving systems of linear equations with the same number of equations and unknowns.

Given the system

$$A_{n \times n}x = b,$$

if A^{-1} exists, then

$$x = A^{-1}b.$$

Proposition 90 (Some other properties of the inverse matrix)

Let the invertible matrix A be given.

1. A^{-1} is invertible and $(A^{-1})^{-1} = A$;

2. A^T is invertible and $(A^T)^{-1} = (A^{-1})^T$;

Proof. 1. We want to verify that the inverse of A^{-1} is A , i.e.,

$$A^{-1}A = I \text{ and } AA^{-1} = I,$$

which is obvious.

2. Observe that

$$A^T \cdot (A^{-1})^T = (A^{-1} \cdot A)^T = I^T = I,$$

and

$$(A^{-1})^T \cdot A^T = (A \cdot A^{-1})^T = I.$$

■

3.3 Elementary matrices

Below, we recall the definition of elementary row operations on a matrix $A \in M_{m \times n}$ presented in Definition 34.

Definition 91 An elementary row operation on a matrix $A \in M_{m \times n}$ is one of the following operations on the rows of A :

$[\mathcal{E}_1]$ (Row interchange) Interchange R^i with R^j , denoted by $R^i \leftrightarrow R^j$;

$[\mathcal{E}_2]$ (Row scaling) Multiply R^i by $k \in \mathbb{R} \setminus \{0\}$, denoted by $kR^i \rightarrow R^i$, $k \neq 0$;

$[\mathcal{E}_3]$ (Row addition) Replace R^i by (k times R^j plus R^i), denoted by $(R^i + kR^j) \rightarrow R^i$.

Sometimes we apply $[\mathcal{E}_2]$ and $[\mathcal{E}_3]$ in one step, i.e., we perform the following operation

$[\mathcal{E}']$ Replace R^i by (k' times R^j and $k \in \mathbb{R} \setminus \{0\}$ times R^i), denoted by $(k'R^j + kR^i) \rightarrow R^i$, $k \neq 0$.

Definition 92 Let \mathfrak{E} be the set of functions $\mathcal{E} : \mathcal{M}_{m,n} \rightarrow \mathcal{M}_{m,n}$ which associate with any matrix $A \in \mathcal{M}_{m,n}$ a matrix $\mathcal{E}(A)$ obtained from A via an elementary row operation presented in Definition 91. For $i \in \{1, 2, 3\}$, let $\mathfrak{E}_i \subseteq \mathfrak{E}$ be the set of elementary row operation functions of type i presented in Definition 91.

Definition 93 For any $\mathcal{E} \in \mathfrak{E}$, define

$$E_{\mathcal{E}} = \mathcal{E}(I_m) \in \mathcal{M}_{m,m}.$$

$E_{\mathcal{E}}$ is called the elementary matrix corresponding to the elementary row operation function \mathcal{E} .

With some abuse of terminology, we call any $\mathcal{E} \in \mathfrak{E}$ an elementary row operation (omitting the word “function”), and we sometimes omit the subscript \mathcal{E} .

Proposition 94 *Each elementary row operations $\mathcal{E}_1, \mathcal{E}_2$ and \mathcal{E}_3 has an inverse, and that inverse is of the same type, i.e., for $i \in \{1, 2, 3\}$, $\mathcal{E} \in \mathfrak{E}_i \Leftrightarrow \mathcal{E}^{-1} \in \mathfrak{E}_i$.*

- Proof.** 1. The inverse of $R^i \leftrightarrow R^j$ is $R^j \leftrightarrow R^i$.
 2. The inverse of $kR^i \rightarrow R^i$, $k \neq 0$ is $k^{-1}R^i \rightarrow R^i$.
 3. The inverse of $(R^i + kR^j) \rightarrow R^i$ is $(-kR^j + R^i) \rightarrow R^i$. ■

Remark 95 *Given the row, canonical¹ vectors e_m^i , for $i \in \{1, \dots, m\}$,*

$$I_m = \begin{bmatrix} e_m^1 \\ \dots \\ e_m^i \\ \dots \\ e_m^j \\ \dots \\ e_m^n \end{bmatrix}$$

The following Proposition shows that the result of applying an elementary row operation \mathcal{E} to a matrix A can be obtained by premultiplying A by the corresponding elementary matrix $E_{\mathcal{E}}$.

Proposition 96 *For any $A \in M_{m,n}$ and for any $\mathcal{E} \in \mathfrak{E}$,*

$$\mathcal{E}(A) = \mathcal{E}(I_m) \cdot A := E_{\mathcal{E}}A. \quad (3.11)$$

Proof. Recall that

$$A = \begin{bmatrix} R^1(A) \\ \dots \\ R^i(A) \\ \dots \\ R^j(A) \\ \dots \\ R^m(A) \end{bmatrix}$$

We have to prove that (3.11) does hold true $\forall \mathcal{E} \in \{\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3\}$.

1. $\mathcal{E} \in \mathfrak{E}_1$.

First of all observe that

$$\mathcal{E}(I_m) = \begin{bmatrix} e_m^1 \\ \dots \\ e_m^j \\ \dots \\ e_m^i \\ \dots \\ e_m^m \end{bmatrix} \quad \text{and} \quad \mathcal{E}(A) = \begin{bmatrix} R^1(A) \\ \dots \\ R^j(A) \\ \dots \\ R^i(A) \\ \dots \\ R^m(A) \end{bmatrix}.$$

From (3.6),

$$\mathcal{E}(I_m) \cdot A = \begin{bmatrix} e_m^1 \cdot A \\ \dots \\ e_m^j \cdot A \\ \dots \\ e_m^i \cdot A \\ \dots \\ e_m^m \cdot A \end{bmatrix} = \begin{bmatrix} R^1(A) \\ \dots \\ R^j(A) \\ \dots \\ R^i(A) \\ \dots \\ R^m(A) \end{bmatrix},$$

as desired.

2. $\mathcal{E} \in \mathfrak{E}_2$.

¹See Definition 55.

Observe that

$$\mathcal{E}(I_m) = \begin{bmatrix} e_m^1 \\ \dots \\ k \cdot e_m^i \\ \dots \\ e_m^j \\ \dots \\ e_m^m \end{bmatrix} \quad \text{and} \quad \mathcal{E}(A) = \begin{bmatrix} R^1(A) \\ \dots \\ k \cdot R^j(A) \\ \dots \\ R^i(A) \\ \dots \\ R^m(A) \end{bmatrix}.$$

$$\mathcal{E}(I_m) \cdot A = \begin{bmatrix} e_m^1 \cdot A \\ \dots \\ k \cdot e_m^i \cdot A \\ \dots \\ e_m^j \cdot A \\ \dots \\ e_m^m \cdot A \end{bmatrix} = \begin{bmatrix} R^1(A) \\ \dots \\ k \cdot R^j(A) \\ \dots \\ R^i(A) \\ \dots \\ R^m(A) \end{bmatrix},$$

as desired.

3. $\mathcal{E} \in \mathfrak{E}_3$.

Observe that

$$\mathcal{E}(I_m) = \begin{bmatrix} e_m^1 \\ \dots \\ e_m^i + k \cdot e_m^j \\ \dots \\ e_m^j \\ \dots \\ e_m^m \end{bmatrix} \quad \text{and} \quad \mathcal{E}(A) = \begin{bmatrix} R^1(A) \\ \dots \\ R^i(A) + k \cdot R^j(A) \\ \dots \\ R^j(A) \\ \dots \\ R^m(A) \end{bmatrix}.$$

$$\mathcal{E}(I_m) \cdot A = \begin{bmatrix} e_m^1 \\ \dots \\ e_m^i + k \cdot e_m^j \\ \dots \\ e_m^j \\ \dots \\ e_m^m \end{bmatrix} A = \begin{bmatrix} e_m^1 \cdot A \\ \dots \\ (e_m^i + k \cdot e_m^j) \cdot A \\ \dots \\ e_m^j \cdot A \\ \dots \\ e_m^m \cdot A \end{bmatrix} = \begin{bmatrix} R^1(A) \\ \dots \\ R^i(A) + k \cdot R^j(A) \\ \dots \\ R^j(A) \\ \dots \\ R^m(A) \end{bmatrix},$$

as desired. ■

Corollary 97 *If A is row equivalent to B , then there exist $k \in \mathbb{N}$ and elementary matrices E_1, \dots, E_k such that*

$$B = E_1 \cdot E_2 \cdot \dots \cdot E_k \cdot A$$

Proof. It follows from the definition of row equivalence and Proposition 96. ■

Proposition 98 *Every elementary matrix $E_{\mathcal{E}}$ is invertible and $(E_{\mathcal{E}})^{-1}$ is an elementary matrix. In fact, $(E_{\mathcal{E}})^{-1} = E_{\mathcal{E}^{-1}}$.*

Proof. Given an elementary matrix E , from Definition 93, $\exists \mathcal{E} \in \mathfrak{E}$ such that

$$E = \mathcal{E}(I) \tag{3.12}$$

Define

$$E' = \mathcal{E}^{-1}(I).$$

Then

$$I \stackrel{\text{def. inv. func.}}{=} \mathcal{E}^{-1}(\mathcal{E}(I)) \stackrel{(3.12)}{=} \mathcal{E}^{-1}(E) \stackrel{\text{Prop. (96)}}{=} \mathcal{E}^{-1}(I) \cdot E \stackrel{\text{def. } E'}{=} E' E$$

and

$$I \stackrel{\text{def. inv.}}{=} \mathcal{E}(\mathcal{E}^{-1}(I)) \stackrel{\text{def. } E'}{=} \mathcal{E}(E') \stackrel{\text{Prop. (96)}}{=} \mathcal{E}(I) \cdot E' \stackrel{(3.12)}{=} E E'.$$

■

Corollary 99 *If E_1, \dots, E_k are elementary matrices, then*

$$P := E_1 \cdot E_2 \cdot \dots \cdot E_k$$

is an invertible matrix.

Proof. It follows from Proposition 88 and Proposition 98. In fact, $(E_k^{-1} \cdot \dots \cdot E_2^{-1} \cdot E_1^{-1})$ is the inverse of P . ■

Proposition 100 *Let $A \in M_{m \times n}$ be given. Then, there exist a matrix $B \in M_{m \times n}$ in row canonical form, $k \in \mathbb{N}$ and elementary matrices E_1, \dots, E_k such that*

$$B = E_1 \cdot E_2 \cdot \dots \cdot E_k \cdot A.$$

Proof. From Proposition 38, there exist $k \in \mathbb{N}$ elementary operations $\mathcal{E}^1, \dots, \mathcal{E}^k$ such that

$$(\mathcal{E}^1 \circ \mathcal{E}^2 \circ \dots \circ \mathcal{E}^k)(A) = B.$$

From Proposition 96, $\forall j \in \{1, \dots, k\}$,

$$\mathcal{E}^j(M) = \mathcal{E}^j(I) \cdot M := E_j \cdot M.$$

Then,

$$\begin{aligned} (\mathcal{E}^1 \circ \mathcal{E}^2 \circ \dots \circ \mathcal{E}^k)(A) &= (\mathcal{E}^1 \circ \mathcal{E}^2 \circ \dots \circ \mathcal{E}^{k-1})(\mathcal{E}^k(A)) = (\mathcal{E}^1 \circ \mathcal{E}^2 \circ \dots \circ \mathcal{E}^{k-1})(E_k \cdot A) = \\ &= (\mathcal{E}^1 \circ \mathcal{E}^2 \circ \dots \circ \mathcal{E}^{k-2}) \circ \mathcal{E}^{k-1}(E_k \cdot A) = (\mathcal{E}^1 \circ \mathcal{E}^2 \circ \dots \circ \mathcal{E}^{k-2})(E_{k-1} \cdot E_k \cdot A) = \\ &= (\mathcal{E}^1 \circ \mathcal{E}^2 \circ \dots \circ \mathcal{E}^{k-3}) \circ \mathcal{E}^{k-2}(E_{k-1} \cdot E_k \cdot A) = (\mathcal{E}^1 \circ \mathcal{E}^2 \circ \dots \circ \mathcal{E}^{k-3})(E_{k-2} \cdot E_{k-1} \cdot E_k \cdot A) = \\ &\dots = E_1 \cdot E_2 \cdot \dots \cdot E_k \cdot A, \end{aligned}$$

as desired. ■

Remark 101 *In fact, in Proposition 159, we will show that the matrix B of the above Corollary is unique.*

Proposition 102 *To be row equivalent is an equivalence relation.*

Proof. Obvious. ■

Proposition 103 $\forall n \in \mathbb{N} \setminus \{0\}$, $A_{n \times n}$ is in row canonical form and it is invertible $\Leftrightarrow A = I$.

Proof. [\Leftarrow] Obvious.

[\Rightarrow]

We proceed by induction on n .

Case 1. $n = 1$.

The case $n = 1$ is obvious. To try to better understand the logic of the proof, take $n = 2$, i.e., suppose that

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

is in row canonical form and invertible. Observe that $A \neq 0$.

1. $a_{11} = 1$. Suppose $a_{11} = 0$. Then, from 1. in the definition of matrix in echelon form - see Definition 28 - $a_{12} \neq 0$ (otherwise, you would have a zero row not on the bottom of the matrix). Then, from 2. in that definition, we must have $a_{21} = 0$. But then the first column is zero, contradicting the fact that A is invertible - see Remark 87. Since $a_{11} \neq 0$, then from 2. in the Definition of row canonical form matrix - see Definition 31 - we get $a_{11} = 1$.

2. $a_{21} = 0$. It follows from the fact that $a_{11} = 1$ and 3. in Definition 31.

3. $a_{22} = 1$. Suppose $a_{22} = 0$, but then the last row would be zero, contradicting the fact that A is invertible and a_{22} is the leading nonzero entry of the second row, i.e., $a_{22} \neq 0$. Then from 2. in the Definition of row canonical form matrix, we get $a_{22} = 1$.

4. $a_{12} = 0$. It follows from the fact that $a_{22} = 1$ and 3. in Definition 31.

Case 2. Assume that statement is true for $n - 1$.

Suppose that

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \dots & & & & & \\ a_{i1} & a_{i2} & \dots & a_{ij} & \dots & a_{in} \\ \dots & & & & & \\ a_{n1} & a_{n2} & \dots & a_{nj} & \dots & a_{nn} \end{bmatrix}$$

is in row canonical form and invertible.

1. $a_{11} = 1$. Suppose $a_{11} = 0$. Then, from 1. in the definition of matrix in echelon form - see Definition 28 -

$$(a_{12}, \dots, a_{1n}) \neq 0.$$

Then, from 2. in that definition, we must have

$$\begin{bmatrix} a_{21} \\ \dots \\ a_{i1} \\ \dots \\ a_{n1} \end{bmatrix} = 0.$$

But then the first column is zero, contradicting the fact that A is invertible - see Remark 87. Since $a_{11} \neq 0$, then from 2. in the Definition of row canonical form matrix - see Definition 31 - we get $a_{11} = 1$.

2. Therefore, we can rewrite the matrix as follows

$$A = \begin{bmatrix} 1 & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ 0 & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \dots & & & & & \\ 0 & a_{i2} & \dots & a_{ij} & \dots & a_{in} \\ \dots & & & & & \\ 0 & a_{n2} & \dots & a_{nj} & \dots & a_{nn} \end{bmatrix} = \begin{bmatrix} 1 & a \\ 0 & A_{22} \end{bmatrix} \quad (3.13)$$

with obvious definitions of a and A_{22} . Since, by assumption, A is invertible, there exists B which we can partition in the same we partitioned A , i.e.,

$$B = \begin{bmatrix} b_{11} & b \\ c & B_{22} \end{bmatrix}.$$

and such that B is invertible. Then,

$$I_n = BA = \begin{bmatrix} b_{11} & b \\ c & B_{22} \end{bmatrix} \begin{bmatrix} 1 & a \\ 0 & A_{22} \end{bmatrix} = \begin{bmatrix} b_{11} & b_{11} + bA_{22} \\ c & ca + B_{22}A_{22} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & I_{n-1} \end{bmatrix};$$

then $c = 0$ and $A_{22}B_{22} = I_{n-1}$.

Moreover,

$$I_n = AB = \begin{bmatrix} 1 & a \\ 0 & A_{22} \end{bmatrix} \begin{bmatrix} b_{11} & b \\ c & B_{22} \end{bmatrix} = \begin{bmatrix} b_{11} + ac & b + aB_{22} \\ c & A_{22}B_{22} \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & I_{n-1} \end{bmatrix}. \quad (3.14)$$

Therefore, A_{22} is invertible. From 3.13, A_{22} can be obtained from A erasing the first row and then erasing a column of zero, from Remark 33, A_{22} is a row reduced form matrix. Then, we can apply the assumption of the induction argument to conclude that $A_{22} = I_{n-1}$. Then, from 3.13,

$$A = \begin{bmatrix} 1 & a \\ 0 & I \end{bmatrix}.$$

Since, by assumption, $A_{n \times n}$ is in row canonical form, from 3. in Definition 31, $a = 0$, and, as desired $A = I$. ■

Proposition 104 *Let A belong to $M_{m,m}$. Then the following statements are equivalent.*

1. A is invertible;
2. A is row equivalent to I_m ;
3. A is the product of elementary matrices.

Proof. 1. \Rightarrow 2.

From Proposition 100, there exist a matrix $B \in M_{m \times m}$ in row canonical form, $k \in \mathbb{N}$ and elementary matrices E_1, \dots, E_k such that

$$B = E_1 \cdot E_2 \cdot \dots \cdot E_k \cdot A.$$

Since A is invertible and, from Corollary 99, $E_1 \cdot E_2 \cdot \dots \cdot E_k$ is invertible as well, from Proposition 88, B is invertible as well. Then, from Proposition 103, $B = I$.

2. \Rightarrow 3.

By assumption and from Corollary 97, there exist $k \in \mathbb{N}$ and elementary matrices E_1, \dots, E_k such that

$$A = E_1 \cdot E_2 \cdot \dots \cdot E_k \cdot I,$$

Since $\forall i \in \{1, \dots, k\}$, E_i is an elementary matrix, the desired result follows.

3. \Rightarrow 1.

By assumption, there exist $k \in \mathbb{N}$ and elementary matrices E_1, \dots, E_k such that

$$A = E_1 \cdot E_2 \cdot \dots \cdot E_k.$$

Since, from Proposition 98, $\forall i \in \{1, \dots, k\}$, E_i is invertible, A is invertible as well, from Proposition 88. ■

Proposition 105 *Let $A_{m \times n}$ be given.*

1. $B_{m \times n}$ is row equivalent to $A_{m \times n} \Leftrightarrow$ there exists an invertible $P_{m \times m}$ such that $B = PA$.
2. $P_{m \times m}$ is an invertible matrix $\Rightarrow PA$ is row equivalent to A .

Proof. 1.

[\Rightarrow] From Corollaries 99 and 97, $B = E_1 \cdot \dots \cdot E_k \cdot A$ with $(E_1 \cdot \dots \cdot E_k)$ invertible matrix. Then, it suffices to take $P = E_1 \cdot \dots \cdot E_k$.

[\Leftarrow] From Proposition 104, P is row equivalent to I , i.e., there exist E_1, \dots, E_k such that $P = E_1 \cdot \dots \cdot E_k \cdot I$. Then by assumption $B = E_1 \cdot \dots \cdot E_k \cdot I \cdot A$, i.e., B is row equivalent to A .

2.

From Proposition 104, P is the product of elementary matrices. Then, the desired result follows from Proposition 96. ■

Proposition 106 *If A is row equivalent to a matrix with a zero row, then A is not invertible.*

Proof. Suppose otherwise, i.e., A is row equivalent to a matrix C with a zero row and A is invertible. From Proposition 105, there exists an invertible P such that $A = PC$ and then $P^{-1}A = C$. Since A and P^{-1} are invertible, then, from Proposition 88, $P^{-1}A$ is invertible, while C , from Remark 87, C is not invertible, a contradiction. ■

Remark 107 *From Proposition 104, we know that if $A_{m \times m}$ is invertible, then there exist E_1, \dots, E_k such that*

$$I = E_1 \cdot \dots \cdot E_k \cdot A \tag{3.15}$$

or

$$A^{-1} = E_1 \cdot \dots \cdot E_k \cdot I. \tag{3.16}$$

Then, from (3.15) and (3.16), if A is invertible then A^{-1} is equal to the finite product of those elementary matrices which “transform” A in I , or, equivalently, can be obtained applying a finite number of corresponding elementary operations to the identity matrix I . That observation leads to the following (Gaussian elimination) algorithm, which either show that an arbitrary matrix $A_{m \times m}$ is not invertible or finds the inverse of A .

An algorithm to find the inverse of a matrix $A_{m \times m}$ or to show the matrix is not invertible.

Step 1. Construct the following matrix $M_{m \times (2m)}$:

$$\begin{bmatrix} A & I_m \end{bmatrix}$$

Step 2. Row reduce M to echelon form. If the process generates a zero row in the part of M corresponding to A , then stop: A is not invertible : A is row equivalent to a matrix with a zero row and therefore, from Proposition 106 is not invertible. Otherwise, the part of M corresponding to A is a triangular matrix.

Step 3. Row reduce M to the row canonical form

$$\begin{bmatrix} I_m & B \end{bmatrix}$$

Then, from Remark 107, $A^{-1} = B$.

Example 108 We find the inverse of

$$A = \begin{bmatrix} 1 & 0 & 2 \\ 2 & -1 & 3 \\ 4 & 1 & 8 \end{bmatrix},$$

applying the above algorithm.

Step 1.

$$M = \begin{bmatrix} 1 & 0 & 2 & 1 & 0 & 0 \\ 2 & -1 & 3 & 0 & 1 & 0 \\ 4 & 1 & 8 & 0 & 0 & 1 \end{bmatrix}.$$

Step 2.

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & -1 & -1 & -2 & 1 & 0 \\ 0 & 1 & 0 & -4 & 0 & 1 \end{array} \right], \left[\begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & -1 & -1 & -2 & 1 & 0 \\ 0 & 0 & -1 & -6 & 1 & 1 \end{array} \right]$$

The matrix is invertible.

Step 3.

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & -1 & -1 & -2 & 1 & 0 \\ 0 & 0 & -1 & -6 & 1 & 1 \end{array} \right], \left[\begin{array}{ccc|ccc} 1 & 0 & 2 & 1 & 0 & 0 \\ 0 & -1 & -1 & -2 & 1 & 0 \\ 0 & 0 & 1 & 6 & -1 & -1 \end{array} \right],$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -11 & 2 & 2 \\ 0 & -1 & 0 & 4 & 0 & -1 \\ 0 & 0 & 1 & 6 & -1 & -1 \end{array} \right], \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & -11 & 2 & 2 \\ 0 & 1 & 0 & -4 & 0 & 1 \\ 0 & 0 & 1 & 6 & -1 & -1 \end{array} \right]$$

Then

$$A^{-1} = \begin{bmatrix} -11 & 2 & 2 \\ -4 & 0 & 1 \\ 6 & -1 & -1 \end{bmatrix}.$$

Example 109

$$\left[\begin{array}{cc|cc} 1 & 3 & 1 & 0 \\ 4 & 2 & 0 & 1 \end{array} \right]$$

$$\left[\begin{array}{cc|cc} 1 & 3 & 1 & 0 \\ 0 & -10 & -4 & 1 \end{array} \right]$$

$$\left[\begin{array}{cc|cc} 1 & 3 & 1 & 0 \\ 0 & 1 & \frac{4}{10} & -\frac{1}{10} \end{array} \right]$$

$$\left[\begin{array}{cc|cc} 1 & 0 & -\frac{2}{10} & \frac{0}{3} \\ 0 & 1 & \frac{4}{10} & -\frac{1}{10} \end{array} \right]$$

3.4 Elementary column operations

This section repeats some of the discussion of the previous section using column instead of rows of a matrix.

Definition 110 *An elementary column operation is one of the following operations on the columns of $A_{m \times n}$:*

[\mathcal{F}_1] (Column interchange) Interchange C_i with C_j , denoted by $C_i \leftrightarrow C_j$;

[\mathcal{F}_2] (Column scaling) Multiply C_i by $k \in \mathbb{R} \setminus \{0\}$, denoted by $kC_i \rightarrow C_i$, $k \neq 0$;

[\mathcal{F}_3] (Column addition) Replace C_i by (k times C_j plus C_i), denoted by $(C_i + kC_j) \rightarrow C_i$.

Each of the above column operation has an inverse operation of the same type just like the corresponding row operations.

Definition 111 *Let \mathcal{F} be an elementary column operation on a matrix $A_{m \times n}$. We denote the resulting matrix by $\mathcal{F}(A)$. We define also*

$$F_{\mathcal{F}} = \mathcal{F}(I_n) \in \mathcal{M}_{n,n}.$$

$F_{\mathcal{F}}$ is then called an elementary matrix corresponding to the elementary column operation \mathcal{F} . We sometimes omit the subscript \mathcal{F} .

Definition 112 *Given an elementary row operation \mathcal{E} , define $\mathcal{F}_{\mathcal{E}}$, if it exists², as the column operation obtained by \mathcal{E} substituting the word row with the word column. Similarly, given an elementary column operation \mathcal{F} define $\mathcal{E}_{\mathcal{F}}$, if it exists, as the row operation obtained by \mathcal{F} substituting the word column with the word row.*

In what follows, \mathcal{F} and \mathcal{E} are such that $\mathcal{F} = \mathcal{F}_{\mathcal{E}}$ and $\mathcal{E}_{\mathcal{F}} = \mathcal{E}$.

Proposition 113 *Let a matrix $A_{m \times n}$ be given. Then*

$$\mathcal{F}(A) = [\mathcal{E}(A^T)]^T.$$

Proof. The above fact is equivalent to $\mathcal{E}(A^T) = (\mathcal{F}(A))^T$ and it is a consequence of the fact that the columns of A are the rows of A^T and vice versa. As an exercise, carefully do the proof in the case of each of the three elementary operation types. ■

Remark 114 *The above Proposition says that applying the column operation \mathcal{F} to a matrix A gives the same result as applying the corresponding row operation $\mathcal{E}_{\mathcal{F}}$ to A^T and then taking the transpose.*

Proposition 115 *Let a matrix $A_{m \times n}$ be given. Then*

1.

$$\mathcal{F}(A) = A \cdot (\mathcal{E}(I))^T = A \cdot \mathcal{F}(I),$$

or, since $E := \mathcal{E}(I)$ and $F := \mathcal{F}(I)$,

$$\mathcal{F}(A) = A \cdot E^T = A \cdot F. \quad (3.17)$$

2. $F = E^T$ and F is invertible.

Proof. 1.

$$\mathcal{F}(A) \stackrel{\text{Lemma 113}}{=} [\mathcal{E}(A^T)]^T \stackrel{\text{Lemma 96}}{=} (\mathcal{E}(I) \cdot A^T)^T = A \cdot (\mathcal{E}(I))^T \stackrel{\text{Lemma 113}}{=} A \cdot \mathcal{F}(I).$$

2. From (3.17), we then get

$$F := \mathcal{F}(I) = I \cdot E^T = E^T.$$

From Proposition 90 and Proposition 98, it follows that F is invertible. ■

²Of course, if you exchange the first and the third row, and the matrix has only two columns, you cannot exchange the first and the third column.

Remark 116 The above Proposition says that says that the result of applying an elementary column operation \mathcal{F} on a matrix A can be obtained by postmultiplying A by the corresponding elementary matrix F .

Definition 117 A matrix $B_{m \times n}$ is said column equivalent to a matrix $A_{m \times n}$ if B can be obtained from A using a finite number of elementary column operations.

Remark 118 By definition of row equivalent, column equivalent and transpose of a matrix, we have that

$$A \text{ and } B \text{ are row equivalent} \Leftrightarrow A^T \text{ and } B^T \text{ are column equivalent,}$$

and

$$A \text{ and } B \text{ are column equivalent} \Leftrightarrow A^T \text{ and } B^T \text{ are row equivalent.}$$

Proposition 119 1. $B_{m \times n}$ is column equivalent to $A_{m \times n} \Leftrightarrow$ there exists an invertible $Q_{n \times n}$ such that $B_{m \times n} = A_{m \times n} Q_{n \times n}$.

2. $Q_{n \times n}$ in invertible matrix $\Rightarrow AQ$ is column equivalent to A .

Proof. It is very similar to the proof of Proposition 105. ■

Definition 120 A matrix $B_{m \times n}$ is said equivalent to a matrix $A_{m \times n}$ if B can be obtained from A using a finite number of elementary row and column operations.

Proposition 121 A matrix $B_{m \times n}$ is equivalent to a matrix $A_{m \times n} \Leftrightarrow$ there exist invertible matrices $P_{m \times m}$ and $Q_{n \times n}$ such that $B_{m \times n} = P_{m \times m} A_{m \times n} Q_{n \times n}$.

Proof. $[\Rightarrow]$

By assumption $B = E_1 \cdot \dots \cdot E_k \cdot A \cdot F_1 \cdot \dots \cdot F_h$.

$[\Leftarrow]$

Similar to the proof of Proposition 105. ■

Proposition 122 For any matrix $A_{m \times n}$ there exists a number $r \in \{0, 1, \dots, \min\{m, n\}\}$ such that A is equivalent to the block matrix of the form

$$\begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}. \quad (3.18)$$

Proof. The proof is constructive in the form of an algorithm.

Step 1. Row reduce A to row canonical form, with leading nonzero entries $a_{11}, a_{2j_2}, \dots, a_{j_r j_r}$.

Step 2. Interchange C^2 and C_{j_2} , C^3 and C_{j_3} and so on up to C_r and C_{j_r} . You then get a matrix of the form

$$\begin{bmatrix} I_r & B \\ 0 & 0 \end{bmatrix}.$$

Step 3. Use column operations to replace entries in B with zeros.

■

Remark 123 From Proposition 159 the matrix in Step 2 is unique and therefore the resulting matrix in Step 3, i.e., matrix (3.18) is unique.

Proposition 124 For any $A \in M_{m,n}$, there exists invertible matrices $P \in M_{m,m}$ and $Q \in M_{n,n}$ and $r \in \{0, 1, \dots, \min\{m, n\}\}$ such that

$$PAQ = \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}$$

Proof. It follows immediately from Propositions 122 and 121. ■

Remark 125 From Proposition 159 the number r in the statement of the previous Proposition is unique.

Example 126 *Take*

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 5 & 7 & 9 \end{bmatrix}$$

Then

$$\begin{aligned} & \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 5 & 7 & 9 \end{bmatrix} \xrightarrow{R^3 - (R^1 + R^2) \rightarrow R^3} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 0 & 0 & 0 \end{bmatrix} \longrightarrow \\ & \xrightarrow{R^2 - 4R^1 \rightarrow R^2} \begin{bmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{-\frac{1}{3}R^2 \rightarrow R^2} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \longrightarrow \\ & \xrightarrow{C^2 - 2C^1 \rightarrow C^2} \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{C^3 - 3C^1 \rightarrow C^3} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \xrightarrow{C^3 - 2C^2 \rightarrow C^3} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

We can find matrices P and Q as follows.

$$\begin{aligned} & \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 5 & 7 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 0 & 0 & 0 \end{bmatrix} \\ & \begin{bmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & 0 & 0 \end{bmatrix} \\ & \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{3} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 0 & -3 & -6 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

Then

$$P = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -\frac{1}{3} & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ -4 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -1 & -1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ \frac{4}{3} & -\frac{1}{3} & 0 \\ -1 & -1 & 1 \end{bmatrix};$$

indeed

$$\begin{aligned} & \begin{bmatrix} 1 & 0 & 0 \\ \frac{4}{3} & -\frac{1}{3} & 0 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 5 & 7 & 9 \end{bmatrix} = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix}. \\ & \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \\ & \begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & -3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \end{aligned}$$

d

Then

$$Q = \begin{bmatrix} 1 & -2 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & -3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix};$$

indeed,

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Summarizing

$$PAQ = \begin{bmatrix} 1 & 0 & 0 \\ \frac{4}{3} & -\frac{1}{3} & 0 \\ -1 & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 5 & 7 & 9 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ 0 & 1 & -2 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

Proposition 127 *If $A_{m \times m} B_{m \times m} = I$, then $BA = I$ and therefore A is invertible and $A^{-1} = B$.*

Proof. Suppose A is not invertible, then from Proposition 104 is not row equivalent to I_m and from Proposition 122, A is equivalent to a block matrix of the form displayed in (3.18) with $r < m$. Then, from Proposition 121, there exist invertible matrices $P_{m \times m}$ and $Q_{m \times m}$ such that

$$PAQ = \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix},$$

and from $AB = I$, we get

$$P = PAQQ^{-1}B$$

and

$$\begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} (Q^{-1}B) = P$$

Therefore, P has some zero rows and columns, contradicting that P is invertible. ■

Remark 128 *The previous Proposition says that to verify that A is invertible it is enough to check that $AB = I$.*

Remark 129 *We will come back to the analysis of further properties of the inverse and on another way of computing it in Section 5.3.*

3.5 Exercises

From Lipschutz (1991),

- starting from page 81: 3.1 - 3.11, 3.14 - 3.16;
- starting from page 111: 4.1, 4.4, 4.5, 4.7, 4.8.

Chapter 4

Vector spaces

4.1 Definition

Definition 130 Let a nonempty set F with the operations of addition which assigns to any $x, y \in F$ an element denoted by $x \oplus y \in F$, and multiplication which assigns to any $x, y \in F$ an element denoted by $x \odot y \in F$ be given. (F, \oplus, \odot) is called a field, if the following properties hold true.

1. (Commutative) $\forall x, y \in F, x \oplus y = y \oplus x$ and $x \odot y = y \odot x$;
2. (Associative) $\forall x, y, z \in F, (x \oplus y) \oplus z = x \oplus (y \oplus z)$ and $(x \odot y) \odot z = x \odot (y \odot z)$;
3. (Distributive) $\forall x, y, z \in F, x \odot (y \oplus z) = (x \odot y) \oplus (x \odot z)$;
4. (Existence of null elements) $\exists f_0, f_1 \in F$ such that $\forall x \in F, f_0 \oplus x = x$ and $f_1 \odot x = x$;
5. (Existence of a negative element) $\forall x \in F \exists y \in F$ such that $x \oplus y = f_0$;
From the above properties, it follows that f_0 and f_1 are unique.¹ We denote f_0 and f_1 by 0 and 1, respectively.
6. (Existence of an inverse element) $\forall x \in F \setminus \{0\}, \exists y \in F$ such that $x \odot y = 1$.
Elements of a field are called scalars.

Example 131 The set \mathbb{R} of real numbers with the standard addition and multiplication is a field. From the above properties all the rules of “elementary” algebra can be deduced.² The set \mathbb{C} of complex numbers is a field.

The sets $\mathbb{N} := \{1, 2, 3, \dots, n, \dots\}$ and \mathbb{Z} of positive integers and integers, respectively, with the standard addition and multiplication are not fields.

Definition 132 Let (F, \oplus, \odot) be a field and V be a nonempty set with the operations of addition which assigns to any $u, v \in V$ an element denoted by $u + v \in V$, and scalar multiplication which assigns to any $u \in V$ and any $\alpha \in F$ an element $\alpha \cdot u \in V$. Then $(V, +, \cdot)$ is called a vector space on the field (F, \oplus, \odot) and its elements are called vectors if the following properties are satisfied.

- A1. (Associative) $\forall u, v, w \in V, (u + v) + w = u + (v + w)$;
- A2. (existence of zero element) there exists an element 0 in V such that $\forall u \in V, u + 0 = u$;
- A3. (existence of inverse element) $\forall u \in V \exists v \in V$ such that $u + v = 0$;
- A4. (Commutative) $\forall u, v \in V, u + v = v + u$;
- M1. (distributive) $\forall \alpha \in F$ and $\forall u, v \in V, \alpha \cdot (u + v) = \alpha \cdot u + \alpha \cdot v$;
- M2. (distributive) $\forall \alpha, \beta \in F$ and $\forall u \in V, (\alpha \oplus \beta) \cdot u = \alpha \cdot u + \beta \cdot u$;
- M3. $\forall \alpha, \beta \in F$ and $\forall u \in V, (\alpha \odot \beta) \cdot u = \alpha \cdot (\beta \cdot u)$;
- M4. $\forall u \in V, 1 \cdot u = u$.

Elements of a vector space are called vectors.

¹The proof of that result is very similar to the proof of Proposition 134.1 and 2.

²See, for example, Apostol (1967), Section 13.2, page 17.

Remark 133 In the remainder of these notes, if no confusion arises, for ease of notation, we will denote a field simply by F and a vector space by V . Moreover, we will write $+$ in the place of \oplus , and we will omit \odot and \cdot , i.e., we will write xy instead of $x \odot y$ and αv instead of $\alpha \cdot v$.

Proposition 134 If V is a vector space, then (as a consequence of the first four properties)

1. The zero vector is unique and it is denoted by 0 .
2. $\forall u \in V$, the inverse element of u is unique and it is denoted by $-u$.
3. (cancellation law) $\forall u, v, w \in V$,

$$u + w = v + w \Rightarrow u = v.$$

Proof. 1. Assume that there exist $0_1, 0_2 \in V$ which are zero vectors. Then by definition of zero vector - see (A2) - we have that

$$0_1 + 0_2 = 0_1 \quad \text{and} \quad 0_2 + 0_1 = 0_2.$$

From (A.4),

$$0_1 + 0_2 = 0_2 + 0_1,$$

and therefore $0_1 = 0_2$.

2. Given $u \in V$, assume there exist $v^1, v^2 \in V$ such that

$$u + v^1 = 0 \quad \text{and} \quad u + v^2 = 0.$$

Then

$$v^2 = v^2 + 0 = v^2 + (u + v^1) = (v^2 + u) + v^1 = (u + v^2) + v^1 = 0 + v^1 = v^1.$$

- 3.

$$u + w = v + w \xrightarrow{(1)} u + w + (-w) = v + w + (-w) \xrightarrow{(2)} u + 0 = v + 0 \xrightarrow{(3)} u = v,$$

where (1) follows from the definition of operation, (2) from the definition of $-w$ and (3) from the definition of 0 . ■

Remark 135 From A2. in Definition 132, we have that for any vector space V , $0 \in V$.

Proposition 136 If V is a vector space over a field F , then

1. For $0 \in F$ and $\forall u \in V$, $0u = 0$.
2. For $0 \in V$ and $\forall \alpha \in F$, $\alpha 0 = 0$.
3. If $\alpha \in F$, $u \in V$ and $\alpha u = 0$, then either $\alpha = 0$ or $u = 0$ or both.
4. $\forall \alpha \in F$ and $\forall u \in V$, $(-\alpha)u = \alpha(-u) = -(\alpha u) := -\alpha u$.

Proof. 1. From (M1),

$$0u + 0u = (0 + 0)u = 0u.$$

Then, adding $-(0u)$ to both sides,

$$0u + 0u + (-(0u)) = 0u + (-(0u))$$

and, using (A3),

$$0u + 0 = 0$$

and, using (A2), we get the desired result.

2. From (A2),

$$0 + 0 = 0;$$

then multiplying both sides by α and using (M1),

$$\alpha 0 = \alpha(0 + 0) = \alpha 0 + \alpha 0;$$

and, using (A3),

$$\alpha 0 + (- (\alpha 0)) = \alpha 0 + \alpha 0 + (- (\alpha 0))$$

and, using (A2), we get the desired result.

3. Assume that $\alpha u = 0$ and $\alpha \neq 0$. Then

$$u = 1u = (\alpha^{-1} \cdot \alpha) u = \alpha^{-1} (\alpha u) = \alpha^{-1} \cdot 0 = 0.$$

Taking the contrapositive of the above result, we get $\langle u \neq 0 \rangle \Rightarrow \langle \alpha u \neq 0 \vee \alpha = 0 \rangle$. Therefore $\langle u \neq 0 \wedge \alpha u = 0 \rangle \Rightarrow \langle \alpha = 0 \rangle$.

4. From $u + (-u) = 0$, we get $\alpha(u + (-u)) = \alpha 0$, and then $\alpha u + \alpha(-u) = 0$, and therefore $-(\alpha u) = \alpha(-u)$.

From $\alpha + (-\alpha) = 0$, we get $(\alpha + (-\alpha))u = 0u$, and then $\alpha u + (-\alpha)u = 0$, and therefore $-(\alpha u) = (-\alpha)u$. ■

Remark 137 From Proposition 136.4, and (M4) in Definition 132, we have

$$(-1)u = 1(-u) = -(1u) = -u.$$

We also define subtraction as follows:

$$v - u := v + (-u)$$

4.2 Examples

Euclidean spaces.³

The Euclidean space \mathbb{R}^n with sum and scalar product defined in Chapter 2 is a vector space over the field \mathbb{R} with the standard addition and multiplication

Matrices on \mathbb{R} .

For any $m, n \in \mathbb{N}$, the set $\mathcal{M}_{m,n}$ of matrices with elements belonging to the field \mathbb{R} with the operation of addition and scalar multiplication, as defined in Section 2.3 is a vector space on the field \mathbb{R} and it is denoted by

$$\mathbb{M}(m, n).$$

Matrices on a field F .

For any $m, n \in \mathbb{N}$, we can also consider the set of matrices whose entries are elements belonging to an arbitrary field F . It is easy to check that set is a vector space on the field F , with the operation of addition and scalar multiplication inherited by F , is a vector space and it is denoted by

$$\mathbb{M}_F(m, n).$$

We do set

$$\mathbb{M}_{\mathbb{R}}(m, n) = \mathbb{M}(m, n).$$

Polynomials

The set of *all* polynomials

$$a_0 + a_1 t + a_2 t^2 + \dots + a_n t^n$$

with $n \in \mathbb{N} \cup \{0\}$ and $a_0, a_1, a_2, \dots, a_n \in \mathbb{R}$ is a vector space on \mathbb{R} with respect to the standard sum between polynomials and scalar multiplication.

Function space $\mathcal{F}(X)$.

Given a nonempty set X , the set of all functions $f : X \rightarrow \mathbb{R}$ with obvious sum and scalar multiplication is a vector space on \mathbb{R} . More generally we can consider the vector space of functions $f : X \rightarrow F$, where X is a nonempty set and F is an arbitrary field, on the same field F .

Sets which are not vector spaces.

$(0, +\infty)$ and $[0, +\infty)$ are **not** a vector spaces in \mathbb{R} .

For any $n \in \mathbb{N}$, the set of all polynomials of degree n is not a vector space on \mathbb{R} .

On the role of the field: put Exercise 5.29 from Lipschutz 2nd edition.

³A detailed proof of some of the statements below is contained, for example, in Hoffman and Kunze (1971), starting from page 28.

4.3 Vector subspaces

In what follows, if no ambiguity may arise, we will say “vector space” instead of “vector space on a field”.

Definition 138 Let W be a subset of a vector space V . W is called a vector subspace of V if W is a vector space with respect to the operation of vector addition and scalar multiplication defined on V restricted to W . In other words, given a vector space $(V, +, \cdot)$ on a field (F, \oplus, \odot) and a subset W of V , we say that W is a vector subspace of V if, defined

$$+_W : W \times W \rightarrow W, \quad (w_1, w_2) \mapsto w_1 + w_2$$

and

$$\cdot_W : F \times W \rightarrow W, \quad (\alpha, w_2) \mapsto \alpha \cdot w_2,$$

then $(W, +_W, \cdot_W)$ is a vector space on the field (F, \oplus, \odot) .

Proposition 139 Let W be a subset of a vector space V . The following three statements are equivalent.

1. W is a vector subspace of V .
2. a. $W \neq \emptyset$;
b. $\forall u, v \in W, u + v \in W$;
c. $\forall u \in W, \alpha \in F, \alpha u \in W$.
3. a. $W \neq \emptyset$;
b. $\forall u, v \in W, \forall \alpha, \beta \in F, \alpha u + \beta v \in W$.

Proof. 2. \Rightarrow 3.

From 2c., $\alpha u \in W$ and $\beta v \in W$. Then, from 2b., $\alpha u + \beta v \in W$, as desired.

2. \Leftarrow 3.

From 3b., identifying α, u, β, v with $1, u, 1, v$, we get $u + v \in W$, i.e., 2b.

From 3b., identifying α, u, β, v with $\frac{1}{2}\alpha, u, \frac{1}{2}\alpha, u$, we get

$$\frac{1}{2}\alpha u + \frac{1}{2}\alpha u \in W. \quad (4.1)$$

Moreover, since $W \subseteq V$, then $u \in V$ and from the distributive property (M1) for V , we get

$$\frac{1}{2}\alpha u + \frac{1}{2}\alpha u = \left(\frac{1}{2}\alpha + \frac{1}{2}\alpha\right)u = \alpha u, \quad (4.2)$$

and from (4.1) and (4.2) the desired result follows.

1. \Rightarrow 2.

All properties follow immediately from the definition of vector space.

1. \Leftarrow 2.

2a., 2b. and 2c. insure that the “preliminary properties” of a vector space are satisfied: $W \neq \emptyset$ and the operations are well defined.

Properties A1, A4, M1, M2, M3 and M4 are satisfied because $W \subseteq V$. Let’s check the remaining two properties.

(A2): We want to show that there exists $0_W \in W$ such that $\forall w \in W$, we have $w + 0_W = w$.

Since $W \neq \emptyset$, we can take $w \in W$. Since $W \subseteq V$, then $w \in V$ and from Proposition 136.1,

$$0_F w = 0_V. \quad (4.3)$$

Moreover, from 2c., identifying α, u with $0_F, w$, we get $0_V \in W$. Then $\forall w \in W \subseteq V, w + 0_V = w$, i.e.,

$$0_W = 0_V := 0. \quad (4.4)$$

(A3): We want to show that $\forall w \in W \exists w' \in W$ such that $w + w' = 0$.

From 2c, we have $(-1)w \in W$ and $1w \in W$. Then

$$(-1)w + w \stackrel{(M4)}{=} (-1)w + 1w \stackrel{(M1)}{=} (-1 + 1)w = 0_F w \stackrel{(?)}{=} 0.$$

Taking $w' = (-1)w$, we have shown the desired result. ■

Remark 140 An often successful way to show that a set S is a vector space is the following one:

1. find a vector space V such that $S \subseteq V$; in Section 4.2 above, we provide a list of “commonly used” vector spaces;
2. use Proposition 139.

Example 141 1. Given an arbitrary vector space V , $\{0\}$ and V are vector subspaces of V .

2. Given \mathbb{R}^3 ,

$$W := \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_3 = 0\}$$

is a vector subspace of \mathbb{R}^3 .

3. Given the space V of polynomials, the set W of all polynomials of degree $\leq n$ is a vector subspace of V .

4. The set of all bounded or continuous or differentiable or integrable functions $f : X \rightarrow \mathbb{R}$ is a vector subspace of $\mathcal{F}(X)$.

5. If V and W are vector spaces, then $V \cap W$ is a vector subspace of V and W .

6. $[0, +\infty)$ is not a vector subspace of \mathbb{R} .

7. Let $V = \{0\} \times \mathbb{R} \subseteq \mathbb{R}^2$ and $W = \mathbb{R} \times \{0\} \subseteq \mathbb{R}^2$. Then $V \cup W$ is not a vector subspace of \mathbb{R}^2 .

4.4 Linear combinations

Notation convention. Unless otherwise stated, a greek (or Latin) letter with a subscript denotes a scalar; a Latin letter with a superscript denotes a vector.

Before introducing the main concept of the section, we need to introduce the definition of sequence. Intuitively, a sequence is an “enumerated collection of objects” in which repetitions are allowed and order is important.

Like a set, it contains members (also called elements, or terms). The number of elements (possibly infinite) is sometimes called the length of the sequence.

Unlike a set, order matters, and exactly the same elements can appear multiple times at different positions in the sequence. Formally, we have what follows

Definition 142 A sequence is a function whose domain is either the set \mathbb{N} of the natural numbers (for infinite sequences) or the set of the first n natural numbers (for a sequence of finite length n). The position of an element in a sequence is its index: it is the integer the element is the image of.

Remark 143 In other words, given a nonempty set X , an infinite sequence in X is a function $x : \mathbb{N} \rightarrow X$, and a finite sequence of length $n \in \mathbb{N}$ is a function $x : \{1, \dots, n\} \rightarrow X$.

Usually, in the case of infinite sequences, for any $i \in \mathbb{N}$, the value $x(i)$ is denoted by x_i , which is called the i -th term of the sequence; the sequence is denoted by $(x_i)_{i \in \mathbb{N}}$. Similarly, for finite sequences of length $n \in \mathbb{N}$, for any $i \in \{1, \dots, n\}$, the value $x(i)$ is denoted by x_i ; the sequence is denoted by $(x_i)_{i \in \{1, \dots, n\}}$ or (x_1, \dots, x_n) .

Definition 144 Let V be a vector space on a field F , $m \in \mathbb{N}$ and $v^1, v^2, \dots, v^m \in V$ be given. The linear combination of the ordered list, or order m -tuple or finite sequence of vectors v^1, v^2, \dots, v^m via scalars $\alpha_1, \alpha_2, \dots, \alpha_m \in F$ is the vector

$$\sum_{i=1}^m \alpha_i v^i.$$

The set of all such combinations

$$\left\{ v \in V : \exists (\alpha_i)_{i=1}^m \in F^m \text{ such that } v = \sum_{i=1}^m \alpha_i v^i \right\}$$

is called span of (v^1, v^2, \dots, v^m) and it is denoted by

$$\text{span}(v^1, v^2, \dots, v^m).$$

add examples

Definition 145 Let V be a vector space and S a subset of V . $\text{span}(S)$ is the set of all linear combinations of a finite sequence of vectors in S , i.e.,

$$\text{span}(S) = \left\{ v \in V : \exists m \in \mathbb{N}, \exists v^1, v^2, \dots, v^m \in S, \exists (\alpha_i)_{i=1}^m \in F^m \text{ such that } v = \sum_{i=1}^m \alpha_i v^i \right\}.$$

$$\text{span}(\emptyset) := \{0\}.$$

Proposition 146 Let V be a vector space and $S \neq \emptyset, S \subseteq V$. Then, “ $\text{span}(S)$ is the smallest vector space containing S ”, i.e.,

1. a. $S \subseteq \text{span}(S)$ and b. $\text{span}(S)$ is a vector subspace of V .
2. If W is a subspace of V and $S \subseteq W$, then $\text{span}(S) \subseteq W$.

Proof. 1a. Given $v \in S$, $1v = v \in \text{span}(S)$. 1b. Since $S \neq \emptyset$, then $\text{span}(S) \neq \emptyset$. Given $\alpha, \beta \in F$ and $v, w \in \text{span}(S)$. Then $\exists \alpha_1, \dots, \alpha_n \in F, v^1, \dots, v^n \in S$ and $\beta_1, \dots, \beta_m \in F, w^1, \dots, w^m \in S$, such that $v = \sum_{i=1}^n \alpha_i v^i$ and $w = \sum_{j=1}^m \beta_j w^j$. Then

$$\alpha v + \beta w = \sum_{i=1}^n (\alpha \alpha_i) v^i + \sum_{j=1}^m (\beta \beta_j) w^j \in \text{span}(S).$$

2. Take $v \in \text{span}(S)$. Then $\exists \alpha_1, \dots, \alpha_n \in F, v^1, \dots, v^n \in S \subseteq W$ such that $v = \sum_{i=1}^n \alpha_i v^i \in W$, as desired. ■

Definition 147 Let V be a vector space and $v^1, v^2, \dots, v^m \in V$. If $V = \text{span}(v^1, v^2, \dots, v^m)$, we say that V is the vector space generated or spanned by the vectors v^1, v^2, \dots, v^m .

Example 148 1. $\mathbb{R}^3 = \text{span}(\{(1, 0, 0), (0, 1, 0), (0, 0, 1)\})$;
 2. $\text{span}(\{(1, 1)\}) = \text{span}(\{(1, 1), (2, 2)\}) = \{(x, y) \in \mathbb{R}^2 : x = y\}$;
 3. $\text{span}(\{(1, 1), (0, 1)\}) = \mathbb{R}^2$.
 2. $\text{span}(\{t^n\}_{n \in \mathbb{N}})$ is equal to the vector space of all polynomials.

Exercise 149 1. If $A \subseteq B$, then $\text{span}(A) \subseteq \text{span}(B)$;
 2. if W is a vector subspace of V , then $\text{span}(W) = W$.

4.5 Row and column space of a matrix

Definition 150 Given $A \in \mathbb{M}(m, n)$,

$$\text{row span } A := \text{span}(R^1(A), \dots, R^i(A), \dots, R^m(A))$$

is called the row space of A or row span of A .

The column space of A or colspan A is

$$\text{col span } A := \text{span}(C^1(A), \dots, C^j(A), \dots, C^m(A)).$$

Remark 151 Given $A \in \mathcal{M}(m, n)$

$$\text{col span } A = \text{row span } A^T.$$

Remark 152 *Linear combinations of columns and rows of a matrix.*

Let $A \in \mathbb{M}(m, n)$, $x \in \mathbb{R}^n$ and $y \in \mathbb{R}^m$. Then, $\forall j \in \{1, \dots, n\}$, $C^j(A) \in \mathbb{R}^m$ and $\forall i \in \{1, \dots, m\}$, $R^i(A) \in \mathbb{R}^n$. Then,

Remark 153

$$Ax = \sum_{j=1}^n x_j \cdot C^j(A),$$

as verified below.

$$Ax = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \dots & & & & & \\ a_{i1} & a_{i2} & \dots & a_{ij} & \dots & a_{in} \\ \dots & & & & & \\ a_{m1} & a_{m2} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \\ \dots \\ x_j \\ \dots \\ x_n \end{bmatrix} = \begin{bmatrix} \sum_{j=1}^n a_{1j}x_j \\ \sum_{j=1}^n a_{2j}x_j \\ \dots \\ \sum_{j=1}^n a_{ij}x_j \\ \dots \\ \sum_{j=1}^n a_{mj}x_j \end{bmatrix};$$

$$\sum_{j=1}^n x_j \cdot C^j(A) = \sum_{j=1}^n x_j \cdot \begin{bmatrix} a_{1j} \\ a_{2j} \\ \dots \\ a_{ij} \\ \dots \\ a_{mj} \end{bmatrix} = \begin{bmatrix} \sum_{j=1}^n x_j a_{1j} \\ \sum_{j=1}^n x_j a_{2j} \\ \dots \\ \sum_{j=1}^n x_j a_{ij} \\ \dots \\ \sum_{j=1}^n x_j a_{mj} \end{bmatrix}.$$

Therefore,

Ax is a linear combination of the columns of A via the components of the vector x .

Moreover,

$$yA = \sum_{i=1}^m y_i \cdot R^i(A),$$

as verified below.

$$yA = [y_1, \dots, y_i, \dots, y_m] \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \dots & & & & & \\ a_{i1} & a_{i2} & \dots & a_{ij} & \dots & a_{in} \\ \dots & & & & & \\ a_{m1} & a_{m2} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix} =$$

$$= \left[\sum_{i=1}^m y_i \cdot a_{i1} \quad \sum_{i=1}^m y_i \cdot a_{i2} \quad \dots \quad \sum_{i=1}^m y_i \cdot a_{ij} \quad \dots \quad \sum_{i=1}^m y_i \cdot a_{in} \right];$$

$$\sum_{i=1}^m y_i \cdot R^i(A) = \sum_{i=1}^m y_i \cdot \begin{bmatrix} a_{i1} & a_{i2} & \dots & a_{ij} & \dots & a_{in} \end{bmatrix} =$$

$$= \left[\sum_{i=1}^m y_i \cdot a_{i1} \quad \sum_{i=1}^m y_i \cdot a_{i2} \quad \dots \quad \sum_{i=1}^m y_i \cdot a_{ij} \quad \dots \quad \sum_{i=1}^m y_i \cdot a_{in} \right].$$

Therefore,

yA is a linear combination of the rows of A via the components of the vector y .

As a consequence of the above observation, we have what follow.

1.

$$\text{row span } A = \{w \in \mathbb{R}^n : \exists y \in \mathbb{R}^m \text{ such that } w = yA\}.$$

2.

$$\text{colspan } A = \{z \in \mathbb{R}^m : \exists x \in \mathbb{R}^n \text{ such that } z = Ax\};$$

Proposition 154 Given $A, B \in \mathbb{M}(m, n)$,

1. if A is row equivalent to B , then $\text{row span } A = \text{row span } B$;
2. if A is column equivalent to B , then $\text{colspan } A = \text{colspan } B$.

Proof. 1. B is obtained by A via elementary row operations. Therefore, $\forall i \in \{1, \dots, m\}$, either
i. $R^i(B) = R^i(A)$, or
ii. $R^i(B)$ is a linear combination of rows of A .

Therefore, $\text{row span } B \subseteq \text{row span } A$. Since A is obtained by B via elementary row operations, $\text{row span } B \supseteq \text{row span } A$.

2. if A is column equivalent to B , then A^T is row equivalent to B^T and therefore, from i. above, $\text{row span } A^T = \text{row span } B^T$. Then the result follows from Remark 151. ■

Remark 155 Let $A \in \mathbb{M}(m, n)$ be given and assume that

$$b := (b_j)_{j=1}^n = \sum_{i=1}^m c_i \cdot R^i(A),$$

i.e., b is a linear combination of the rows of A . Then,

$$\forall j \in \{1, \dots, n\}, \quad b_j = \sum_{i=1}^m c_i \cdot R^{ij}(A),$$

where $\forall i \in \{1, \dots, m\}$ and $\forall j \in \{1, \dots, n\}$, $R^{ij}(A)$ is the j -th component of the i -th row $R^i(A)$ of A .

Lemma 156 Assume that $A, B \in \mathbb{M}(m, n)$ are in echelon form with pivots

$$a_{1j_1}, \dots, a_{ij_i}, \dots, a_{rj_r},$$

and

$$b_{1k_1}, \dots, b_{ik_i}, \dots, b_{sk_s},$$

respectively, and⁴ $r, s \leq \min\{m, n\}$. Then

$$\langle \text{row span } A = \text{row span } B \rangle \Rightarrow \langle s = r \text{ and for } i \in \{1, \dots, s\}, \quad j_i = k_i \rangle.$$

Proof. Preliminary remark 1. If $A = 0$, then $A = B$ and $s = r = 0$.

Preliminary remark 2. Assume that $A, B \neq 0$ and then $s, r \geq 1$. We want to verify that $j_1 = k_1$. Suppose $j_1 < k_1$. Then, by definition of echelon matrix, $C_{j_1}(B) = 0$, otherwise you would contradict Property 2 of the Definition 28 of echelon matrix. Then, from the assumption that $\text{row span } A = \text{row span } B$, we have that $R^1(A)$ is a linear combination of the rows of B , via some coefficients c_1, \dots, c_m , and from Remark 155 and the fact that $C^{j_1}(B) = 0$, we have that $a_{1j_1} = c_1 \cdot 0 + \dots + c_m \cdot 0 = 0$, contradicting the fact that a_{1j_1} is a pivot for A . Therefore, $j_1 \geq k_1$. A perfectly symmetric argument shows that $j_1 \leq k_1$.

We can now prove the result by induction on the number m of rows.

Step 1. $m = 1$.

It is basically the proof of Preliminary Remark 2.

Step 2.

Given $A, B \in \mathbb{M}(m, n)$, define $A', B' \in \mathbb{M}(m-1, n)$ as the matrices obtained erasing the first row in matrix A and B respectively. From Remark 33, A' and B' are still in echelon form. If we show that $\text{row span } A' = \text{row span } B'$, from the induction assumption, and using Preliminary Remark 2, we get the desired result.

Let $R = (a_1, \dots, a_n)$ be any row of A' . Since $R \in \text{row span } B$, $\exists (d_i)_{i=1}^m$ such that

$$R = \sum_{i=1}^m d_i R^i(B).$$

Since A is in echelon form and we erased its first row, we have that if $i \leq j_1 = k_1$, then $a_i = 0$, otherwise you would contradict the definition of j_1 . Since B is in echelon form, each entry in its k_1 -th column are zero, but b_{1k_1} which is different from zero. Then,

$$a_{1k_1} = 0 = \sum_{i=1}^m d_i \cdot b_{ik_1} = d_1 \cdot b_{1k_1},$$

and therefore $d_1 = 0$, i.e., $R = \sum_{i=2}^m d_i R^i(B)$, or $R \in \text{row span } B'$, as desired. Symmetric argument shows the other inclusion. ■

⁴See Remark 30.

Remark 157 *Given*

$$A = \begin{bmatrix} 1 & 1 \\ 2 & 1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 2 & 2 \\ 6 & 3 \end{bmatrix},$$

clearly $A \neq B$ and $\text{row span } A = \text{row span } B$.

Proposition 158 *Assume that $A, B \in \mathbb{M}(m, n)$ are in row canonical form. Then,*

$$\langle \text{row span } A = \text{row span } B \rangle \Leftrightarrow \langle A = B \rangle.$$

Proof. $[\Leftarrow]$ Obvious.

$[\Rightarrow]$ From Lemma 156, the number of pivots in A and B is the same. Therefore, A and B have the same number s of nonzero rows, which in fact are the first s rows. Take $i \in \{1, \dots, s\}$. Since $\text{row span } A = \text{row span } B$, there exists $(c_h)_{h=1}^s$ such that

$$R^i(A) = \sum_{h=1}^s c_h \cdot R^h(B). \quad (4.5)$$

We want then to show that $c_i = 1$ and $\forall l \in \{1, \dots, s\} \setminus \{i\}, c_l = 0$.

Let a_{ij_i} be the pivot of $R^i(A)$, i.e., a_{ij_i} is the nonzero j_i -th component of $R^i(A)$. Then, from Remark 155,

$$a_{ij_i} = \sum_{h=1}^s c_h \cdot R^{hj_i}(B) = \sum_{h=1}^s c_h \cdot b_{hj_i}. \quad (4.6)$$

From Lemma 156, for $i \in \{1, \dots, s\}$, $j_i = k_i$, and therefore b_{ij_i} is a pivot entry for B , and since B is in row reduced form, b_{ij_i} is the only nonzero element in the j_i column of B . Therefore, from (4.6),

$$a_{ij_i} = \sum_{h=1}^s c_h \cdot R^{hj_i}(B) = c_i \cdot b_{ij_i}.$$

Since A and B are in row canonical form $a_{ij_i} = b_{ij_i} = 1$ and therefore

$$c_i = 1.$$

Now take $l \in \{1, \dots, s\} \setminus \{i\}$ and consider the pivot element b_{lj_l} in $R^l(B)$. From (4.5) and Remark 155,

$$a_{lj_l} = \sum_{h=1}^s c_h \cdot b_{hj_l} = c_l, \quad (4.7)$$

where the last equalities follow from the fact that B is in row reduced form and therefore b_{lj_l} is the only nonzero element in the j_l -th column of B , in fact, $b_{lj_l} = 1$. From Lemma 156, since b_{lj_l} is a pivot element for B , a_{lj_l} is a pivot element for A . Since A is in row reduced form, a_{lj_l} is the only nonzero element in column j_l of A . Therefore, since $l \neq i$, $a_{ij_l} = 0$, and from (4.7), the desired result,

$$\forall l \in \{1, \dots, s\} \setminus \{i\}, \quad c_l = 0.,$$

does follow. ■

Proposition 159 *For every $A \in \mathbb{M}(m, n)$, there exists a unique $B \in \mathbb{M}(m, n)$ which is in row canonical form and row equivalent to A .*

Proof. The existence of at least one matrix with the desired properties is the content of Proposition 38. Suppose that there exists B_1 and B_2 with those properties. Then from Proposition 154, we get

$$\text{row span } A = \text{row span } B_1 = \text{row span } B_2.$$

From Proposition 158,

$$B_1 = B_2.$$

■

Corollary 160 1. For any matrix $A \in \mathbb{M}(m, n)$ there exists a unique number $r \in \{0, 1, \dots, \min\{m, n\}\}$ such that A is equivalent to the block matrix of the form

$$\begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}.$$

2. For any $A \in \mathbb{M}(m, n)$, there exist invertible matrices $P \in \mathbb{M}(m, m)$ and $Q \in \mathbb{M}(n, n)$ and a unique number $r \in \{0, 1, \dots, \min\{m, n\}\}$ such that

$$PAQ = \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}$$

Proof. 1.

From Step 1 in the proof of Proposition 122 and from Proposition 159, there exists a unique matrix A^* which is row equivalent to A and it is in row canonical form.

From Step 2 and 3 in the proof of Proposition 122 and from Proposition 159, there exist a unique matrix

$$\begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}^T$$

which is row equivalent to A^{*T} and it is in row canonical form. Therefore the desired result follows.

2.

From Proposition 105.2, PA is row equivalent to A ; from Proposition 119.2, PAQ is column equivalent to PA . Therefore, PAQ is equivalent to A . From Proposition 124,

$$\begin{bmatrix} I_{r'} & 0 \\ 0 & 0 \end{bmatrix}$$

is equivalent to A . From part 1 of the present Proposition, the desired result then follows. ■

Example 161 Let $A = \begin{bmatrix} 2 & 2 \\ 1 & 1 \end{bmatrix}$ be given. Then,

$$\begin{bmatrix} 2 & 2 \\ 1 & 1 \end{bmatrix} \xrightarrow{\frac{1}{2}R^1 \rightarrow R^1} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \xrightarrow{R^2 - R^1 \rightarrow R^2} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \xrightarrow{C^2 - C^1 \rightarrow C^1} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}.$$

Indeed,

$$\begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 2 & 2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix},$$

and

$$P = \begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} & 0 \\ -\frac{1}{2} & 1 \end{bmatrix},$$

$$Q = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix}.$$

4.6 Linear dependence and independence

Definition 162 Let a vector space V on a field F , a number $m \in \mathbb{N}$ and vectors $v^1, v^2, \dots, v^m \in V$ be given. The finite sequence of vectors (v^1, v^2, \dots, v^m) is linearly dependent, or the vectors v^1, v^2, \dots, v^m are linearly dependent, if

either $m = 1$ and $v^1 = 0$,

or $m > 1$ and $\exists k \in \{1, \dots, m\}$ and there exist $(m - 1) \geq 1$ coefficients $\alpha_j \in F$ with $j \in \{1, \dots, m\} \setminus \{k\}$ such that

$$v^k = \sum_{j \in \{1, \dots, m\} \setminus \{k\}} \alpha_j v^j$$

or, shortly,

$$v^k = \sum_{j \neq k} \alpha_j v^j$$

i.e., there exists a vector equal to a linear combination of the other vectors.

Geometrical interpretation of linear (in)dependence in \mathbb{R}^2 .

Proposition 163 Let a vector space V on a field F and vectors $v^1, v^2, \dots, v^m \in V$ be given. The vectors $v^1, v^2, \dots, v^m \in V$ are linearly dependent vectors if and only if

$$\exists (\beta_1, \dots, \beta_i, \dots, \beta_m) \in F^m \setminus \{0\} \text{ such that } \sum_{i=1}^m \beta_i \cdot v^i = 0, \quad (4.8)$$

i.e., there exists a linear combination of the vectors equal to the null vector and with some nonzero coefficient.

Proof. $[\Rightarrow]$

If $m = 1$, any $\beta \in \mathbb{R} \setminus \{0\}$ is such that $\beta \cdot 0 = 0$. Assume then that $m > 1$. Take

$$\beta_i = \begin{cases} \alpha_i & \text{if } i \neq j \\ -1 & \text{if } i = j \end{cases}$$

$[\Leftarrow]$

If $m = 1$, $\exists \beta \in \mathbb{R} \setminus \{0\}$ is such that $\beta \cdot v^1 = 0$, then from Proposition 58.3 $v^1 = 0$. Assume then that $m > 1$. Without loss of generality take $\beta_1 \neq 0$. Then,

$$\beta_1 v^1 + \sum_{i \neq 1} \beta_i v^i = 0$$

and

$$v^1 = \sum_{i \neq 1} \frac{\beta_i}{\beta_1} v^i.$$

■

Proposition 164 Let $m \geq 2$ and (v^1, \dots, v^m) be a finite sequence of nonzero linearly dependent vectors. Then, one of the vectors is a linear combination of the preceding vectors, i.e., $\exists k > 1$ and $(\alpha^i)_{i=1}^{k-1}$ such that $v^k = \sum_{i=1}^{k-1} \alpha_i v^i$.

Proof. Since (v^1, \dots, v^m) is linearly dependent, $\exists (\beta_i)_{i=1}^m \in \mathbb{R}^m \setminus \{0\}$ such that $\sum_{i=1}^m \beta_i v^i = 0$. Let k be the largest i such that $\beta_i \neq 0$, i.e.,

$$k \in \{1, \dots, m\} \text{ is such that } \beta_k \neq 0 \text{ and } \forall i \in \{k+1, \dots, m\}, \beta_i = 0. \quad (4.9)$$

Consider the case $k = 1$. Then we would have $\beta_1 \neq 0$ and $\forall i > 1, \beta_i = 0$ and therefore $0 = \sum_{i=1}^m \beta_i v^i = \beta_1 v^1$, and $v^1 = 0$, contradicting the assumption that v^1, \dots, v^m are nonzero vectors. Then, we must have $k > 1$, and from (4.9), we have

$$0 = \sum_{i=1}^m \beta_i v^i = \sum_{i=1}^k \beta_i v^i$$

and

$$\beta_k v^k = \sum_{i=1}^{k-1} \beta_i v^i,$$

or, as desired,

$$v^k = \sum_{i=1}^{k-1} \frac{-\beta_i}{\beta_k} v^i.$$

It is then enough to choose $\alpha_i = \frac{-\beta_i}{\beta_k}$ for any $i \in \{1, \dots, k-1\}$. ■

Example 165 Take the vectors $x^1 = (1, 2)$, $x^2 = (-1, -2)$ and $x^3 = (0, 4)$. x^1, x^2, x^3 are linearly dependent vectors: $x^1 = -1 \cdot x^2 + 0 \cdot x^3$. Observe that there are no $\alpha_1, \alpha_2 \in \mathbb{R}$ such that $x^3 = \alpha_1 \cdot x^1 + \alpha_2 \cdot x^2$.

Definition 166 The finite sequence of vectors (v^1, v^2, \dots, v^m) is linearly independent, or the vectors v^1, v^2, \dots, v^m are linearly independent vectors if they are not linearly dependent.

Remark 167 The finite sequence of vectors (v^1, v^2, \dots, v^m) is linearly independent if $\langle \neg(4.8) \rangle$ holds true, i.e., if

$$\forall (\beta_1, \dots, \beta_i, \dots, \beta_m) \in F^m \setminus \{0\} \text{ it is the case that } \sum_{i=1}^m \beta_i \cdot v^i \neq 0,$$

or

$$(\beta_1, \dots, \beta_i, \dots, \beta_m) \in F^m \setminus \{0\} \Rightarrow \sum_{i=1}^m \beta_i \cdot v^i \neq 0$$

or

$$\sum_{i=1}^m \beta_i \cdot v^i = 0 \Rightarrow (\beta_1, \dots, \beta_i, \dots, \beta_m) = 0$$

or the only linear combination of the vectors which is equal to the null vector has each coefficient equal to zero.

Example 168 The vectors $(1, 2)$, $(1, 5)$ are linearly independent. The vectors $(1, 5)$, $(1, 2)$ are linearly independent.

Remark 169 Let L_1 be a finite sequence (or an ordered list) of vectors and L_2 a finite sequence obtained from L_1 changing the order of the vectors. Then, since sum is commutative, L_1 is linearly independent if and only if L_2 is linearly independent, and therefore, L_1 is linearly dependent if and only if L_2 is linearly dependent.

Remark 170 From Remark 152, we have what follows:

$$\langle Ax = 0 \Rightarrow x = 0 \rangle \Leftrightarrow \langle \text{the column vectors of } A \text{ are linearly independent} \rangle \quad (4.10)$$

$$\langle yA = 0 \Rightarrow y = 0 \rangle \Leftrightarrow \langle \text{the row vectors of } A \text{ are linearly independent} \rangle \quad (4.11)$$

Remark 171 We also need to speak of linearly dependent and independent **sets**. As already said, the difference between an ordered list or a finite sequence on one side, and a set on the other side is that in the latter concept (the concept of set) both order and repetitions “do not count”. We distinguish sets from finite sequence using curly brackets, i.e., we use $\{\}$ for sets and round brackets, i.e., $()$ or no brackets at all, for finite sequences. Then

$$\{v, u, w\} = \{w, u, v\} = \{v, v, u, w, w, w, w\}, \quad (4.12)$$

⁵ and

$$(v, u, w) \neq (w, u, v),$$

$$(v, u, w) \neq (v, v, u, w, w, w, w).$$

Observe also that the cardinality of each set in (4.12) is 3: in computing the cardinality of sets, “repetitions and order do not count”.

In the definition of linearly dependent and linearly independent sets we require to consider **distinct** vectors.

Definition 172 Let a vector space V on a field F and a set S (of arbitrary cardinality) contained in V be given.

S is linearly dependent if there exists a finite sequence of **distinct** vectors in S which is linearly dependent.

S is linearly independent if any finite sequence of **distinct** vectors in S is linearly independent.

⁵Recall that given two sets A and B , we have that $A = B$ if and only if

1. $a \in A \Rightarrow a \in B$ and 2. $b \in B \Rightarrow b \in A$.

Recall that a function is a triple: domain, codomain and “rule”. Two functions are equal if the corresponding triples are equal.

Proposition 173 Let $m \in \mathbb{N}$ and a set $S = \{v^1, \dots, v^m\}$ be given.

1. any finite sequence of all distinct elements of S is linearly independent. \Leftrightarrow a finite sequence of all distinct elements of S is linearly independent.
2. any finite sequence of all distinct elements of S is linearly dependent. \Leftrightarrow a finite sequence of all distinct elements of S is linearly dependent.

Proof. Without loss of generality, let (v^1, v^2, \dots, v^n) with $n \leq m$ be a finite sequence of all the distinct elements of S . Observe that there are many (indeed, $n! := n \cdot (n-1) \cdot \dots \cdot 3 \cdot 2$) different finite sequences of all distinct elements of S . For example, given $S = \{v^1, v^2, v^1\}$ with $v^1 \neq v^2$, then both (v^1, v^2) and (v^2, v^1) are finite sequences of all distinct elements of S .

1. $[\Rightarrow]$ Obvious.

$[\Leftarrow]$ It follows from Remark 169.

2. The proof is similar to the one in point 1. above. ■

Proposition 174 Let $m \in \mathbb{N}$ and a set $S = \{v^1, \dots, v^m\}$ be given.

1. S is linearly independent \Leftrightarrow a finite sequence of all distinct elements of S is linearly independent.
2. S is linearly dependent \Leftrightarrow a finite sequence of all distinct elements of S is linearly dependent.

Proof. 1.

$[\Rightarrow]$

Obvious.

$[\Leftarrow]$

Without loss of generality, let (v^1, v^2, \dots, v^n) with $n \leq m$ be a finite sequence of all the distinct elements of S . Assume our claim is false and, without loss of generality, $\exists k < n$ such that $\{v^1, \dots, v^k\} \subsetneq \{v^1, \dots, v^k, v^{k+1}, \dots, v^n\}$ and $\exists \beta \in \mathbb{R}^k \setminus \{0\}$ such that $\sum_{i=1}^k \beta_i v^i = 0$. Then, chosen $\beta_{k+1} = \dots = \beta_n = 0$, we have that there exists $\beta \in \mathbb{R}^n \setminus \{0\}$ such that $\sum_{i=1}^n \beta_i v^i = 0$, i.e., the finite sequence $(v^1, \dots, v^k, v^{k+1}, \dots, v^n)$ is linearly dependent, contradicting the assumption.

2.

It is enough to take the contrapositive of each implication in statement 1. above. In other words, the desired result follows from the fact that $p \Leftrightarrow q$ is logically equivalent to $(\neg p) \Leftrightarrow (\neg q)$.

$[\Leftarrow]$

From 1. above, we have

S is linearly independent \Rightarrow a finite sequence of all distinct elements of S is linearly independent,

and therefore

any finite sequence of all distinct elements of S is linearly dependent $\Rightarrow S$ is linearly dependent. (4.13)

From Proposition 173,

a finite sequence of all distinct elements of S is linearly dependent

\Leftrightarrow

any finite sequence of all distinct elements of S is linearly dependent. (4.14)

From (4.14) and (4.13), the desired result follows.

$[\Rightarrow]$

Similar argument applies. ■

Remark 175 \emptyset is a set of linearly independent vectors.

Remark 176 (Recipe for analyzing linear(in)dependence).

Summarizing, we can say what follow to deal with the problem of determining linear (in)dependence.

1. If you have a finite sequence of vectors, try to show all of them are linearly (in)dependent, using Definition 162 or Proposition 163, or Remark 167;
2. If you a finite cardinality set, then eliminate the repetitions (i.e., consider a finite ordered list of all pairwise distinct vectors belonging to the set) and then use 1. above (recall that from Remark 169, the order of the chosen list is irrelevant with respect to linear (in)dependence). Sometimes it can hard to eliminate repetitions; if you are able to show that the finite sequence you are dealing with is linearly independent, then the set is linearly independent as well.
3. If you have an infinite cardinality set, then apply Definition 172.

Remark 177 *To greatly simplify the exposition, when we consider a finite set $S = \{v^1, \dots, v^m\}$ in a vector space V , unless otherwise clearly specified, we assume that the cardinality of S is m , i.e., all elements in S are pairwise distinct.*

Example 178 *Consider the vectors $x^1 = (1, 0, 0, 0) \in \mathbb{R}^4$ and $x^2 = (0, 1, 0, 0)$. Observe that $\alpha_1 x^1 + \alpha_2 x^2 = 0$ means $(\alpha_1, \alpha_2, 0, 0) = (0, 0, 0, 0)$.*

Exercise 179 *Say if the following set is a set of linearly dependent or independent vectors:*

$$S = \{x^1, x^2, x^3\} \text{ with } x^1 = (3, 2, 1), \quad x^2 = (4, 1, 3), \quad x^3 = (3, -3, 6).$$

Exercise 180 *Let a vector space V and $u, v \in V$ such that $u \neq v$ be given. Show that if $\{u, v\}$ is linearly independent, then $\{u + v, v\}$ is linearly independent.*

First of all, observe that, by assumption,

$$\forall \alpha_1, \alpha_2 \in F \text{ such that } \alpha_1 v + \alpha_2 u = 0, \text{ we have } \alpha_1 = \alpha_2 = 0. \quad (4.15)$$

We now want to apply the “Recipe” described in Remark 176.

Observe that $u + v \neq v$: suppose otherwise, then we would have $u + v = v$ and $u = 0$, contradicting the fact that $\{u, v\}$ is linearly independent. Then, following the Recipe, we want to show that

$$\forall \beta_1, \beta_2 \in F \text{ such that } \beta_1 (u + v) + \beta_2 v = 0, \text{ we have } \beta_1 = \beta_2 = 0.$$

Indeed, $0 = \beta_1 (u + v) + \beta_2 v = \beta_1 u + (\beta_1 + \beta_2) v$, and from (4.15), we have $\beta_1 = \beta_1 + \beta_2 = 0$, which easily implies the desired result.

Example 181 *Let V be the vector space of all functions $f : \mathbb{R} \rightarrow \mathbb{R}$. f, g, h defined below are linearly independent:*

$$f(x) = e^x, \quad g(x) = x^2, \quad h(x) = x.$$

Suppose there exists $(\alpha_1, \alpha_2, \alpha_3) \in \mathbb{R}^3$ such that

$$\alpha_1 \cdot f + \alpha_2 \cdot g + \alpha_3 \cdot h = 0_V,$$

which means that

$$\forall x \in \mathbb{R}, \quad \alpha_1 \cdot f(x) + \alpha_2 \cdot g(x) + \alpha_3 \cdot h(x) = 0.$$

We want to show that $(\alpha_1, \alpha_2, \alpha_3) = (0, 0, 0)$. The trick is to find appropriate values of x to get the desired value of $(\alpha_1, \alpha_2, \alpha_3)$. Choose x to take values $0, 1, -1$. We obtain the following system of equations

$$\begin{cases} \alpha_1 & & & = & 0 \\ e \cdot \alpha_1 & + & \alpha_2 & + & \alpha_3 & = & 0 \\ e^{-1} \alpha_1 & + & \alpha_2 & + & (-1) \alpha_3 & = & 0 \end{cases}$$

It then follows that $(\alpha_1, \alpha_2, \alpha_3) = (0, 0, 0)$, as desired.

Remark 182 *The following statements are easy consequences of Definition 172.*

1. *Any set which contains a linearly dependent set is linearly dependent.*
2. *Any subset of a linearly independent set is linearly independent.*

Remark 183 *Consider vectors in \mathbb{R}^n .*

1. *Adding components to linearly dependent vectors gives raise to linearly dependent or independent vectors;*
2. *Eliminating components from linearly independent vectors gives raise to linearly dependent or independent vectors;*
3. *Adding components to linearly independent vectors gives raise to linearly independent vectors;*
4. *Eliminating components from linearly dependent vectors gives raise to linearly dependent vectors.*

To verify 1. and 2. above consider the following two vectors:

$$\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 1 \end{bmatrix}$$

To verify 4., if you have a set of linearly dependent vectors, it is possible to express one vector as a linear combination of the others and then eliminate components leaving the equality still true. The proof of 3 is contained in the following Proposition.

Proposition 184 If $S_x = \{x^1, \dots, x^m\} \subseteq \mathbb{R}^n$ is a set of linearly independent vectors and $S_y = \{y^1, \dots, y^m\} \subseteq \mathbb{R}^k$ is a set of vectors, then $S = \{(x^1, y^1), \dots, (x^m, y^m)\} \subseteq \mathbb{R}^{n+k}$ is a set of linearly independent vectors.

Proof. By assumption

$$\sum_{i=1}^m \beta_i \cdot v^i = 0 \quad \Rightarrow \quad (\beta_1, \dots, \beta_i, \dots, \beta_m) = 0.$$

S is a set of linearly independent vectors if

$$\sum_{i=1}^m \beta_i \cdot (v^i, y^i) = 0 \quad \Rightarrow \quad (\beta_1, \dots, \beta_i, \dots, \beta_m) = 0.$$

Since

$$\sum_{i=1}^m \beta_i \cdot (v^i, y^i) = 0 \quad \Leftrightarrow \quad \sum_{i=1}^m \beta_i \cdot v^i = 0 \text{ and } \sum_{i=1}^m \beta_i \cdot y^i = 0$$

the desired result follows. ■

Corollary 185 If $S_x = \{x^1, \dots, x^m\} \subseteq \mathbb{R}^n$ is a set of linearly independent vectors and $S_y = \{y^1, \dots, y^m\} \subseteq \mathbb{R}^k$ and $S_z = \{z^1, \dots, z^m\} \subseteq \mathbb{R}^l$ are sets of vectors, then

$$S = \{(z_1, x^1, y^1), \dots, (z_m, x^m, y^m)\} \subseteq \mathbb{R}^{l+n+k}$$

is a set of linearly independent vectors.

Example 186 1. The set of vectors $\{v^1, \dots, v^m\}$ is a linearly dependent set if $\exists k \in \{1, \dots, m\}$ such that $v^k = 0$:

$$v^k + \sum_{i \neq k} 0 \cdot v^i = 0$$

2. The set of vectors $\{v^1, \dots, v^m\}$ is a linearly dependent set if $\exists k, k' \in \{1, \dots, m\}$ and $\alpha \in \mathbb{R} \setminus \{1\}$ such that $v^{k'} = \alpha v^k$:

$$v^{k'} - \alpha v^k + \sum_{i \neq k, k'} 0 \cdot v^i = 0$$

Remark 187 In many textbook, the expression

$$(v^1, \dots, v^m) \text{ is a finite sequence of linearly (in)dependent vectors,}$$

is written as

$$v^1, \dots, v^m \text{ are linearly (in)dependent vectors.}$$

Remark 188 When we talk of linear (in)dependence of rows or columns of a matrix we refer to the ordered list or finite sequence of rows or columns, **not to the set of rows or columns**. Indeed, while the rows of the matrix

$$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$

are linearly dependent, the set $\{(1, 1), (1, 1)\} = \{(1, 1)\}$ is linearly independent.

Proposition 189 *The nonzero rows of a matrix A in echelon form are linearly independent.*

Proof. We will show that each row of A starting from the first one is not a linear combination of the subsequent rows. Then, as a consequence of Proposition 164, the desired result will follow.

Since A is in echelon form, the first row has a pivot below which all the elements are zero. Then that row cannot be a linear combination of the following rows. Similar argument applies to the other rows. ■

Remark 190 (Herstein (1975), page 178) *“We point out that linear dependence is a function not only of the vectors but also of the field. For instance, the field of complex numbers is a vector space over the field of real numbers and it is also a vector space over the field of complex numbers. The elements $v_1 = 1, v_2 = i$ in it are linearly independent over the reals but linearly dependent over the complexes, since $iv_1 + (-1)v_2 = 0$.”*

4.7 Basis and dimension

Definition 191 *A set S (of arbitrary cardinality) in a vector space V on a field F is a basis of V if*

1. S is a linearly independent set;
2. $\text{span}(S) = V$.

Lemma 192 *Suppose that given a vector space V , $\text{span}(\{v^1, \dots, v^m\}) = V$.*

1. If $w \in V$ and $w \notin \{v^1, \dots, v^m\}$, then a. $\{w, v^1, \dots, v^m\}$ is linearly dependent and
b. $\text{span}(\{w, v^1, \dots, v^m\}) = V$;
2. If v^i is a linear combination of $\{v^j\}_{j=1}^{i-1}$, then $\text{span}(\{v^1, \dots, v^{i-1}, v^{i+1}, \dots, v^m\}) = V$.

Proof. 1.a.

Since, by assumption, $w \notin \{v^1, \dots, v^m\}$, then all elements in the set $\{w, v^1, \dots, v^m\}$ are distinct; then, from the Recipe in Remark 176, it is enough to show that (w, v^1, \dots, v^m) is linearly dependent.

Since $w \in V \stackrel{\text{Assn.}}{=} \text{span}(\{v^1, \dots, v^m\})$, then by definition of span, there exist $\beta_1, \dots, \beta_m \in F$ such that $w = \sum_{i=1}^m \beta_i v^i$, as desired.

b.

By assumption, there exist $\beta_1, \dots, \beta_m \in F$ such that $w = \sum_{i=1}^m \beta_i v^i$. We want to show that $\text{span}(\{v^1, \dots, v^m\}) = \text{span}(\{w, v^1, \dots, v^m\})$

[\subseteq] We want to show that $v = \sum_{i=1}^m \alpha_i v^i$ implies that there exist $\alpha_0, \alpha_1, \dots, \alpha_m \in F$ such that $v = \alpha_0 w + \sum_{i=1}^m \alpha_i v^i$; it is enough to take $\alpha_0 = 0$ and for $i \in \{1, \dots, m\}$, $\alpha_i = \delta_i$.

[\supseteq] We want to show that $v = \alpha_0 w + \sum_{i=1}^m \alpha_i v^i$ implies that there exist $\delta_1, \dots, \delta_m \in F$ such that $v = \sum_{i=1}^m \delta_i v^i$.

Since, by assumption $w \in V = \text{span}(\{v^1, \dots, v^m\})$, then there exist $\beta_1, \dots, \beta_m \in F$ such that

$$w = \sum_{i=1}^m \beta_i v^i. \text{ Then, } v = \alpha_0 w + \sum_{i=1}^m \alpha_i v^i = v = \alpha_0 \sum_{i=1}^m \beta_i v^i + \sum_{i=1}^m \alpha_i v^i = \sum_{i=1}^m (\alpha_0 \beta_i + \alpha_i) v^i,$$

as desired.

2.

By assumption, there exists $\{\beta_j \in F : j \in \{1, \dots, m\} \setminus \{i\}\}$ such that $v^i = \sum_{j \neq i} \beta_j v^j$. We want to show that $\text{span}(\{v^1, \dots, v^{i-1}, v^{i+1}, \dots, v^m\}) = \text{span}(\{v^1, \dots, v^m\})$. Then, the proof is very similar to the proof of 1b. above. ■

Lemma 193 (Replacement Lemma) *Let a vector space V and sets of vectors $\{v^1, \dots, v^n\}$ and $\{w^1, \dots, w^m\}$ contained in V be given. If*

1. $\text{span}(\{v^1, \dots, v^n\}) = V$,

2. $\{w^1, \dots, w^m\}$ is linearly independent,

then

1. $n \geq m$,

2. a. If $n = m$, then $\text{span}(\{w^1, \dots, w^m\}) = V$.

b. if $n > m$, there exists $\{v^{i_1}, \dots, v^{i_{n-m}}\} \subseteq \{v^1, \dots, v^n\}$ such that

$$\text{span}(\{w^1, \dots, w^m, v^{i_1}, \dots, v^{i_{n-m}}\}) = V.$$

Proof. Observe preliminary that since $\{w^1, \dots, w^m\}$ is linearly independent, for any $j \in \{1, \dots, m\}$, $w^j \neq 0$.

We now distinguish 2 cases: Case 1. For any $i \in \{1, \dots, n\}$, $v^i \neq 0$, and Case 2. There exists $i \in \{1, \dots, n\}$ such that $v^i = 0$.

Case 1.

Now consider the case $n = 1$. $\{w^1, \dots, w^m\} \subseteq V$ implies that there exists $(\alpha_j)_{j=1}^m \in \mathbb{R}^m$ such that $\forall j$, $\alpha_j \neq 0$ and $w^j = \alpha^j v^1$, then it has to be $m = 1$ (and conclusion 1 holds) and since $w^1 = \alpha_1 v^1$, $\text{span}(w^1) = V$ (and conclusion 2 holds).

Consider now the case $n \geq 2$.

If $w^1 \in \{v^1, \dots, v^n\}$, we immediately obtain that there exists $k_1 \in \{1, \dots, n\}$ such that

$$\text{span}(\{w^1\} \cup \{v^i : i \in \{1, \dots, n\} \setminus \{k_1\}\}) = V.$$

If $w^1 \notin \{v^1, \dots, v^n\}$, from Lemma 192.1, $\{w^1, v^1, \dots, v^n\}$ is linearly dependent and

$$\text{span}(\{w^1, v^1, \dots, v^n\}) = V.$$

Since elements in $\{w^1, v^1, \dots, v^n\}$ are pairwise distinct, from the Recipe in Remark 176, we have that $\{w^1, v^1, \dots, v^n\}$ is linearly dependent iff (w^1, v^1, \dots, v^n) is linearly dependent and from Lemma 164, there exists $k_1 \in \{1, \dots, n\}$ such that v^{k_1} is a linear combination of the preceding vectors. Then from Lemma 192.2, we have

$$\text{span}(\{w^1\} \cup \{v^i : i \in \{1, \dots, n\} \setminus \{k_1\}\}) = V.$$

If $w^2 \in \{v^i : i \in \{1, \dots, n\} \setminus \{k_1\}\}$, we immediately obtain that there exists $k_2 \in \{1, \dots, n\} \setminus \{k_1\}$ such that

$$\text{span}(\{w^1, w^2\} \cup \{v^i : i \in \{1, \dots, n\} \setminus \{k_1, k_2\}\}) = V.$$

If $w^2 \notin \{v^i : i \in \{1, \dots, n\} \setminus \{k_1\}\}$, then from Lemma 192.1, $\{w^1, w^2\} \cup \{v^i : i \in \{1, \dots, n\} \setminus \{k_1\}\}$ is linearly dependent and

$$\text{span}(\{w^1, w^2\} \cup \{v^i : i \in \{1, \dots, n\} \setminus \{k_1\}\}) = V.$$

Then, as above, by Lemma 164, there exists $k_2 \in \{2, \dots, n\} \setminus \{k_1\}$ such that v^{k_2} or w^2 is a linear combination of the preceding vectors. w^2 cannot be a linear combination of w^1 because of assumption 2. Therefore, from Lemma 192.2, we have

$$\text{span}(\{w^1, w^2\} \cup \{v^i : i \in \{1, \dots, n\} \setminus \{k_1, k_2\}\}) = V.$$

We can now distinguish three cases: $m < n$, $m = n$ and $m > n$.

Now if $m < n$, after m steps of the above procedure we get

$$\text{span}(\{w^1, \dots, w^m\} \cup \{v^i : i \in \{1, \dots, n\} \setminus \{k_1, k_2, \dots, k_m\}\}) = V,$$

which shows 2.a. If $m = n$, we have

$$\text{span}(\{w^1, \dots, w^m\}) = V,$$

which shows 2.b.

Let's now show that it cannot be $m > n$. Suppose that is the case. Then, after n of the above steps, we get $\text{span}(w^1, \dots, w^n) = V$ and therefore w^{n+1} is a linear combination of (w^1, \dots, w^n) , contradicting assumption 2.

Case 2.

In the present case, we assume that there exists a set I_0 such that $I_0 \neq \emptyset$ and $I_0 \subseteq \{1, \dots, n\}$. Define also $I_1 = \{1, \dots, n\} \setminus I_0$ and $n_1 = \#I_1$. Clearly,

$$\text{span}(\{v^i : i \in I_1\}) = \text{span}(\{v^1, \dots, v^n\}) = V$$

and $n_1 < n$. Then, repeating the “replacement” argument described in Case 1 applied to $\{v^i : i \in I_1\}$ and $\{w^1, \dots, w^m\}$, we get

- (i) $n > n_1 \geq m$, i.e., Conclusion 1 holds and Conclusion 2a does not hold true, and
- (ii) there exists

$$\{v^{i_1}, \dots, v^{i_{\#I_1-m}}\} \subseteq \{v^i : i \in I_1\} \subseteq \{v^1, \dots, v^n\}$$

such that

$$\text{span}(\{w^1, \dots, w^m, v^{i_1}, \dots, v^{i_{\#I_1-m}}\}) = V,$$

and therefore Conclusion 2b does hold true. ■

Proposition 194 Assume that $S = \{u^1, u^2, \dots, u^n\}$ and $T = \{v^1, v^2, \dots, v^m\}$ are bases of V . Then $n = m$.

Proof. By definition of basis we have that

$$\text{span}(\{u^1, u^2, \dots, u^n\}) = V \quad \text{and} \quad \{v^1, v^2, \dots, v^m\} \text{ are linearly independent.}$$

Then from Lemma 193, $m \leq n$. Similarly,

$$\text{span}(\{v^1, v^2, \dots, v^m\}) = V \quad \text{and} \quad \{u^1, u^2, \dots, u^n\} \text{ are linearly independent,}$$

and from Lemma 193, $n \leq m$. ■

The above Proposition allows to give the following Definition.

Definition 195 A vector space V has dimension $n \in \mathbb{N}$ if there exists a basis of V whose cardinality is n . In that case, we say that V has finite dimension (equal to n) and we write $\dim V = n$. If a vector space does not have finite dimension, it is said to be of infinite dimension.

Definition 196 The vector space $\{0\}$ has dimension 0.

Example 197 1. A basis of \mathbb{R}^n is $\{e^1, \dots, e^i, \dots, e^n\}$, where e^i is defined in Definition 55. That basis is called canonical basis. Then $\dim \mathbb{R}^n = n$.

2. Consider the vector space $\mathbb{P}_n(t)$ of polynomials of degree $\leq n$. The set $\{t^0, t^1, \dots, t^n\}$ of polynomials is a basis of $\mathbb{P}_n(t)$ and therefore $\dim \mathbb{P}_n(t) = n + 1$.

Example 198 Put the example of infinite dimensional space from Hoffman and Kunze, page 43.

Proposition 199 Let V be a vector space of dimension n .

1. $m > n$ vectors in V are linearly dependent;
2. If $S = \{u^1, \dots, u^n\} \subseteq V$ is a linearly independent set, then it is a basis of V ;
3. If $\text{span}(u^1, \dots, u^n) = V$, then $\{u^1, \dots, u^n\}$ is a basis of V .

Proof. Let $\{w^1, \dots, w^n\}$ be a basis of V .

1. We want to show that $\{v^1, \dots, v^m\}$ arbitrary vectors in V are linearly dependent. Suppose otherwise, then by Lemma 193, we would have $m \leq n$, a contradiction.
2. It is the content of Lemma 193.2.a.

3. We have to show that $\{u^1, \dots, u^n\}$ are linearly independent. Suppose otherwise. Then there exists $k \in \{1, \dots, n\}$ such that $\text{span}\left(\left\{(u^i)_{i \neq k}\right\}\right) = V$, but since $\{w^1, \dots, w^n\}$ is linearly independent, from Proposition 193 (the Replacement Lemma), we have $n \leq n-1$, a contradiction.

■

Remark 200 The above Proposition 199 shows that in the case of finite dimensional vector spaces, one of the two conditions defining a basis is sufficient to obtain a basis.

Proposition 201 (Completion Lemma) Let V be a vector space of dimension n and $\{w^1, \dots, w^m\} \subseteq V$ be a linearly independent set, with⁶ $m \leq n$. If $m < n$, then, there exists a set $\{u^1, \dots, u^{n-m}\}$ such that

$$\{w^1, \dots, w^m, u^1, \dots, u^{n-m}\}$$

is a basis of V .

Proof. Take a basis $\{v^1, \dots, v^n\}$ of V . Then from Conclusion 2.b in Lemma 193,

$$\text{span}(w^1, \dots, w^m, v^{i_1}, \dots, v^{i_{n-m}}) = V.$$

Then from Proposition 199.3, we get the desired result. ■

Proposition 202 Let W be a subspace of an n -dimensional vector space V . Then

1. $\dim W \leq n$;
2. If $\dim W = n$, then $W = V$.

Proof. 1. From Proposition 199.1, $m > n$ vectors in V are linearly dependent. Since a basis of W is a set of linearly independent vectors then $\dim W \leq n$.

2. If $\{w^1, \dots, w^n\}$ is a basis of W , then $\text{span}(w^1, \dots, w^n) = W$. Moreover, those vectors are n linearly independent vectors in V . Therefore from Proposition 199.2., $\text{span}(w^1, \dots, w^n) = V$. ■

Remark 203 As a trivial consequence of Proposition 201, $V = \text{span}\{u^1, \dots, u^r\} \Rightarrow \dim V \leq r$.

Example 204 Let W be a subspace of \mathbb{R}^3 , whose dimension is 3. Then from the previous Proposition, $\dim W \in \{0, 1, 2, 3\}$. In fact,

1. If $\dim W = 0$, then $W = \{0\}$, i.e., a point,
2. if $\dim W = 1$, then W is a straight line through the origin,
3. if $\dim W = 2$, then W is a plane through the origin,
4. if $\dim W = 3$, then $W = \mathbb{R}^3$.

4.8 Coordinates

Definition 205 If V is a finite-dimensional vector space, an ordered basis for V is a finite sequence of vectors which is linearly independent and spans V .

Remark 206 If the finite sequence v^1, \dots, v^n is an ordered basis for V , then the vectors in $\{v^1, \dots, v^n\}$ have to be pairwise distinct, otherwise the finite sequence (v^1, \dots, v^n) would be linearly dependent. We will make some abuse of notation saying that the set

$$\{v^1, \dots, v^n\}$$

is an ordered basis for V , i.e., a basis of V is a linearly independent “ordered set” which spans V .

Proposition 207 $S = \{u^1, u^2, \dots, u^n\} \subseteq V$ is an ordered basis of V on a field F if and only if $\forall v \in V$ there exists a unique $(\alpha_i)_{i=1}^n \in F^n$ such that $v = \sum_{i=1}^n \alpha_i u^i$.

⁶The inequality $m \leq n$ follows from Proposition 199.1.

Proof. $[\Rightarrow]$ Suppose there exist $(\alpha_i)_{i=1}^n, (\beta_i)_{i=1}^n \in \mathbb{R}^n$ such that $v = \sum_{i=1}^n \alpha_i u^i = \sum_{i=1}^n \beta_i u^i$. Then

$$0 = \sum_{i=1}^n \alpha_i u^i - \sum_{i=1}^n \beta_i u^i = \sum_{i=1}^n (\alpha_i - \beta_i) u^i.$$

Since $\{u^1, u^2, \dots, u^n\}$ are linearly independent,

$$\forall i \in \{1, \dots, n\}, \alpha_i - \beta_i = 0,$$

as desired.

$[\Leftarrow]$

Clearly $V = \text{span}(S)$; we are left with showing that $\{u^1, u^2, \dots, u^n\}$ are linearly independent. Consider $\sum_{i=1}^n \alpha_i u^i = 0$. Moreover, $\sum_{i=1}^n 0 \cdot u^i = 0$. But since there exists a unique $(\alpha_i)_{i=1}^n \in \mathbb{R}^n$ such that $v = \sum_{i=1}^n \alpha_i u^i$, it must be the case that $\forall i \in \{1, \dots, n\}, \alpha_i = 0$. ■

The above proposition allows to give the following definition.

Definition 208 Given a vector space V on a field F with an ordered basis $\mathcal{V} = \{v^1, \dots, v^n\}$, the associated coordinate function is

$$cr_{\mathcal{V}} : V \rightarrow F^n, \quad v \mapsto cr_{\mathcal{V}}(v) := [v]_{\mathcal{V}}$$

where $cr_{\mathcal{V}}(v)$ is the unique vectors of coefficients which give v as a linear combination of the element of the ordered basis, i.e., $cr_{\mathcal{V}}(v) := (\alpha_i)_{i=1}^n$ such that $v = \sum_{i=1}^n \alpha_i v^i$.

4.9 Row and column span

We start our analysis with a needed lemma.

Lemma 209 Given a vector space V , if

1. $(v^1, \dots, v^k) \subseteq V$ is linearly independent, and
 2. $v^{k+1} \in V$ and $(v^1, \dots, v^k, v^{k+1})$ is linearly dependent,
- then v^{k+1} is a linear combination of the vectors in (v^1, \dots, v^k) .

Proof. Since (v^1, \dots, v^k) is linearly dependent,

$$\exists i \in \{1, \dots, k+1\}, (\beta_j)_{j \in \{1, \dots, k+1\} \setminus \{i\}} \text{ such that } v^i = \sum_{j \in \{1, \dots, k+1\} \setminus \{i\}} \beta_j v^j.$$

If $i = k+1$, we are done. If $i \neq k+1$, without loss of generality, take $i = 1$. Then

$$\exists (\gamma_j)_{j \in \{1, \dots, k+1\} \setminus \{1\}} \text{ such that } v^1 = \sum_{j=2}^{k+1} \gamma_j v^j.$$

If $\gamma_{k+1} = 0$, we would have $v^1 - \sum_{j=2}^k \gamma_j v^j = 0$, contradicting Assumption 1. Then

$$v^{k+1} = \frac{1}{\gamma_{k+1}} \left(v^1 - \sum_{j=2}^k \gamma_j v^j \right).$$

■

Remark 210 Let V be a vector space and $S \subseteq T \subseteq V$. Then,

1. $\text{span } S \subseteq \text{span } T$;
2. $\text{span}(\text{span } S) = \text{span } S$.

Definition 211 For any $A \in \mathbb{M}(m, n)$,

the row rank of A is the $\dim(\text{row span of } A)$;
the column rank of A is the $\dim(\text{col span of } A)$.

Proposition 212 For any $A \in \mathbb{M}(m, n)$, row rank of A is equal to column rank of A .

Proof. Let A be an arbitrary $m \times n$ matrix

$$\begin{bmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ \dots & & & & \\ a_{i1} & \dots & a_{ij} & \dots & a_{in} \\ \dots & & & & \\ a_{m1} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix}$$

Suppose the row rank is $r \leq m$ and the following r vectors form a basis of the row space:

$$\begin{bmatrix} S_1 & = & [b_{11} & \dots & b_{1j} & \dots & b_{1n}] \\ \dots & & & & & & \\ S_k & = & [b_{k1} & \dots & b_{kj} & \dots & b_{kn}] \\ \dots & & & & & & \\ S_r & = & [b_{r1} & \dots & b_{rj} & \dots & b_{rn}] \end{bmatrix}$$

Then, each row vector of A is a linear combination of the above vectors, i.e., we have

$$\forall i \in \{1, \dots, m\}, \quad R^i = \sum_{k=1}^r \alpha_{ik} S_k,$$

or

$$\forall i \in \{1, \dots, m\}, \quad [a_{i1} \quad \dots \quad a_{ij} \quad \dots \quad a_{in}] = \sum_{k=1}^r \alpha_{ik} [b_{k1} \quad \dots \quad b_{kj} \quad \dots \quad b_{kn}],$$

and setting the j component of each of the above vector equations equal to each other, we have

$$\forall j \in \{1, \dots, n\} \quad \text{and} \quad \forall i \in \{1, \dots, m\}, \quad a_{ij} = \sum_{k=1}^r \alpha_{ik} b_{kj},$$

and

$$\forall j \in \{1, \dots, n\}, \quad \begin{cases} a_{1j} & = & \sum_{k=1}^r \alpha_{1k} b_{kj}, \\ \dots & & \\ a_{ij} & = & \sum_{k=1}^r \alpha_{ik} b_{kj}, \\ \dots & & \\ a_{mj} & = & \sum_{k=1}^r \alpha_{mk} b_{kj}, \end{cases}$$

or

$$\forall j \in \{1, \dots, n\}, \quad \begin{bmatrix} a_{1j} \\ \dots \\ a_{ij} \\ \dots \\ a_{mj} \end{bmatrix} = \sum_{k=1}^r b_{kj} \begin{bmatrix} \alpha_{1k} \\ \dots \\ \alpha_{ik} \\ \dots \\ \alpha_{mk} \end{bmatrix},$$

i.e., each column of A is a linear combination of the r vectors

$$\left\{ \begin{pmatrix} \alpha_{11} \\ \dots \\ \alpha_{i1} \\ \dots \\ \alpha_{m1} \end{pmatrix}, \dots, \begin{pmatrix} \alpha_{1k} \\ \dots \\ \alpha_{ik} \\ \dots \\ \alpha_{mk} \end{pmatrix}, \dots, \begin{pmatrix} \alpha_{1r} \\ \dots \\ \alpha_{ir} \\ \dots \\ \alpha_{mr} \end{pmatrix} \right\}.$$

Then, from Remark 203,

$$\dim \text{col span } A \leq r = \text{row rank } A, \quad (4.16)$$

i.e.,

$$\text{col rank } A \leq \text{row rank } A.$$

From (4.16), which holds for arbitrary matrix A , we also get

$$\dim \text{col span } A^T \leq \text{row rank } A^T. \quad (4.17)$$

Moreover,

$$\dim \operatorname{col} \operatorname{span} A^T \stackrel{\text{Rmk. 151}}{=} \dim \operatorname{row} \operatorname{span} A := \operatorname{row} \operatorname{rank} A$$

and

$$\operatorname{row} \operatorname{rank} A^T := \dim \operatorname{row} \operatorname{span} A^T \stackrel{\text{Rmk. 151}}{=} \dim \operatorname{col} \operatorname{span} A.$$

Then, from the two above equalities and (4.17), we get

$$\operatorname{row} \operatorname{rank} A \leq \dim \operatorname{col} \operatorname{span} A, \quad (4.18)$$

and (4.16) and (4.18) gives the desired result. ■

Proposition 213 For any $A \in \mathbb{M}(m, n)$,

1. the row rank of A is equal to the maximum number of linearly independent rows of A ;
2. the column rank of A is equal to the maximum number of linearly independent columns of A .

Proof. 1.

Let $k_1 := \operatorname{row} \operatorname{rank} A$ and $k_2 :=$ maximum number of linearly independent rows of A . Assume our claim is false and therefore, either a. $k_1 > k_2$, or b. $k_1 < k_2$.

a.

Let (v^1, \dots, v^{k_2}) a finite sequence of linearly independent rows of A . Since, by assumption, k_2 is the maximum number of linearly independent rows of A , then, from Lemma 209, the other rows of A are a linear combination of v^1, \dots, v^{k_2} and from Lemma 192, $\operatorname{span}(v^1, \dots, v^{k_2}) = \operatorname{row} \operatorname{span} A$. Then, (v^1, \dots, v^{k_2}) is a basis of $\operatorname{span} A$ and therefore $k_1 := \operatorname{row} \operatorname{rank} A := \dim \operatorname{row} \operatorname{span} A = k_2$, contradicting the assumption of case a.

b.

In this case we would have k_2 linearly independent vectors in a vector space of dimension k_1 , with $k_2 > k_1$, contradicting Proposition 199.1.

2.

The proof is basically the same as the above one. ■

Corollary 214 For every $A \in \mathbb{M}(m, n)$,

$$\begin{aligned} \operatorname{row} \operatorname{rank} A &= \text{maximum number of linearly independent rows of } A = \\ &= \text{maximum number of linearly independent columns of } A = \operatorname{col} \operatorname{rank} A. \end{aligned} \quad (4.19)$$

Proof. It follows from Propositions 212 and 213. ■

Exercise 215 Check the above result on the following matrix

$$\begin{bmatrix} 1 & 2 & 3 \\ 5 & 1 & 6 \\ 3 & 7 & 10 \end{bmatrix}.$$

4.10 Exercises

Problem sets: 1,2,3,4,5,6,7.

From Lipschutz (1991), starting from page 162:

5.3, 5.7, 5.8, 5.9, 5.10, 5.12 \rightarrow 5.15, 5.17 \rightarrow 5.23, 5.24 \rightarrow 5.29, 5.31 \rightarrow 5.34, 5.46 \rightarrow 5.49.

Chapter 5

Determinant and rank of a matrix

In this chapter we are going to introduce the definition of determinant, an useful tool to study linear dependence, invertibility of matrices and solutions to systems of linear equations.

5.1 Definition and properties of the determinant of a matrix

To motivate the definition of determinant, we present an informal discussion of a way to find solutions to the linear system with two equations and two unknowns, shown below.

$$\begin{cases} a_{11}x_1 + a_{12}x_2 = b_1 \\ a_{21}x_1 + a_{22}x_2 = b_2 \end{cases} \quad (5.1)$$

The system can be rewritten as follows

$$Ax = b$$

where

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}, \quad x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \quad b = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}.$$

Let's informally discuss how to find solutions to system (5.1). If $a_{22} \neq 0$ and $a_{12} \neq 0$, multiplying both sides of the first equation by a_{22} , of the second equation by $-a_{12}$ and adding up, we get

$$a_{11}a_{22}x_1 + a_{12}a_{22}x_2 - a_{12}a_{21}x_1 - a_{12}a_{22}x_2 = a_{22}b_1 - a_{12}b_2$$

Therefore, if

$$a_{11}a_{22} - a_{12}a_{21} \neq 0$$

we have

$$x_1 = \frac{b_1a_{22} - b_2a_{12}}{a_{11}a_{22} - a_{12}a_{21}} \quad (5.2)$$

In a similar manner¹ we have

$$x_2 = \frac{b_2a_{11} - b_1a_{21}}{a_{11}a_{22} - a_{12}a_{21}} \quad (5.3)$$

We can then the following preliminary definition: given $A \in \mathcal{M}_{22}$, the determinant of A is

$$\det A = \det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} := a_{11}a_{22} - a_{12}a_{21}$$

Using the definition of determinant, we can rewrite (5.2) and (5.3) as follows.

$$x_1 = \frac{\det \begin{bmatrix} b_1 & a_{12} \\ b_2 & a_{22} \end{bmatrix}}{\det A} \quad \text{and} \quad x_2 = \frac{\det \begin{bmatrix} a_{11} & b_1 \\ a_{21} & b_2 \end{bmatrix}}{\det A}$$

¹Assuming $a_{21} \neq 0$ and $a_{11} \neq 0$, multiply both sides of the first equation by a_{21} , of the second equation by $-a_{11}$ and then add up.

We can now present the definition of the determinant of a **square** matrix $A_{n \times n}$ for arbitrary $n \in \mathbb{N}$.

Definition 216 Given $n > 1$ and $A \in \mathbb{M}(n, n)$, $\forall i, j \in \{1, \dots, n\}$, we call $A_{ij} \in \mathbb{M}(n-1, n-1)$ the matrix obtained from A erasing the i -th row and the j -th column.

Definition 217 Given $A \in \mathbb{M}(1, 1)$, i.e., $A = [a]$ with $a \in \mathbb{R}$. The determinant of A is denoted by $\det A$ and we let $\det A := a$. For $\mathbb{N} \setminus \{0, 1\}$, given $A \in \mathbb{M}(n, n)$, we define the determinant of A as

$$\det A := \sum_{j=1}^n (-1)^{1+j} a_{1j} \det A_{1j}$$

Observe that $[a_{1j}]_{j=1}^n$ is the first row of A , i.e.,

$$\det A := \sum_{j=1}^n (-1)^{1+j} R^{1j}(A) \det A_{1j}$$

Example 218 For $n = 2$, we have

$$\begin{aligned} \det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} &= \sum_{j=1}^2 (-1)^{1+j} a_{1j} \det A_{1j} = (-1)^{1+1} a_{11} \det A_{11} + (-1)^{1+2} a_{12} \det A_{12} = \\ &= a_{11}a_{22} - a_{12}a_{21} \end{aligned}$$

and we get the informal definition given above.

Example 219 For $n = 3$, we have

$$\det A = \det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = (-1)^{1+1} a_{11} \det A_{11} + (-1)^{1+2} a_{12} \det A_{12} + (-1)^{1+3} a_{13} \det A_{13}$$

Definition 220 Given $A = [a_{ij}] \in \mathbb{M}(n, n)$, $\forall i, j$, $\det A_{ij}$ is called minor of a_{ij} in A ;

$$(-1)^{i+j} \det A_{ij}$$

is called cofactor of a_{ij} in A .

Theorem 221 Given $A \in \mathbb{M}(n, n)$, $\det A$ is equal to the sum of the products of the elements of any rows or column for the corresponding cofactors, i.e.,

$$\forall i \in \{1, \dots, n\}, \det A = \sum_{j=1}^n (-1)^{i+j} R^{ij}(A) \det A_{ij} \quad (5.4)$$

and

$$\forall j \in \{1, \dots, n\}, \det A = \sum_{i=1}^n (-1)^{i+j} C^{ji}(A) \det A_{ij} \quad (5.5)$$

Proof. Omitted. We are going to omit several proofs about determinants. There are different ways of introducing the concept of determinant of a square matrix. One of them uses the concept of permutations - see, for example, Lipschutz (1991), Chapter 7. Another one is an axiomatic approach - see, for example, Lang (1971) - he introduces (three) properties that a function $f : \mathbb{M}(n, n) \rightarrow \mathbb{R}$ has to satisfy and then shows that there exists a unique such function, called determinant. Following the first approach the proof of the present theorem can be found on page 252, in Lipschutz (1991) Theorem 7.8, or following the second approach, in Lang (1971), page 128, Theorem 4. ■

Definition 222 The expression used above for the computation of $\det A$ in (5.4) is called “(Laplace) expansion” of the determinant by row i .

The expression used above for the computation of $\det A$ in (5.5) is called “(Laplace) expansion” of the determinant by column j .

Definition 223 Consider a matrix $A_{n \times n}$. Let $1 \leq k \leq n$. A k -th order principal submatrix (minor) of A is the (determinant of the) square submatrix of A obtained deleting $(n - k)$ rows and $(n - k)$ columns in the same position.

Theorem 224 (Properties of determinants)

Let the matrix $A = [a_{ij}] \in \mathbb{M}(n, n)$ be given. Properties presented below hold true even if words “column, columns” are substituted by the words “row, rows”.

1. $\det A = \det A^T$.
2. if two columns are interchanged, the determinant changes its sign,.
3. if there exists $j \in \{1, \dots, n\}$ such that $C^j(A) = \sum_{k=1}^p \beta_k b^k$, then

$$\det \left[C^1(A), \dots, \sum_{k=1}^p \beta_k b^k, \dots, C^n(A) \right] = \sum_{k=1}^p \beta_k \det [C^1(A), \dots, b^k, \dots, C^n(A)],$$

i.e., the determinant of a matrix which has a column equal to the linear combination of some vectors is equal to the linear combination of the determinants of the matrices in which the column under analysis is each of the vector of the initial linear combination, and, therefore, $\forall \beta \in \mathbb{R}$ and $\forall j \in \{1, \dots, n\}$,

$$\det [C^1(A), \dots, \beta C^j(A), \dots, C^n(A)] = \beta \det [C^1(A), \dots, C^j(A), \dots, C^n(A)] = \beta \det A.$$

4. if $\exists j \in \{1, \dots, n\}$ such that $C^j(A) = 0$, then $\det A = 0$, i.e., if a matrix has column equal to zero, then the determinant is zero.
5. if $\exists j, k \in \{1, \dots, n\}$ and $\beta \in \mathbb{R}$ such that $C^j(A) = \beta C^k(A)$, then $\det A = 0$, i.e., the determinant of a matrix with two columns proportional one to the other is zero.
6. If $\exists k \in \{1, \dots, n\}$ and $\beta_1, \dots, \beta_{k-1}, \beta_{k+1}, \dots, \beta_n \in \mathbb{R}$ such that $C^k(A) = \sum_{j \neq k} \beta_j C^j(A)$, then $\det A = 0$, i.e., if a column is equal to a linear combination of the other columns, then $\det A = 0$.
- 7.

$$\det \left[C^1(A), \dots, C^k(A) + \sum_{j \neq k} \beta_j \cdot C^j(A), \dots, C^n(A) \right] = \det A$$

8. $\forall j, j^* \in \{1, \dots, n\}$, $\sum_{i=1}^n a_{ij} \cdot (-1)^{i+j^*} \det A_{ij^*} = 0$, i.e., the sum of the products of the elements of a column times the cofactor of the analogous elements of another column is equal to zero.
9. If A is triangular, $\det A = a_{11} \cdot \dots \cdot a_{22} \cdot \dots \cdot a_{nn}$, i.e., if A is triangular (for example, diagonal), the determinant is the product of the elements on the diagonal.

Proof. 1.

Consider the expansion of the determinant by the first row for the matrix A and the expansion of the determinant by the first column for the matrix A^T .

2.

We proceed by induction. Let A be the starting matrix and A' the matrix with the interchanged columns.

$\mathcal{P}(2)$ is obviously true.

$\mathcal{P}(n-1) \Rightarrow \mathcal{P}(n)$

Expand $\det A$ and $\det A'$ by a column which is not any of the interchanged ones (say column j):

$$\det A = \sum_{i=1}^n (-1)^{i+j} C^{ij}(A) \det A_{ij}$$

$$\det A' = \sum_{i=1}^n (-1)^{i+j} C^{ij}(A) \det A'_{ij}$$

Since $\forall i \in \{1, \dots, n\}$, A_{ij} , $A'_{ij} \in \mathbb{M}(n-1, n-1)$, and they have interchanged column, by the induction argument, $\det A_{ij} = -\det A'_{ij}$, and the desired result follows.

3.

Observe that

$$\sum_{k=1}^p \beta_k b^k = \left(\sum_{k=1}^p \beta_k b_i^k \right)_{i=1}^n.$$

Then,

$$\begin{aligned} \det [C^1(A), \dots, \sum_{k=1}^p \beta_k b^k, \dots, C^n(A)] &= \\ &= \det \left[C^1(A), \dots, \sum_{k=1}^p \beta_k \begin{bmatrix} b_1^k \\ \dots \\ b_i^k \\ \dots \\ b_n^k \end{bmatrix}, \dots, C^n(A) \right] = \\ &= \det \left[C^1(A), \dots, \begin{bmatrix} \sum_{k=1}^p \beta_k b_1^k \\ \dots \\ \sum_{k=1}^p \beta_k b_i^k \\ \dots \\ \sum_{k=1}^p \beta_k b_n^k \end{bmatrix}, \dots, C^n(A) \right] = \\ &= \sum_{i=1}^n (-1)^{i+j} \left(\sum_{k=1}^p \beta_k b_i^k \right) \det A_{ij} = \\ &= \sum_{k=1}^p \beta_k \sum_{i=1}^n (-1)^{i+j} b_i^k \det A_{ij} = \sum_{k=1}^p \beta_k \det [C^1(A), \dots, b^k, \dots, C^n(A)]. \end{aligned}$$

4.

It is sufficient to expand the determinant by the column equal to zero.

5.

Let $A := [C^1(A), \beta C^1(A), C^3(A), \dots, C^n(A)]$ and $\tilde{A} := [C^1(A), C^1(A), C^3(A), \dots, C^n(A)]$ be given. Then $\det A = \beta \det [C^1(A), C^1(A), C^3(A), \dots, C^n(A)] = \beta \det \tilde{A}$. Interchanging the first column with the second column of the matrix \tilde{A} , from property 2, we have that $\det \tilde{A} = -\det \tilde{A}$ and therefore $\det \tilde{A} = 0$, and $\det A = \beta \det \tilde{A} = 0$.

6.

It follows from 3 and 5.

7.

It follows from 3 and 6.

8.

It follows from the fact that the obtained expression is the determinant of a matrix with two equal columns.

9.

It can be shown by induction and expanding the determinant by the first row or column, choosing one which has all the elements equal to zero excluding at most the first element. In other words, in the case of an upper triangular matrix, we can say what follows.

$$\det \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ 0 & a_{22} & a_{23} & \dots & a_{2n} \\ 0 & 0 & a_{33} & \dots & a_{3n} \\ \dots & & & \ddots & \\ 0 & \dots & 0 & \dots & a_{nn} \end{bmatrix} = a_{11} \det \begin{bmatrix} a_{22} & a_{23} & \dots & a_{2n} \\ 0 & a_{33} & \dots & a_{3n} \\ & & \ddots & \\ \dots & 0 & \dots & a_{nn} \end{bmatrix} = a_{11} \cdot a_{22} \cdot a_{33} \cdot \dots a_{nn}.$$

■

Theorem 225 For any $A, B \in \mathbb{M}(n, n)$, $\det(AB) = \det A \cdot \det B$.

Proof. Exercise. ■

Definition 226 $A \in \mathbb{M}(n, n)$ is called nonsingular if $\det A \neq 0$.

5.2 Rank of a matrix

Definition 227 Given $A \in \mathbb{M}(m, n)$, a square submatrix of A of order $k \leq \min\{m, n\}$ is a matrix obtained considering the elements belonging to k rows and k columns of A .

Definition 228 Given $A \in \mathbb{M}(m, n)$, the rank of A is the greatest order of square nonsingular submatrices of A .

Remark 229 $\text{rank } A \leq \min\{m, n\}$.

To compute $\text{rank } A$, with $A \in \mathbb{M}(m, n)$, we can proceed as follows.

1. Consider $k = \min\{m, n\}$, and the set of square submatrices of A of order k . If there exists a nonsingular matrix among them, then $\text{rank } A = k$. If all the square submatrices of A of order k are singular, go to step 2 below.

2. Consider $k - 1$, and then the set of the square submatrices of A of order $k - 1$. If there exists a nonsingular matrix among them, then $\text{rank } A = k - 1$. If all square submatrices of order $k - 1$ are singular, go to step 3.

3. Consider $k - 2 \dots$
and so on.

Remark 230 1. $\text{rank } I_n = n$.

2. The rank of a matrix with a zero row or column is equal to the rank of that matrix without that row or column, i.e.,

$$\text{rank} \begin{bmatrix} A \\ 0 \end{bmatrix} = \text{rank} \begin{bmatrix} A & 0 \end{bmatrix} = \text{rank} \begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix} = \text{rank } A$$

That result follows from the fact that the determinant of any square submatrix of A involving that zero row or columns is zero.

3. From the above results, we also have that

$$\text{rank} \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} = r$$

We now describe an easier way to compute the rank of A , which in fact involves elementary row and column operations we studied in Chapter 1.

Proposition 231 Given $A, A' \in \mathbb{M}(m, n)$,

$$\langle A \text{ is equivalent to } A' \rangle \Leftrightarrow \langle \text{rank } A = \text{rank } A' \rangle$$

Proof. $[\Rightarrow]$ Since A is equivalent to A' , it is possible to go from A to A' through a finite number of elementary row or column operations. In each step, in any square submatrix A^* of A which has been changed accordingly to those operations, the elementary row or column operations 1, 2 and 3 (i.e., 1. row or column interchange, 2. row or column scaling and 3. row or column addition) are such that the determinant of A^* remains unchanged or changes its sign (Property 2, Theorem 224), it is multiplied by a nonzero constant (Property 3), remains unchanged (Property 7), respectively.

Therefore, each submatrix A^* whose determinant is different from zero remains with determinant different from zero and any submatrix A^* whose determinant is zero remains with zero determinant.

$[\Leftarrow]$

From Corollary 160.2², we have that there exist unique \hat{r} and \hat{r}' such that

$$A \text{ is equivalent to } \begin{bmatrix} I_{\hat{r}} & 0 \\ 0 & 0 \end{bmatrix}$$

²That results says what follows:

For any $A \in \mathbb{M}(m, n)$, there exist invertible matrices $P \in \mathbb{M}(m, m)$ and $Q \in \mathbb{M}(n, n)$ and a unique number $r \in \{0, 1, \dots, \min\{m, n\}\}$ such that A is equivalent to

$$PAQ = \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}$$

and

$$A' \text{ is equivalent to } \begin{bmatrix} I_{\widehat{r}'} & 0 \\ 0 & 0 \end{bmatrix}.$$

Moreover, from $[\Rightarrow]$ part of the present proposition, and Remark 230

$$\text{rank} A = \text{rank} \begin{bmatrix} I_{\widehat{r}} & 0 \\ 0 & 0 \end{bmatrix} = \widehat{r}$$

and

$$\text{rank} A' = \text{rank} \begin{bmatrix} I_{\widehat{r}'} & 0 \\ 0 & 0 \end{bmatrix} = \widehat{r}'.$$

Then, by assumption, $\widehat{r} = \widehat{r}' := r$, and A and A' are equivalent to

$$\begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}$$

and therefore A is equivalent to A' . ■

Example 232 *Given*

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 5 & 7 \end{bmatrix}$$

we can perform the following elementary rows and column operations, and cancellation of zero row and columns on the matrix:

$$\begin{aligned} & \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 3 & 5 & 7 \end{bmatrix}, \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \\ 0 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 4 \end{bmatrix}, \begin{bmatrix} 1 & 1 & 3 \\ 2 & 1 & 4 \end{bmatrix}, \\ & \begin{bmatrix} 1 & 1 & 0 \\ 2 & 1 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}. \end{aligned}$$

Therefore, the rank of the matrix is 2.

5.3 Inverse matrices (continued)

Using the notion of determinant, we can find another way of analyzing the problems of i. existence and ii. computation of the inverse matrix. To do that, we introduce the concept of adjoint matrix.

Definition 233 *Given a matrix $A_{n \times n}$, we call adjoint matrix of A , and we denote it by $\text{Adj } A$, the matrix whose elements are the cofactors³ of the corresponding elements of A^T .*

Remark 234 *In other words to construct $\text{Adj } A$,*

1. *construct A^T ,*
2. *consider the cofactors of each element of A^T .*

Example 235

$$A = \begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix}, \quad A^T = \begin{bmatrix} 1 & 0 \\ 2 & 2 \end{bmatrix}, \quad \text{Adj } A = \begin{bmatrix} 2 & -2 \\ 0 & 1 \end{bmatrix}$$

Observe that

$$\begin{bmatrix} 1 & 2 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 2 & -2 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} = (\det A) \cdot I.$$

³From Definition 220, recall that given $A = [a_{ij}] \in \mathbb{M}(m, m)$, $\forall i, j$,

$$(-1)^{i+j} \det A_{ij}$$

is called cofactor of a_{ij} in A .

Example 236

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 1 & 2 & 0 \end{bmatrix}, \quad A^T = \begin{bmatrix} 1 & 0 & 1 \\ 2 & 1 & 2 \\ 3 & 2 & 0 \end{bmatrix}, \quad \text{Adj } A = \begin{bmatrix} -4 & 6 & 1 \\ 2 & -3 & -2 \\ -1 & 0 & 1 \end{bmatrix}$$

Proposition 237 Given $A_{n \times n}$, we have

$$A \cdot \text{Adj } A = \text{Adj } A \cdot A = \det A \cdot I \quad (5.6)$$

Proof. Making the product $A \cdot \text{Adj } A := B$, we have

1. $\forall i \in \{1, \dots, n\}$, the i -th element on the diagonal of B is the expansion of the determinant by the i -th row and therefore is equal to $\det A$.

2. any element not on the diagonal of B is the product of the elements of a row times the corresponding cofactor a parallel row and it is therefore equal to zero due to Property 8 of the determinants stated in Theorem 224). ■

Example 238

$$\det \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 1 & 2 & 0 \end{bmatrix} = -3$$

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 1 & 2 & 0 \end{bmatrix} \begin{bmatrix} -4 & 6 & 1 \\ 2 & -3 & -2 \\ -1 & 0 & 1 \end{bmatrix} = \begin{bmatrix} -3 & 0 & 0 \\ 0 & -3 & 0 \\ 0 & 0 & -3 \end{bmatrix} = -3 \cdot I$$

Proposition 239 Given an $n \times n$ matrix A , the following statements are equivalent:

1. $\det A \neq 0$, i.e., A is nonsingular;
2. A^{-1} exists, i.e., A is invertible;
3. $\text{rank } A = n$;
4. $\text{row rank } A = n$;
5. the column vectors of the matrix A are linearly independent;
6. $\text{col rank } A = n$;
7. the row vectors of the matrix A are linearly independent;

Proof. $1 \Rightarrow 2$

From (5.6) and from the fact that $\det A \neq 0$, we have

$$A \cdot \frac{\text{Adj } A}{\det A} = \frac{\text{Adj } A}{\det A} \cdot A = I$$

and therefore

$$A^{-1} = \frac{\text{Adj } A}{\det A} \quad (5.7)$$

$1 \Leftarrow 2$

$$AA^{-1} = I \Rightarrow \det(AA^{-1}) = \det I \Rightarrow \det A \cdot \det A^{-1} = 1 \Rightarrow \det A \neq 0 \text{ (and } \det A^{-1} \neq 0).$$

$1 \Leftarrow 3$

It follows from the definition of rank and the fact that A is $n \times n$ matrix.

$2 \Rightarrow 4$

From (4.10), it suffices to show that $\langle Ax = 0 \Rightarrow x = 0 \rangle$. Since A^{-1} exists, $Ax = 0 \Rightarrow A^{-1}Ax = A^{-1}0 \Rightarrow x = 0$.

$2 \Leftarrow 4$

From Proposition 199.2, the n linearly independent column vectors $[C^1(A), \dots, C_i(A), \dots, C^n(A)]$ are a basis of \mathbb{R}^n . Therefore, each vector in \mathbb{R}^n is equal to a linear combination of those vectors.

Then $\forall k \in \{1, \dots, n\} \exists b^k \in \mathbb{R}^n$ such that the k -th vector e^k in the canonical basis is equal to $[C^1(A), \dots, C_i(A), \dots, C^n(A)] \cdot b^k = Ab^k$, i.e.,

$$\begin{bmatrix} e^1 & \dots & e^k & \dots & e^n \end{bmatrix} = \begin{bmatrix} Ab^1 & \dots & Ab^k & \dots & Ab^n \end{bmatrix}$$

or, from (3.4) in Remark 70, defined

$$B := \begin{bmatrix} b^1 & \dots & b^k & \dots & b^n \end{bmatrix}$$

$$I = AB$$

i.e., A^{-1} exists (and it is equal to B).

The remaining equivalences follow from Corollary 214. ■

Remark 240 From the proof of the previous Proposition, we also have that, if $\det A \neq 0$, then $\det A^{-1} = (\det A)^{-1}$.

Remark 241 The previous theorem gives a way to compute the inverse matrix as explained in (5.7).

Example 242 1.

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 2 \\ 1 & 2 & 0 \end{bmatrix}^{-1} = -\frac{1}{3} \begin{bmatrix} -4 & 6 & 1 \\ 2 & -3 & -2 \\ -1 & 0 & 1 \end{bmatrix}$$

2.

$$\begin{bmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}^{-1} = \begin{bmatrix} -1 & 0 & 1 \\ 1 & 1 & -1 \\ 0 & -1 & 1 \end{bmatrix}$$

3.

$$\begin{bmatrix} 0 & 1 & 2 \\ 3 & 4 & 5 \\ 6 & 7 & 8 \end{bmatrix}^{-1} \text{ does not exist because } \det \begin{bmatrix} 0 & 1 & 2 \\ 3 & 4 & 5 \\ 6 & 7 & 8 \end{bmatrix} = 0$$

4.

$$\begin{bmatrix} 0 & 1 & 0 \\ 2 & 0 & 2 \\ 1 & 2 & 3 \end{bmatrix}^{-1} = \begin{bmatrix} 1 & \frac{3}{4} & -\frac{1}{2} \\ 1 & 0 & 0 \\ -1 & -\frac{1}{4} & \frac{1}{2} \end{bmatrix}$$

5.

$$\begin{bmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{bmatrix}^{-1} = \begin{bmatrix} \frac{1}{a} & 0 & 0 \\ 0 & \frac{1}{b} & 0 \\ 0 & 0 & \frac{1}{c} \end{bmatrix}$$

if $a, b, c \neq 0$.

5.4 Span of a matrix, linearly independent rows and columns, rank

Proposition 243 Given $A \in \mathbb{M}(m, n)$, then

$$\text{row rank } A = \text{rank } A.$$

Proof. First proof.

The following result which is the content of Corollary 160.5 plays a crucial role in the proof:

For any $A \in \mathbb{M}(m, n)$, there exist invertible matrices $P \in \mathbb{M}(m, m)$ and $Q \in \mathbb{M}(n, n)$ and a unique number $r \in \{0, 1, \dots, \min\{m, n\}\}$ such that

$$A \text{ is equivalent to } PAQ = \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}.$$

From the above result, Proposition 231 and Remark 230 , we have that

$$\text{rank } A = \text{rank} \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} = r. \quad (5.8)$$

From Propositions 105 and 154, we have

$$\text{row span } A = \text{row span } PA;$$

from Propositions 119 and 154, we have

$$\text{col span } PA = \text{col span } PAQ = \text{col span} \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}.$$

From Corollary 214,

$$\dim \text{row span } PA = \dim \text{col span } PA.$$

Therefore

$$\dim \text{row span } A = \dim \text{col span} \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} = r, \quad (5.9)$$

where the last equality follows simply because

$$\text{col span} \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} = \text{col span} \begin{bmatrix} I_r \\ 0 \end{bmatrix},$$

and the r column vectors of the matrix $\begin{bmatrix} I_r \\ 0 \end{bmatrix}$ are linearly independent and therefore, from Proposition ?? , they are a basis of $\text{span} \begin{bmatrix} I_r \\ 0 \end{bmatrix}$.

From (5.8) and (5.9), the desired result follows.

Second proof.

We are going to show that $\text{row rank } A = \text{rank } A$.

Recall that

$$\text{row rank } A := \left\langle \begin{array}{l} r \in \mathbb{N} \text{ such that} \\ \text{i. } r \text{ row vectors of } A \text{ are linearly independent,} \\ \text{ii. if } m > r, \text{ any set of rows of } A \text{ of cardinality } > r \text{ is linearly dependent.} \end{array} \right\rangle$$

We ant to show that

1. if $\text{row rank } A = r$, then $\text{rank } A = r$, and
2. if $\text{rank } A = r$, then $\text{row rank } A = r$.

1.

Consider the r l.i. row vectors of A . Since r is the maximal number of l.i. row vectors, from Lemma 209, each of the remaining $(m - r)$ row vectors is a linear combination of the r l.i. ones. Then, up to reordering of the rows of A , which do not change either $\text{row rank } A$ or $\text{rank } A$, there exist matrices $A_1 \in \mathbb{M}(r, n)$ and $A_2 \in \mathbb{M}(m - r, n)$ such that

$$\text{rank } A = \text{rank} \begin{bmatrix} A_1 \\ A_2 \end{bmatrix} = \text{rank } A_1$$

where the last equality follows from Proposition 231. Then r is the maximum number of l.i. row vectors of A_1 and therefore, from Proposition 212, the maximum number of l.i. column vectors of A_1 . Then, again from Lemma 209, we have that there exist $A_{11} \in \mathbb{M}(r, r)$ and $A_{12} \in \mathbb{M}(r, n - r)$ such that

$$\text{rank } A_1 = \text{rank} \begin{bmatrix} A_{11} & A_{12} \end{bmatrix} = \text{rank } A_{11}$$

Then the square $r \times r$ matrix A_{11} contains r linearly independent vectors. Then from Proposition 239, the result follows.

2.

By Assumption, up to reordering of rows, which do not change either row rank A or rank A ,

$$A = \begin{bmatrix} & s & r & n-r-s \\ r & A_{11} & A_{12} & A_{13} \\ m-r & A_{21} & A_{22} & A_{23} \end{bmatrix}$$

with

$$\text{rank } A_{11} = r.$$

Then from Proposition 239, row, and column, vectors of A_{12} are linearly independent. Then from Corollary 185, the r row vectors of

$$\begin{bmatrix} A_{11} & A_{12} & A_{13} \end{bmatrix}$$

are l.i.

Now suppose that the maximum number of l.i. row vectors of A are $r' > r$ (and the other $m-r'$ row vectors of A are linear combinations of them). Then from part 1 of the present proof, $\text{rank } A = r' > r$, contradicting the assumption. ■

Remark 244 From Corollary 214 and the above Proposition, we have for any matrix $A_{m \times n}$, the following numbers are equal.

1. $\text{rank } A := \text{greatest order of square nonsingular submatrices of } A$,
2. $\text{row rank } A := \dim \text{row span } A$,
3. $\max \text{ number of linear independent rows of } A$,
4. $\text{col rank } A := \dim \text{col span } A$,
5. $\max \text{ number of linear independent columns of } A$.

5.5 Exercises

Problem sets: 9,10

From Lipschutz (1991), starting from page 115:

4.13, 4.14;

starting from page 258:

7.1 \rightarrow 7.10, 7.14 \rightarrow 7.16, 7.44, 7.48.

Chapter 6

Linear functions

6.1 Definition

Definition 245 Given the vector spaces V and U over the same field F , a function $l : V \rightarrow U$ is linear if

1. $\forall v, w \in V, l(v + w) = l(v) + l(w)$, and
2. $\forall \alpha \in F, \forall v \in V, l(\alpha v) = \alpha l(v)$.

Call $L(V, U)$ the set of all such functions. Any time we write $L(V, U)$, we implicitly assume that V and U are vector spaces on the same field F .

In other words, l is linear if it “preserves” the two basic operations of vector spaces.

Remark 246 1. $l \in L(V, U)$ iff $\forall v, w \in V$ and $\forall \alpha, \beta \in F, l(\alpha v + \beta w) = \alpha l(v) + \beta l(w)$;
 2. If $l \in L(V, U)$, then $l(0) = 0$: for arbitrary $x \in V, l(0) = l(0x) = 0l(x) = 0$.

Example 247 Let V and U be vector spaces. The following functions are linear.

1. (identity function)

$$l_1 : V \rightarrow V, \quad l_1(v) = v.$$

2. (null function)

$$l_2 : V \rightarrow U, \quad l_2(v) = 0.$$

- 3.

$$\forall a \in F, \quad l_a : V \rightarrow V, \quad l_a(v) = av.$$

4. (projection function)

$$\forall n \in \mathbb{N}, \forall k \in \mathbb{N},$$

$$\text{proj}_{n+k, n} : \mathbb{R}^{n+k} \rightarrow \mathbb{R}^n, \quad \text{proj}_{n+k, n} : (x_i)_{i=1}^{n+k} \mapsto (x_i)_{i=1}^n;$$

5. (immersion function)

$$\forall n \in \mathbb{N}, \forall k \in \mathbb{N},$$

$$i_{n, n+k} : \mathbb{R}^n \rightarrow \mathbb{R}^{n+k}, \quad i_{n, n+k} : (x_i)_{i=1}^n \mapsto ((x_i)_{i=1}^n, 0) \quad \text{with } 0 \in \mathbb{R}^k.$$

Example 248 Taken $A \in \mathbb{M}(m, n)$, then

$$l : \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad l(x) = Ax$$

is a linear function, as shown in part 3 in Remark 76.

Remark 249 Let V and U be vector spaces. If $l \in L(V, U)$, then

$$\text{graph } l := \{(v, u) \in V \times U : u = l(v)\}$$

is a vector subspace of $V \times U$.

Example 250 Let V be the vector space of polynomials in the variable t . The following functions are linear

1. The derivative defines a function $\mathbb{D} : V \rightarrow V$ as

$$\mathbb{D} : p \mapsto p'$$

where p' is the derivative function of p .

2. The definite integral from 0 to 1 defines a function $i : V \rightarrow \mathbb{R}$ as

$$i : p \mapsto \int_0^1 p(t) dt.$$

Proposition 251 If $l \in L(V, U)$ is invertible, then its inverse l^{-1} is linear.

Proof. Take arbitrary $u, u' \in U$ and $\alpha, \beta \in F$. Then, since l is onto, there exist $v, v' \in V$ such that

$$l(v) = u \quad \text{and} \quad l(v') = u'$$

and by definition of inverse function

$$l^{-1}(u) = v \quad \text{and} \quad l^{-1}(u') = v'.$$

Then

$$\alpha u + \beta u' = \alpha l(v) + \beta l(v') = l(\alpha v + \beta v')$$

where last equality comes from the linearity of l . Then again by definition of inverse,

$$l^{-1}(\alpha u + \beta u') = \alpha v + \beta v' = \alpha l^{-1}(u) + \beta l^{-1}(u').$$

■

6.2 Kernel and Image of a linear function

Definition 252 Assume that $l \in L(V, U)$. The kernel of l , denoted by $\ker l$ is the set

$$\{v \in V : l(v) = 0\} = l^{-1}(0).$$

The Image of l , denoted by $\text{Im } l$ is the set

$$\{u \in U : \exists v \in V \text{ such that } f(v) = u\} = l(V).$$

Proposition 253 Given $l \in L(V, U)$, then

1. $\ker l$ is a vector subspace of V , and
2. $\text{Im } l$ is a vector subspace of U .

Proof. 1.

Since $l(0) = 0$, then $0 \in \ker l$.

Take $v^1, v^2 \in \ker l$ and $\alpha, \beta \in F$. Then

$$l(\alpha v^1 + \beta v^2) = \alpha l(v^1) + \beta l(v^2) = 0$$

i.e., $\alpha v^1 + \beta v^2 \in \ker l$.

2.

Since $0 \in V$ and $l(0) = 0$, then $0 \in \text{Im } l$.

Take $w^1, w^2 \in \text{Im } l$ and $\alpha, \beta \in F$. Then for $i \in \{1, 2\}$, $\exists v^i \in V$ such that $l(v^i) = w_i$. Moreover,

$$\alpha w^1 + \beta w^2 = \alpha l(v^1) + \beta l(v^2) = l(\alpha v^1 + \beta v^2)$$

i.e., $\alpha w^1 + \beta w^2 \in \text{Im } l$. ■

Proposition 254 If $\text{span}(\{v^1, \dots, v^n\}) = V$ and $l \in L(V, U)$, then $\text{span}(\{l(v^i)\}_{i=1}^n) = \text{Im } l$.

Proof. Taken $u \in \text{Im } l$, there exists $v \in V$ such that $l(v) = u$. Moreover, $\exists (\alpha_i)_{i=1}^n \in \mathbb{R}^n$ such that $v = \sum_{i=1}^n \alpha_i v^i$. Then

$$u = l(v) = l\left(\sum_{i=1}^n \alpha_i v^i\right) = \sum_{i=1}^n \alpha_i l(v^i),$$

as desired. ■

Remark 255 From the previous proposition, we have that if $\{v^1, \dots, v^n\}$ is a basis of V , then

$$n \geq \dim \text{span}\left(\{l(v^i)\}_{i=1}^n\right) = \dim \text{Im } l.$$

Example 256 Let V the vector space of polynomials and $\mathbb{D}^3 : V \rightarrow V$, $p \mapsto p'''$, i.e., the third derivative of p . Then

$$\ker \mathbb{D}^3 = \text{set of polynomials of degree } \leq 2,$$

since $\mathbb{D}^3(at^2 + bt + c) = 0$ and $\mathbb{D}^3(t^n) \neq 0$ for $n > 2$. Moreover,

$$\text{Im } \mathbb{D}^3 = V,$$

since every polynomial is the third derivative of some polynomial.

Proposition 257 (Dimension Theorem) If V is a finite dimensional vector space and $l \in L(V, U)$, then

$$\dim V = \dim \ker l + \dim \text{Im } l$$

Proof. (Idea of the proof.

1. Using a basis of $\ker l$ (with n_1 elements) and a basis of $\text{Im } l$ (with n_2 elements) construct a set with $n_1 + n_2$ elements which generates V .

2. Show that set is linearly independent (by contradiction), and therefore a basis of V , and therefore $\dim V = n_1 + n_2$.)

Since $\ker l \subseteq V$ and from Remark 255, $\ker l$ and $\text{Im } l$ have finite dimension. Therefore, we can define $n_1 = \dim \ker l$ and $n_2 = \dim \text{Im } l$.

Take an arbitrary $v \in V$. Let

$$\{w^1, \dots, w^{n_1}\} \text{ be a basis of } \ker l \quad (6.1)$$

and

$$\{u^1, \dots, u^{n_2}\} \text{ be a basis of } \text{Im } l \quad (6.2)$$

Then,

$$\forall i \in \{1, \dots, n_2\}, \exists v^i \in V \text{ such that } u^i = l(v^i) \quad (6.3)$$

From (6.2),

$$\exists c = (c_i)_{i=1}^{n_2} \text{ such that } l(v) = \sum_{i=1}^{n_2} c_i u^i \quad (6.4)$$

Then, from (6.4) and (6.3), we get

$$l(v) = \sum_{i=1}^{n_2} c_i u^i = \sum_{i=1}^{n_2} c_i l(v^i)$$

and from linearity of l

$$0 = l(v) - \sum_{i=1}^{n_2} c_i l(v^i) = l(v) - l\left(\sum_{i=1}^{n_2} c_i v^i\right) = l\left(v - \sum_{i=1}^{n_2} c_i v^i\right)$$

i.e.,

$$v - \sum_{i=1}^{n_2} c_i v^i \in \ker l \quad (6.5)$$

From (6.1),

$$\exists (d_j)_{j=1}^{n_2} \text{ such that } v - \sum_{i=1}^{n_2} c_i v^i = \sum_{j=1}^{n_1} d_j w^j$$

Summarizing, we have

$$\forall v \in V, \exists (c_i)_{i=1}^{n_2} \text{ and } (d_j)_{j=1}^{n_1} \text{ such that } v = \sum_{i=1}^{n_2} c_i v^i + \sum_{j=1}^{n_1} d_j w^j$$

Therefore, we found $n_1 + n_2$ vectors which generate V ; if we show that they are l.i., then, by definition, they are a basis and therefore $n = n_1 + n_2$ as desired.

We want to show that

$$\sum_{i=1}^{n_2} \alpha_i v^i + \sum_{j=1}^{n_1} \beta_j w^j = 0 \Rightarrow ((\alpha_i)_{i=1}^{n_2}, (\beta_j)_{j=1}^{n_1}) = 0 \quad (6.6)$$

Then

$$0 = l \left(\sum_{i=1}^{n_2} \alpha_i v^i + \sum_{j=1}^{n_1} \beta_j w^j \right) = \sum_{i=1}^{n_2} \alpha_i l(v^i) + \sum_{j=1}^{n_1} \beta_j l(w^j)$$

From (6.1), i.e., $\{w^1, \dots, w^{n_1}\}$ is a basis of $\ker l$, and from (6.3), we get

$$\sum_{i=1}^{n_2} \alpha_i u_i = 0$$

From (6.2), i.e., $\{u^1, \dots, u^{n_2}\}$ be a basis of $\text{Im } l$,

$$(\alpha_i)_{i=1}^{n_2} = 0 \quad (6.7)$$

But from the assumption in (6.6) and (6.7) we have that

$$\sum_{j=1}^{n_1} \beta_j w^j = 0$$

and since $\{w^1, \dots, w^{n_1}\}$ is a basis of $\ker l$, we get also that

$$(\beta_j)_{j=1}^{n_1} = 0,$$

as desired. ■

Example 258 Let V and U be vector spaces, with $\dim V = n$.

In 1. and 2. below, we verify the statement of the Dimension Theorem: in 3. and 4., we use that statement.

1. Identity function id_V .

$$\begin{aligned} \dim \text{Im } \text{id}_V &= n \\ \dim \ker l &= 0. \end{aligned}$$

2. Null function $0 \in L(V, U)$

$$\begin{aligned} \dim \text{Im } 0 &= 0 \\ \dim \ker 0 &= n. \end{aligned}$$

3. $l \in L(\mathbb{R}^2, \mathbb{R})$,

$$l((x_1, x_2)) = x_1.$$

Since $\ker l = \{(x_1, x_2) \in \mathbb{R}^2 : x_1 = 0\}$, $\{(0, 1)\}$ is a basis¹ of $\ker l$ and

$$\begin{aligned} \dim \ker l &= 1, \\ \dim \text{Im } l &= 2 - 1 = 1. \end{aligned}$$

¹In Remark 337 we will present an algorithm to compute a basis of $\ker l$.

4. $l \in L(\mathbb{R}^3, \mathbb{R}^2)$,

$$l((x_1, x_2, x_3)) = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}.$$

Defined

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 0 \end{bmatrix},$$

since

$$\text{Im } l = \{y \in \mathbb{R}^2 : \exists x \in \mathbb{R}^3 \text{ such that } Ax = y\} = \text{span col } A = \text{rank } A,$$

and since the first two column vectors of A are linearly independent, we have that

$$\begin{aligned} \dim \text{Im } l &= 2 \\ \dim \ker l &= 3 - 2 = 1. \end{aligned}$$

Exercise 259 Let $l \in \mathcal{L}(\mathbb{R}^3, \mathbb{R}^3)$ such that

$$l \begin{pmatrix} x \\ y \\ z \end{pmatrix} \mapsto \begin{pmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \\ z \end{pmatrix}$$

be given. Find $\ker l$ and $\text{Im } l$ (Answer: $\ker l = \{0\}$; $\text{Im } l = \mathbb{R}^3$).

6.3 Nonsingular functions and isomorphisms

Definition 260 $l \in L(V, U)$ is singular if $\exists v \in V \setminus \{0\}$ such that $l(v) = 0$.

Remark 261 Thus $l \in L(V, U)$ is nonsingular if $\forall v \in V \setminus \{0\}$, $l(v) \neq 0$ i.e., $\ker l = \{0\}$. Briefly,

$$l \in L(V, U) \text{ is nonsingular} \Leftrightarrow \ker l = \{0\}.$$

Remark 262 In Remark 300, we will discuss the relationship between singular matrices and singular linear functions.

Example 263 1. Let $l \in \mathcal{L}(\mathbb{R}^3, \mathbb{R}^3)$ be the projection mapping into the xy plane, i.e.,

$$l : \begin{matrix} \mathbb{R}^3 & \rightarrow & \mathbb{R}^3 \\ \begin{pmatrix} x \\ y \\ z \end{pmatrix} & \mapsto & \begin{pmatrix} x \\ y \\ 0 \end{pmatrix} \end{matrix}$$

Then l is singular, since for any $a \in \mathbb{R}$, $l(0, 0, a) = (0, 0, 0)$.

2. $l \in \mathcal{L}(\mathbb{R}^3, \mathbb{R}^3)$ defined in Example 259 is nonsingular, since $\ker l = \{0\}$.

Proposition 264 If $l \in L(V, U)$ is nonsingular, then the image of any linearly independent set is linearly independent.

Proof. Suppose $\{v^1, \dots, v^n\}$ are linearly independent. We want to show that $\{l(v^1), \dots, l(v^n)\}$ are linearly independent as well. Suppose

$$\sum_{i=1}^n \alpha_i \cdot l(v^i) = 0.$$

Then

$$l \left(\sum_{i=1}^n \alpha_i \cdot v^i \right) = 0.$$

and therefore $(\sum_{i=1}^n \alpha_i \cdot v^i) \in \ker l = \{0\}$, where the last equality comes from the fact that l is nonsingular. Then $\sum_{i=1}^n \alpha_i \cdot v^i = 0$ and, since $\{v^1, \dots, v^n\}$ are linearly independent, $(\alpha_i)_{i=1}^n = 0$, as desired. ■

Definition 265 Let two vector spaces V and U be given. U is isomorphic to V if there exists a function $l \in L(V, U)$ which is one-to-one and onto. l is called an isomorphism from V to U .

Remark 266 By definition of isomorphism, if l is an isomorphism, the l is invertible and therefore, from Proposition 251, l^{-1} is linear.

Remark 267 “To be isomorphic” is an equivalence relation.

Proposition 268 Any n -dimensional vector space V on a field F is isomorphic to F^n .

Proof. Since V and F^n are vector spaces, we are left with showing that there exists an isomorphism between them. Let $\mathbf{v} = \{v^1, \dots, v^n\}$ be a basis of V . Recall that we define

$$cr : V \rightarrow F^n, \quad v \mapsto [v]_{\mathbf{v}},$$

where cr stands for “coordinates”.

1. cr is linear. Given $v, w \in V$, suppose

$$v = \sum_{i=1}^n a_i v^i \quad \text{and} \quad w = \sum_{i=1}^n b_i v^i$$

i.e.,

$$[v]_{\mathbf{v}} = [a_i]_{i=1}^n \quad \text{and} \quad [w]_{\mathbf{v}} = [b_i]_{i=1}^n.$$

$\forall \alpha, \beta \in F$ and $\forall v^1, v^2 \in V$,

$$\alpha v + \beta w = \alpha \sum_{i=1}^n a_i v^i + \beta \sum_{i=1}^n b_i v^i = \sum_{i=1}^n (\alpha a_i + \beta b_i) v^i$$

i.e.,

$$[\alpha v + \beta w]_{\mathbf{v}} = \alpha [a_i]_{i=1}^n + \beta [b_i]_{i=1}^n = \alpha [v]_{\mathbf{v}} + \beta [w]_{\mathbf{v}}.$$

2. cr is onto. $\forall (a)_{i=1}^n \in \mathbb{R}^n$, $cr(\sum_{i=1}^n a_i v^i) = (a)_{i=1}^n$.

3. cr is one-to-one. $cr(v) = cr(w)$ implies that $v = w$, simply because $v = \sum_{i=1}^n cr_i(v) u^i$ and $w = \sum_{i=1}^n cr_i(w) u^i$. ■

Proposition 269 Let V and U be finite dimensional vectors spaces on the same field F such that $S = \{v^1, \dots, v^n\}$ is a basis of V and $\{u^1, \dots, u^n\}$ is a set of arbitrary vectors in U . Then there exists a unique linear function $l : V \rightarrow U$ such that $\forall i \in \{1, \dots, n\}$, $l(v^i) = u^i$.

Proof. The proof goes the following three steps.

1. Define l ;
 2. Show that l is linear;
 3. Show that l is unique.
1. Using the definition of coordinates, $\forall v \in V$, define

$$l : V \rightarrow U, \quad v \mapsto \sum_{i=1}^n [v]_S^i \cdot u^i,$$

where $[v]_S^i$ denotes the i -th component of the vector $[v]_S$. Recall that $e_n^j \in \mathbb{R}^n$ is the j -th element in the canonical basis of \mathbb{R}^n and defined $e_n^j := (e_{n,i}^j)_{i=1}^n$, we have

$$e_{n,i}^j = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j. \end{cases}$$

Then $\forall j \in \{1, \dots, n\}$, $[v^j]_S = e_n^j$ and

$$l(v^j) = \sum_{i=1}^n [v^j]_S^i \cdot u^i = \sum_{i=1}^n e_{n,i}^j \cdot u^i = u^j.$$

2. Let $v, w \in V$ and $\alpha, \beta \in F$. Then

$$l(\alpha v + \beta w) = \sum_{i=1}^n [\alpha v + \beta w]_S^i \cdot u^i = \sum_{i=1}^n \left(\alpha [v]_S^i + \beta [w]_S^i \right) \cdot u^i = \alpha \sum_{i=1}^n [v]_S^i \cdot u^i + \beta \sum_{i=1}^n [w]_S^i \cdot u^i,$$

where the before the last equality follows from the linearity of $[\cdot]$ - see the proof of Proposition 268

3. Suppose $g \in L(V, U)$ and $\forall i \in \{1, \dots, n\}$, $g(v^i) = u^i$. Then $\forall v \in V$,

$$g(v) = g\left(\sum_{i=1}^n [v]_S^i \cdot v^i\right) = \sum_{i=1}^n [v]_S^i \cdot g(v^i) = \sum_{i=1}^n [v]_S^i \cdot u^i = l(v)$$

where the last equality follows from the definition of l . ■

Remark 270 Observe that if V and U are finite(nonzero) dimensional vector spaces, there is a multitude of functions from V to U . The above Proposition says that linear functions are completely determined by what “they do to the elements of a basis” of V .

Example 271 We want to find the unique linear mapping $l : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that

$$l(1, 2) = (2, 3) \quad \text{and} \quad l(0, 1) = (1, 4).$$

Observe that $\mathcal{B} := \{(1, 2), (0, 1)\}$ is a basis of \mathbb{R}^2 . For any $(a, b) \in \mathbb{R}^2$, there exist $x, y \in \mathbb{R}$ such that

$$(a, b) = x(1, 2) + y(0, 1) = (x, 2x + y),$$

i.e., $a = x$ and $b = 2x + y$ and therefore $x = a$ and $y = -2a + b$. Then,

$$l(a, b) = l(x(1, 2) + y(0, 1)) \stackrel{l \text{ linear}}{=} xl(1, 2) + yl(0, 1) = a(2, 3) + (-2a + b)(1, 4) = (b, -5a + 4b),$$

i.e.,

$$l : \begin{matrix} \mathbb{R}^2 & \rightarrow & \mathbb{R}^2 \\ \begin{pmatrix} a \\ b \end{pmatrix} & \mapsto & \begin{pmatrix} b \\ -5a + 4b \end{pmatrix} \end{matrix}$$

Proposition 272 Assume that $l \in L(V, U)$. Then,

$$l \text{ is one-to-one} \Leftrightarrow l \text{ is nonsingular}$$

Proof. $[\Rightarrow]$

Take $v \in \ker l$. Then

$$l(v) = 0 = l(0)$$

where last equality follows from Remark 246. Since l is one-to-one, $v = 0$.

$[\Leftarrow]$ If $l(v) = l(w)$, then $l(v - w) = 0$ and, since l is nonsingular, $v - w = 0$. ■

Proposition 273 Assume that V and U are finite dimensional vector spaces and $l \in L(V, U)$. Then,

1. l is one-to-one $\Rightarrow \dim V \leq \dim U$;
2. l is onto $\Rightarrow \dim V \geq \dim U$;
3. l is invertible $\Rightarrow \dim V = \dim U$.

Proof. The main ingredient in the proof is Proposition 257, i.e., the Dimension Theorem.

1. Since l is one-to-one, from the previous Proposition, $\dim \ker l = 0$. Then, from Proposition 257 (the Dimension Theorem), $\dim V = \dim \text{Im } l$. Since $\text{Im } l$ is a subspace of U , then $\dim \text{Im } l \leq \dim U$.

2. Since l is onto iff $\text{Im } l = U$, from Proposition 257 (the Dimension Theorem), we get

$$\dim V = \dim \ker l + \dim U \geq \dim U.$$

3. l is invertible iff l is one-to-one and onto. ■

Proposition 274 *Let V and U be finite dimensional vector space on the same field F . Then,*

$$U \text{ and } V \text{ are isomorphic} \Leftrightarrow \dim U = \dim V.$$

Proof. $[\Rightarrow]$

It follows from the definition of isomorphism and part 3 in the previous Proposition.

$[\Leftarrow]$

Assume that V and U are vector spaces such that $\dim V = \dim U = n$. Then, from Proposition 268, V and U are isomorphic to F^n and from Remark 267, the result follows. ■

Proposition 275 *Suppose V and U are vector spaces such that $\dim V = \dim U = n$ and $l \in L(V, U)$. Then the following statements are equivalent.*

1. l is nonsingular, i.e., $\ker l = \{0\}$,
2. l is one-to-one,
3. l is onto,
4. l is an isomorphism.

Proof. $[1 \Leftrightarrow 2]$.

It is the content of Proposition 272.

$[1 \Rightarrow 3]$

Since l is nonsingular, then $\ker l = \{0\}$ and $\dim \ker l = 0$. Then, from Proposition 257 (the Dimension Theorem), i.e., $\dim V = \dim \ker l + \dim \operatorname{Im} l$, and the fact $\dim V = \dim U$, we get $\dim U = \dim \operatorname{Im} l$. Since $\operatorname{Im} l \subseteq U$ and U is finite dimensional, from Proposition 202, $\operatorname{Im} l = U$, i.e., l is onto, as desired.

$[3 \Rightarrow 1]$

Since l is onto, $\dim \operatorname{Im} l = \dim V$ and from Proposition 257 (the Dimension Theorem), $\dim V = \dim \ker l + \dim \operatorname{Im} l$, and therefore $\dim \ker l = 0$, i.e., l is nonsingular.

$[1 \Rightarrow 4]$

It follows from the definition of isomorphism and the facts that $[1 \Leftrightarrow 2]$ and $[1 \Rightarrow 3]$.

$[4 \Rightarrow 1]$

It follows from the definition of isomorphism and the facts that $[2 \Leftrightarrow 1]$. ■

Definition 276 *A vector space endomorphism is a linear function from a vector space V into itself.*

6.4 Exercises

From Lipschutz (1991), starting from page 325:

9.3, 9.6, 9.9 \rightarrow 9.11, 9.16 \rightarrow 9.21, 9.26, 9.27, 9.31 \rightarrow 9.35, 9.42 \rightarrow 9.44; observe that Lipschutz denotes $\mathcal{L}(V, U)$ by $\operatorname{Hom}(V, U)$.

Chapter 7

Linear functions and matrices

In Remark 65 we have seen that the set of $m \times n$ matrices with the standard sum and scalar multiplication is a vector space, called $\mathbb{M}(m, n)$. We are going to show that:

1. the set $\mathcal{L}(V, U)$ with naturally defined sum and scalar multiplication is a vector space, called $\mathcal{L}(V, U)$;
2. If $\dim V = n$ and $\dim U = m$, then $\mathcal{L}(V, U)$ and $\mathbb{M}(m, n)$ are isomorphic.

7.1 From a linear function to the associated matrix

Definition 277 Suppose V and U are vector spaces over a field F and $l_1, l_2 \in \mathcal{L}(V, U)$ and $\alpha \in F$.

$$l_1 + l_2 : V \rightarrow U, \quad v \mapsto l_1(v) + l_2(v)$$

$$\alpha l_1 : V \rightarrow U, \quad v \mapsto \alpha l_1(v)$$

Proposition 278 $\mathcal{L}(V, U)$ with the above defined operations is a vector space on F , denoted by $\mathcal{L}(V, U)$.

Proof. Exercise.¹ ■

Proposition 279 Compositions of linear functions are linear.

Proof. Suppose V, U, W are vector spaces over a field F , $l_1 \in \mathcal{L}(V, U)$ and $l_2 \in \mathcal{L}(U, W)$. We want to show that $l := l_2 \circ l_1 \in \mathcal{L}(V, W)$. Indeed, for any $\alpha_1, \alpha_2 \in F$ and for any $v^1, v^2 \in V$, we have that

$$\begin{aligned} l(\alpha_1 v^1 + \alpha_2 v^2) &:= (l_2 \circ l_1)(\alpha_1 v^1 + \alpha_2 v^2) = l_2(l_1(\alpha_1 v^1 + \alpha_2 v^2)) = \\ &= l_2(\alpha_1 l_1(v^1) + \alpha_2 l_1(v^2)) = \alpha_1 l_2(l_1(v^1)) + \alpha_2 l_2(l_1(v^2)) = \alpha_1 l(v^1) + \alpha_2 l(v^2), \end{aligned}$$

as desired. ■

Definition 280 Suppose $l \in \mathcal{L}(V, U)$, $\mathbf{v} = \{v^1, \dots, v^n\}$ is an ordered basis of V , $\mathbf{u} = \{u^1, \dots, u^m\}$ is an ordered basis of U . Then,

$$[l]_{\mathbf{v}}^{\mathbf{u}} := \begin{bmatrix} [l(v^1)]_{\mathbf{u}} & \dots & [l(v^j)]_{\mathbf{u}} & \dots & [l(v^n)]_{\mathbf{u}} \end{bmatrix} \in \mathbb{M}(m, n), \quad (7.1)$$

where for any $j \in \{1, \dots, n\}$, $[l(v^j)]_{\mathbf{u}}$ is a column vector, is called the matrix representation of l relative to the basis \mathbf{v} and \mathbf{u} . In words, $[l]_{\mathbf{v}}^{\mathbf{u}}$ is the matrix whose columns are the coordinates relative to the basis of the codomain of l of the images of each vector in the basis of the domain of l .

Remark 281 Observe that by definition of coordinates, there is a unique matrix representation of a linear function.

¹For a detailed proof see Lipschutz (1989), page 270.

Definition 282 Suppose $l \in \mathcal{L}(V, U)$, $\mathbf{v} = \{v^1, \dots, v^n\}$ is a ordered basis of V , $\mathbf{u} = \{u^1, \dots, u^m\}$ is a ordered basis of U .

$$\varphi_{\mathbf{v}}^{\mathbf{u}} : \mathcal{L}(V, U) \rightarrow \mathbb{M}(m, n), \quad l \mapsto [l]_{\mathbf{v}}^{\mathbf{u}} \text{ defined in (7.1)}.$$

If no confusion may arise, we will denote $\varphi_{\mathbf{v}}^{\mathbf{u}}$ simply by φ .

Example 283 Given $\text{id}_V \in \mathcal{L}(V, V)$ and a ordered basis $\mathbf{v} = \{v^1, \dots, v^n\}$ of V , then $[l]_{\mathbf{v}}^{\mathbf{v}} = I_n$.

The proposition below shows that multiplying the coordinate vector of v relative to the ordered basis \mathbf{v} by the matrix $[l]_{\mathbf{v}}^{\mathbf{u}}$, we get the coordinate vector of $l(v)$ relative to the ordered basis \mathbf{u} .

Proposition 284 $\forall v \in V$,

$$[l]_{\mathbf{v}}^{\mathbf{u}} \cdot [v]_{\mathbf{v}} = [l(v)]_{\mathbf{u}} \quad (7.2)$$

Proof. Assume $v \in V$. First of all observe that

$$[l]_{\mathbf{v}}^{\mathbf{u}} \cdot [v]_{\mathbf{v}} = \begin{bmatrix} [l(v^1)]_{\mathbf{u}} & \dots & [l(v^j)]_{\mathbf{u}} & \dots & [l(v^n)]_{\mathbf{u}} \end{bmatrix} \begin{bmatrix} [v]_{\mathbf{v}}^1 \\ \dots \\ [v]_{\mathbf{v}}^j \\ \dots \\ [v]_{\mathbf{v}}^n \end{bmatrix} = \sum_{j=1}^n [v]_{\mathbf{v}}^j \cdot [l(v^j)]_{\mathbf{u}}.$$

Moreover, from the linearity of the function $cr_{\mathbf{u}} := [\cdot]_{\mathbf{u}}$, and using the fact that the composition of linear functions is a continuous function, we get:

$$[l(v)]_{\mathbf{u}} = cr_{\mathbf{u}}(l(v)) = (cr_{\mathbf{u}} \circ l) \left(\sum_{j=1}^n [v]_{\mathbf{v}}^j \cdot v^j \right) = \sum_{j=1}^n [v]_{\mathbf{v}}^j \cdot (cr_{\mathbf{u}} \circ l)(v^j) = \sum_{j=1}^n [v]_{\mathbf{v}}^j \cdot [l(v^j)]_{\mathbf{u}}.$$

■

Example 285 Let's verify equality (7.2) in the case in which

a.

$$l : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad (x_1, x_2) \mapsto \begin{pmatrix} x_1 + x_2 \\ x_1 - x_2 \end{pmatrix}$$

b. the ordered basis \mathbf{v} of the domain of l is

$$\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\},$$

c. the ordered basis \mathbf{u} of the codomain of l is

$$\left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\},$$

d.

$$v = \begin{pmatrix} 3 \\ 4 \end{pmatrix}.$$

The main needed computations are presented below.

$$[l]_{\mathbf{v}}^{\mathbf{u}} := \left[\left[l \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right]_{\mathbf{u}}, \left[l \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right]_{\mathbf{u}} \right] = \left[\left[\begin{pmatrix} 1 \\ 1 \end{pmatrix} \right]_{\mathbf{u}}, \left[\begin{pmatrix} 1 \\ -1 \end{pmatrix} \right]_{\mathbf{u}} \right] = \begin{bmatrix} 1 & -3 \\ 0 & 2 \end{bmatrix},$$

$$[v]_{\mathbf{v}} = \begin{bmatrix} 3 \\ 4 \end{bmatrix},$$

$$[l(v)]_{\mathbf{u}} = \left[\begin{pmatrix} 7 \\ -1 \end{pmatrix} \right]_{\mathbf{u}} = \begin{bmatrix} -9 \\ 8 \end{bmatrix},$$

$$[l]_{\mathbf{v}}^{\mathbf{u}} \cdot [v]_{\mathbf{v}} = \begin{bmatrix} 1 & -3 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \begin{bmatrix} -9 \\ 8 \end{bmatrix}.$$

7.2 From a matrix to the associated linear function

Given $A \in \mathbb{M}(m, n)$, recall that $\forall i \in \{1, \dots, m\}$, $R^i(A)$ denotes the i -th row vector of A , i.e.,

$$A = \begin{bmatrix} R^1(A) \\ \vdots \\ R^i(A) \\ \vdots \\ R^m(A) \end{bmatrix}_{m \times n}$$

Definition 286 Consider vector spaces V and U with ordered basis $\mathbf{v} = \{v^1, \dots, v^n\}$ and $\mathbf{u} = \{u^1, \dots, u^m\}$, respectively. Given $A \in \mathbb{M}(m, n)$, define

$$l_{A, \mathbf{v}}^{\mathbf{u}} : V \rightarrow U, \quad v \mapsto \sum_{i=1}^m (R^i(A) \cdot [v]_{\mathbf{v}}) \cdot u^i.$$

Example 287 Take $U = V = \mathbb{R}^2$, $\mathbf{v} = \mathcal{E}_2$, $\mathbf{u} = \left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\}$ and $A = \begin{bmatrix} 1 & -3 \\ 0 & 2 \end{bmatrix}$. Then,

$$\begin{aligned} l_{A, \mathbf{v}}^{\mathbf{u}}(x_1, x_2) &:= \sum_{i=1}^2 (R^i(A) \cdot [v]_{\mathcal{E}_2}) \cdot u^i = \left(\begin{bmatrix} 1 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \right) \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \left(\begin{bmatrix} 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \right) \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \\ &= (x_1 - 3x_2) \begin{pmatrix} 1 \\ 1 \end{pmatrix} + 2x_2 \begin{pmatrix} 2 \\ 1 \end{pmatrix} = \begin{pmatrix} x_1 - 3x_2 + 4x_2 \\ x_1 - 3x_2 + 2x_2 \end{pmatrix} = \begin{pmatrix} x_1 + x_2 \\ x_1 - x_2 \end{pmatrix} \end{aligned}$$

Proposition 288 $l_{A, \mathbf{v}}^{\mathbf{u}}$ defined above is linear, i.e., $l_{A, \mathbf{v}}^{\mathbf{u}} \in \mathcal{L}(V, U)$.

Proof. $\forall \alpha, \beta \in \mathbb{R}$ and $\forall v^1, v^2 \in V$,

$$\begin{aligned} l_{A, \mathbf{v}}^{\mathbf{u}}(\alpha v^1 + \beta v^2) &= \sum_{i=1}^m R^i(A) \cdot [\alpha v^1 + \beta v^2]_{\mathbf{v}} \cdot u^i = \sum_{i=1}^m R^i(A) \cdot (\alpha [v^1]_{\mathbf{v}} + \beta [v^2]_{\mathbf{v}}) \cdot u^i = \\ &= \alpha \sum_{i=1}^m R^i(A) \cdot [v^1]_{\mathbf{v}} \cdot u^i + \beta \sum_{i=1}^m R^i(A) \cdot [v^2]_{\mathbf{v}} \cdot u^i = \alpha l_{A, \mathbf{v}}^{\mathbf{u}}(v^1) + \beta l_{A, \mathbf{v}}^{\mathbf{u}}(v^2). \end{aligned}$$

where the second equality follows from the proof of Proposition 268. ■

Definition 289 Given the vector spaces V and U with ordered basis $\mathbf{v} = \{v^1, \dots, v^n\}$ and $\mathbf{u} = \{u^1, \dots, u^m\}$, respectively, define

$$\psi_{\mathbf{v}}^{\mathbf{u}} : \mathbb{M}(m, n) \rightarrow \mathcal{L}(V, U) \quad : A \mapsto l_{A, \mathbf{v}}^{\mathbf{u}}.$$

If no confusion may arise, we will denote $\psi_{\mathbf{v}}^{\mathbf{u}}$ simply by ψ .

Proposition 290 ψ defined above is linear.

Proof. We want to show that $\forall \alpha, \beta \in \mathbb{R}$ and $\forall A, B \in \mathbb{M}(m, n)$,

$$\psi(\alpha A + \beta B) = \alpha \psi(A) + \beta \psi(B)$$

i.e.,

$$l_{\alpha A + \beta B, \mathbf{u}}^{\mathbf{v}} = \alpha l_{A, \mathbf{v}}^{\mathbf{u}} + \beta l_{B, \mathbf{u}}^{\mathbf{v}}$$

i.e., $\forall v \in V$,

$$l_{\alpha A + \beta B, \mathbf{u}}^{\mathbf{v}}(v) = \alpha l_{A, \mathbf{v}}^{\mathbf{u}}(v) + \beta l_{B, \mathbf{u}}^{\mathbf{v}}(v).$$

Now,

$$\begin{aligned} l_{\alpha A + \beta B, \mathbf{u}}^{\mathbf{v}}(v) &= \sum_{i=1}^m (\alpha \cdot R^i(A) + \beta \cdot R^i(B)) \cdot [v]_{\mathbf{v}} \cdot u^i = \\ &= \alpha \sum_{i=1}^m R^i(A) \cdot [v]_{\mathbf{v}} \cdot u^i + \beta \sum_{i=1}^m R^i(B) \cdot [v]_{\mathbf{v}} \cdot u^i = \alpha l_{A, \mathbf{v}}^{\mathbf{u}}(v) + \beta l_{B, \mathbf{u}}^{\mathbf{v}}(v), \end{aligned}$$

where the first equality come from Definition 286. ■

7.3 $\mathbb{M}(m, n)$ and $\mathcal{L}(V, U)$ are isomorphic

Proposition 291 *Given the vector space V and U with dimension n and m , respectively,*

$$\mathbb{M}(m, n) \text{ and } \mathcal{L}(V, U) \text{ are isomorphic,}$$

and

$$\dim \mathcal{L}(V, U) = mn.$$

Proof. Linearity of the two spaces was proved above. We want now to check that ψ presented in Definition 289 is an isomorphism, i.e., ψ is linear, one-to-one and onto. In fact, thanks to Proposition 290, it is enough to show that ψ is invertible.

First proof.

1. ψ is one-to-one: see Theorem 2, page 105 in Lang (1971);
2. ψ is onto: see bottom of page 107 in Lang (1970).

Second proof.

1. $\psi \circ \varphi = id_{\mathcal{L}(V, U)}$.

Given $l \in \mathcal{L}(V, U)$, we want to show that $\forall v \in V$,

$$l(v) = ((\psi \circ \varphi)(l))(v)$$

i.e., from Proposition 268,

$$[l(v)]_{\mathbf{u}} = [((\psi \circ \varphi)(l))(v)]_{\mathbf{u}}.$$

First of all, observe that from (7.2), we have

$$[l(v)]_{\mathbf{u}} = [l]_{\mathbf{v}}^{\mathbf{u}} [v]_{\mathbf{v}}.$$

Moreover,

$$\begin{aligned} [((\psi \circ \varphi)(l))(v)]_{\mathbf{u}} &\stackrel{(1)}{=} [\psi([l]_{\mathbf{v}}^{\mathbf{u}}(v))]_{\mathbf{u}} \stackrel{(2)}{=} [\sum_{i=1}^n (i\text{-th row of } [l]_{\mathbf{v}}^{\mathbf{u}}) \cdot [v]_{\mathbf{v}} \cdot u^i]_{\mathbf{u}} \stackrel{(3)}{=} \\ &= [(i\text{-th row of } [l]_{\mathbf{v}}^{\mathbf{u}}) \cdot [v]_{\mathbf{v}}]_{i=1}^n \stackrel{(4)}{=} [l]_{\mathbf{v}}^{\mathbf{u}} \cdot [v]_{\mathbf{v}} \end{aligned}$$

where (1) comes from the definition of φ , (2) from the definition of ψ , (3) from the definition of $[\cdot]_{\mathbf{u}}$, (4) from the definition of product between matrices.

2. $\varphi \circ \psi = id_{\mathbb{M}(m, n)}$.

Given $A \in \mathbb{M}(m, n)$, we want to show that $(\varphi \circ \psi)(A) = A$. By definition of ψ ,

$$\psi(A) = l_{A, \mathbf{v}}^{\mathbf{u}} \text{ such that } \forall v \in V, l_{A, \mathbf{v}}^{\mathbf{u}}(v) = \sum_{i=1}^m R^i(A) \cdot [v]_{\mathbf{v}} \cdot u^i. \quad (7.3)$$

By definition of φ ,

$$\varphi(\psi(A)) = [l_{A, \mathbf{v}}^{\mathbf{u}}]_{\mathbf{v}}^{\mathbf{u}}.$$

Therefore, we want to show that $[l_{A, \mathbf{v}}^{\mathbf{u}}]_{\mathbf{v}}^{\mathbf{u}} = A$. Observe that from 7.3,

$$\begin{aligned} l_{A, \mathbf{v}}^{\mathbf{u}}(v^1) &= \sum_{i=1}^m R^i(A) \cdot [v^1]_{\mathbf{v}} \cdot u^i = \sum_{i=1}^m [a_{i1}, \dots, a_{ij}, \dots, a_{in}] \cdot [1, \dots, 0, \dots, 0] \cdot u^i = a_{11}u^1 + \dots + a_{i1}u^i + \dots + a_{m1}u^m \\ &\dots \\ l_{A, \mathbf{v}}^{\mathbf{u}}(v^j) &= \sum_{i=1}^m R^i(A) \cdot [v^j]_{\mathbf{v}} \cdot u^i = \sum_{i=1}^m [a_{i1}, \dots, a_{ij}, \dots, a_{in}] \cdot [0, \dots, 1, \dots, 0] \cdot u^i = a_{1j}u^1 + \dots + a_{ij}u^i + \dots + a_{mj}u^m \\ &\dots \\ l_{A, \mathbf{v}}^{\mathbf{u}}(v^n) &= \sum_{i=1}^m R^i(A) \cdot [v^n]_{\mathbf{v}} \cdot u^i = \sum_{i=1}^m [a_{i1}, \dots, a_{ij}, \dots, a_{in}] \cdot [0, \dots, 0, \dots, 1] \cdot u^i = a_{1n}u^1 + \dots + a_{in}u^i + \dots + a_{mn}u^m \end{aligned}$$

(From the above, it is clear why in definition 280 we take the transpose.) Therefore,

$$[l_{A, \mathbf{v}}^{\mathbf{u}}]_{\mathbf{v}}^{\mathbf{u}} = \begin{bmatrix} a_{11} & \dots & a_{1j} & \dots & a_{1n} \\ \dots & & & & \\ a_{i1} & & a_{ij} & & a_{in} \\ \dots & & & & \\ a_{m1} & \dots & a_{mj} & \dots & a_{mn} \end{bmatrix} = A,$$

as desired.

The fact that $\dim \mathcal{L}(V, U)$ follows from Proposition 274. ■

Proposition 292 *Let the following objects be given.*

1. Vector spaces V with ordered basis $\mathbf{v} = \{v^1, \dots, v^j, \dots, v^n\}$, U with ordered basis $\mathbf{u} = \{u^1, \dots, u^i, \dots, u^m\}$ and W with ordered basis $\mathbf{w} = \{w^1, \dots, w^k, \dots, w^p\}$;
2. $l_1 \in \mathcal{L}(V, U)$ and $l_2 \in \mathcal{L}(U, W)$.

Then

$$[l_2 \circ l_1]_{\mathbf{v}}^{\mathbf{w}} = [l_2]_{\mathbf{u}}^{\mathbf{w}} \cdot [l_1]_{\mathbf{v}}^{\mathbf{u}},$$

or

$$\varphi_{\mathbf{v}}^{\mathbf{w}}(l_2 \circ l_1) = \varphi_{\mathbf{u}}^{\mathbf{w}}(l_2) \cdot \varphi_{\mathbf{v}}^{\mathbf{u}}(l_1).$$

Proof. By definition²

$$\begin{aligned} [l_1]_{\mathbf{v}}^{\mathbf{u}} &= \begin{bmatrix} [l_1(v^1)]_{\mathbf{u}} & \dots & [l_1(v^j)]_{\mathbf{u}} & \dots & [l_1(v^n)]_{\mathbf{u}} \end{bmatrix}_{m \times n} := \\ &:= \begin{bmatrix} l_1^1(v^1) & \dots & l_1^1(v^j) & \dots & l_1^1(v^n) \\ \vdots & & \vdots & & \vdots \\ l_1^i(v^1) & & l_1^i(v^j) & & l_1^i(v^n) \\ \vdots & & \vdots & & \vdots \\ l_1^m(v^1) & \dots & l_1^m(v^j) & \dots & l_1^m(v^n) \end{bmatrix} := \begin{bmatrix} l_1^{11} & \dots & l_1^{1j} & \dots & l_1^{1n} \\ \vdots & & \vdots & & \vdots \\ l_1^{i1} & & l_1^{ij} & & l_1^{in} \\ \vdots & & \vdots & & \vdots \\ l_1^{m1} & \dots & l_1^{mj} & \dots & l_1^{mn} \end{bmatrix} := \\ &:= [l_1^{ij}]_{i \in \{1, \dots, m\}, j \in \{1, \dots, n\}} := A \in \mathbb{M}(m, n), \end{aligned}$$

and therefore $\forall j \in \{1, \dots, n\}$, $l_1(v^j) = \sum_{i=1}^m l_1^{ij} \cdot u^i$.

Similarly,

$$\begin{aligned} [l_2]_{\mathbf{u}}^{\mathbf{w}} &= \begin{bmatrix} [l_2(u^1)]_{\mathbf{w}} & \dots & [l_2(u^i)]_{\mathbf{w}} & \dots & [l_2(u^m)]_{\mathbf{w}} \end{bmatrix}_{p \times m} := \\ &:= \begin{bmatrix} l_2^1(u^1) & \dots & l_2^1(u^i) & \dots & l_2^1(u^m) \\ \vdots & & \vdots & & \vdots \\ l_2^k(u^1) & & l_2^k(u^i) & & l_2^k(u^m) \\ \vdots & & \vdots & & \vdots \\ l_2^p(u^1) & \dots & l_2^p(u^i) & \dots & l_2^p(u^m) \end{bmatrix} := \begin{bmatrix} l_2^{11} & \dots & l_2^{1i} & \dots & l_2^{1m} \\ \vdots & & \vdots & & \vdots \\ l_2^{k1} & & l_2^{ki} & & l_2^{km} \\ \vdots & & \vdots & & \vdots \\ l_2^{p1} & \dots & l_2^{pi} & \dots & l_2^{pm} \end{bmatrix} := \\ &:= [l_2^{ki}]_{k \in \{1, \dots, p\}, i \in \{1, \dots, m\}} := B \in \mathbb{M}(p, m), \end{aligned}$$

and therefore $\forall i \in \{1, \dots, m\}$, $l_2(u^i) = \sum_{k=1}^p l_2^{ki} \cdot w^k$.

Moreover, defined $l := (l_2 \circ l_1)$, we get

$$\begin{aligned} [l_2 \circ l_1]_{\mathbf{v}}^{\mathbf{w}} &= \begin{bmatrix} [l(v^1)]_{\mathbf{w}} & \dots & [l(v^j)]_{\mathbf{w}} & \dots & [l(v^n)]_{\mathbf{w}} \end{bmatrix}_{p \times n} := \\ &:= \begin{bmatrix} l^1(v^1) & \dots & l^1(v^j) & \dots & l^1(v^n) \\ \vdots & & \vdots & & \vdots \\ l^k(v^1) & & l^k(v^j) & & l^k(v^n) \\ \vdots & & \vdots & & \vdots \\ l^p(v^1) & \dots & l^p(v^j) & \dots & l^p(v^n) \end{bmatrix} := \begin{bmatrix} l^{11} & \dots & l^{1j} & \dots & l^{1n} \\ \vdots & & \vdots & & \vdots \\ l^{k1} & & l^{kj} & & l^{kn} \\ \vdots & & \vdots & & \vdots \\ l^{p1} & \dots & l^{pj} & \dots & l^{pn} \end{bmatrix} := \\ &:= [l^{kj}]_{k \in \{1, \dots, p\}, j \in \{1, \dots, n\}} := C \in \mathbb{M}(p, n), \end{aligned}$$

and therefore $\forall j \in \{1, \dots, n\}$, $l(v^j) = \sum_{k=1}^p l^{kj} \cdot w^k$.

Now, $\forall j \in \{1, \dots, n\}$

$$\begin{aligned} l(v^j) &= (l_2 \circ l_1)(v^j) = l_2(l_1(v^j)) = l_2\left(\sum_{i=1}^m l_1^{ij} \cdot u^i\right) = \\ &= \sum_{i=1}^m l_1^{ij} \cdot l_2(u^i) = \sum_{i=1}^m l_1^{ij} \cdot \sum_{k=1}^p l_2^{ki} \cdot w^k = \sum_{k=1}^p \sum_{i=1}^m l_2^{ki} \cdot l_1^{ij} \cdot w^k. \end{aligned}$$

² $[l_1(v^1)]_{\mathbf{u}}, \dots, [l_1(v^n)]_{\mathbf{u}}$ are column vectors.

The above says that $\forall j \in \{1, \dots, n\}$, the j -th column of C is

$$\begin{bmatrix} \sum_{i=1}^m l_2^{1i} \cdot l_1^{ij} \\ \sum_{i=1}^m l_2^{ki} \cdot l_1^{ij} \\ \sum_{i=1}^m l_2^{pi} \cdot l_1^{ij} \end{bmatrix}$$

On the other hand, the j -th column of $B \cdot A$ is

$$\begin{bmatrix} [1st \text{ row of } B] \cdot [j-th \text{ column of } A] \\ [k-th \text{ row of } B] \cdot [j-th \text{ column of } A] \\ [p-th \text{ row of } B] \cdot [j-th \text{ column of } A] \end{bmatrix} = \begin{bmatrix} \begin{bmatrix} l_2^{11} & \dots & l_2^{1i} & \dots & l_2^{1m} \end{bmatrix} \cdot \begin{bmatrix} l_1^{1j} \\ l_1^{ij} \\ l_1^{mj} \end{bmatrix} \\ \dots \\ \begin{bmatrix} l_2^{k1} & l_2^{ki} & l_2^{km} \end{bmatrix} \cdot \begin{bmatrix} l_1^{1j} \\ l_1^{ij} \\ l_1^{mj} \end{bmatrix} \\ \dots \\ \begin{bmatrix} l_2^{p1} & \dots & l_2^{pi} & \dots & l_2^{pm} \end{bmatrix} \cdot \begin{bmatrix} l_1^{1j} \\ l_1^{ij} \\ l_1^{mj} \end{bmatrix} \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^m l_2^{1i} \cdot l_1^{ij} \\ \sum_{i=1}^m l_2^{ki} \cdot l_1^{ij} \\ \sum_{i=1}^m l_2^{pi} \cdot l_1^{ij} \end{bmatrix}$$

as desired. ■

7.4 Some related properties of a linear function and associated matrix

In this section, the following objects will be given: a vector space V with a ordered basis $\mathbf{v} = \{v^1, \dots, v^n\}$; a vector space U with a ordered basis $\mathbf{u} = \{u^1, \dots, u^n\}$; $l \in \mathcal{L}(V, U)$ and $A \in \mathbb{M}(m, n)$.

From Remark 152, recall that

$$\text{colspan } A = \{z \in \mathbb{R}^m : \exists x \in \mathbb{R}^n \text{ such that } z = Ax\};$$

Lemma 293 $cr_{\mathbf{u}}(\text{Im } l) = \text{colspan } [l]_{\mathbf{v}}^{\mathbf{u}}$.

Proof. $[\subseteq]$

$y \in cr_{\mathbf{u}}(\text{Im } l) \xrightarrow{\text{def } cr} \exists u \in \text{Im } l \text{ such that } cr_{\mathbf{u}}(u) = [u]_{\mathbf{u}} = y \xrightarrow{\text{def } \text{Im } l} \exists v \in V \text{ such that } l(v) = u \Rightarrow \exists v \in V \text{ such that } [l(v)]_{\mathbf{u}} = y \xrightarrow{\text{Prop. 284}} \exists v \in V \text{ such that } [l]_{\mathbf{v}}^{\mathbf{u}} \cdot [v]_{\mathbf{v}} = y \Rightarrow y \in \text{colspan } [l]_{\mathbf{v}}^{\mathbf{u}}.$

$[\supseteq]$

We want to show that $y \in \text{colspan } [l]_{\mathbf{v}}^{\mathbf{u}} \Rightarrow y \in cr_{\mathbf{u}}(\text{Im } l)$, i.e., $\exists u \in \text{Im } l$ such that $y = [u]_{\mathbf{u}}$.

$y \in \text{colspan } [l]_{\mathbf{v}}^{\mathbf{u}} \Rightarrow \exists x_y \in \mathbb{R}^n \text{ such that } [l]_{\mathbf{v}}^{\mathbf{u}} \cdot x_y = y \xrightarrow{\text{def } cr} \exists v = \sum_{j=1}^n x_{y,j} v^j \text{ such that } [l]_{\mathbf{v}}^{\mathbf{u}} \cdot [v]_{\mathbf{v}} = y \xrightarrow{\text{Prop. 284}} \exists v \in V \text{ such that } [l(v)]_{\mathbf{u}} = y \xrightarrow{u=l(v)} \exists u \in \text{Im } l \text{ such that } y = [u]_{\mathbf{u}}, \text{ as desired. } \blacksquare$

Lemma 294 $\dim \text{Im } l = \dim \text{colspan } [l]_{\mathbf{v}}^{\mathbf{u}} = \text{rank } [l]_{\mathbf{v}}^{\mathbf{u}}$.

Proof. It follows from the above Lemma, the proof of Proposition 268, which says that $cr_{\mathbf{u}}$ is an isomorphism and Proposition 274, which says that isomorphic spaces have the same dimension. ■

Proposition 295 Given $l \in \mathcal{L}(V, U)$,

1. l onto $\Leftrightarrow \text{rank } [l]_{\mathbf{v}}^{\mathbf{u}} = \dim U$;
2. l one-to-one $\Leftrightarrow \text{rank } [l]_{\mathbf{v}}^{\mathbf{u}} = \dim V$;
3. l invertible $\Leftrightarrow [l]_{\mathbf{v}}^{\mathbf{u}}$ invertible, and in that case $[l^{-1}]_{\mathbf{u}}^{\mathbf{v}} = [l]_{\mathbf{v}}^{\mathbf{u}}^{-1}$,
i.e.,

Proof. Recall that from Remark 261 and Proposition 272, l one-to-one $\Leftrightarrow l$ nonsingular $\Leftrightarrow \ker l = \{0\}$.

1. l onto $\Leftrightarrow \text{Im } l = U \Leftrightarrow \dim \text{Im } l = \dim U \stackrel{\text{Lemma 294}}{\Leftrightarrow} \text{rank } [l]_{\mathbf{v}}^{\mathbf{u}} = \dim U$.
2. l one-to-one $\stackrel{\text{Proposition 257}}{\Leftrightarrow} \dim \text{Im } l = \dim V \stackrel{\text{Lemma 294}}{\Leftrightarrow} \text{rank } [l]_{\mathbf{v}}^{\mathbf{u}} = \dim V$.
3. The first part of the statement follows from 1. and 2. above. The second part is proven below. First of all observe that for any vector space W with ordered basis \mathbf{w} , $\text{id}_W \in \mathcal{L}(W, W)$ and if W has a ordered basis $\mathbf{w} = \{w^1, \dots, w^k\}$, we also have that

$$[\text{id}_W]_{\mathbf{w}}^{\mathbf{w}} = [[\text{id}_W(w^1)]_{\mathbf{w}}, \dots, [\text{id}_W(w^k)]_{\mathbf{w}}] = I_k.$$

Moreover, if l is invertible

$$l^{-1} \circ l = \text{id}_V$$

and

$$[l^{-1} \circ l]_{\mathbf{v}}^{\mathbf{v}} = [\text{id}_V]_{\mathbf{v}}^{\mathbf{v}} = I_m.$$

Since

$$[l^{-1} \circ l]_{\mathbf{v}}^{\mathbf{v}} = [l^{-1}]_{\mathbf{u}}^{\mathbf{v}} \cdot [l]_{\mathbf{v}}^{\mathbf{u}},$$

the desired result follows.

■

Remark 296 From the definitions of φ and ψ , we have what follows:

1.

$$l = \psi(\varphi(l)) = \psi([l]_{\mathbf{v}}^{\mathbf{u}}) = l_{[l]_{\mathbf{v}}^{\mathbf{u}}, \mathbf{v}}^{\mathbf{u}}.$$

2.

$$A = \varphi(\psi(A)) = \varphi(l_{A, \mathbf{v}}^{\mathbf{u}}) = [l_{A, \mathbf{v}}^{\mathbf{u}}]_{\mathbf{v}}^{\mathbf{u}}.$$

Lemma 297 $\text{cr}_{\mathbf{u}}(\text{Im } l_{A, \mathbf{v}}^{\mathbf{u}}) = \text{colspan } A$.

Proof. Recall that $\varphi(l) = [l]_{\mathbf{v}}^{\mathbf{u}}$ and $\psi(A) = l_{A, \mathbf{v}}^{\mathbf{u}}$. For any $l \in \mathcal{L}(V, U)$,

$$\text{cr}_{\mathbf{u}}(\text{Im } l) \stackrel{\text{Lemma 293}}{=} \text{colspan } [l]_{\mathbf{v}}^{\mathbf{u}} \stackrel{\text{Def. 282}}{=} \text{colspan } \varphi(l) \quad (7.4)$$

Take $l = l_{A, \mathbf{v}}^{\mathbf{u}}$. Then from (7.4), we have

$$\text{cr}_{\mathbf{u}}(\text{Im } l_{A, \mathbf{v}}^{\mathbf{u}}) = \text{colspan } \varphi(l_{A, \mathbf{v}}^{\mathbf{u}}) \stackrel{\text{Rmk. 296.2}}{=} \text{colspan } A.$$

■

Lemma 298 $\dim \text{Im } l_{A, \mathbf{v}}^{\mathbf{u}} = \dim \text{colspan } A = \text{rank } A$.

Proof. Since Lemma 294 holds for any $l \in \mathcal{L}(V, U)$ and $l_{A, \mathbf{v}}^{\mathbf{u}} \in \mathcal{L}(V, U)$, we have that

$$\dim \text{Im } l_{A, \mathbf{v}}^{\mathbf{u}} = \dim \text{colspan } [l_{A, \mathbf{v}}^{\mathbf{u}}]_{\mathbf{v}}^{\mathbf{u}} \stackrel{\text{Rmk. 296.2}}{=} \dim \text{colspan } A \stackrel{\text{Rmk. 244}}{=} \text{rank } A.$$

■

Proposition 299 Let $A \in \mathbb{M}(m, n)$ be given.

1. $\text{rank} A = m \Leftrightarrow l_{A,\mathbf{v}}^{\mathbf{u}}$ onto;
2. $\text{rank} A = n \Leftrightarrow l_{A,\mathbf{v}}^{\mathbf{u}}$ one-to-one;
3. A invertible $\Leftrightarrow l_{A,\mathbf{v}}^{\mathbf{u}}$ invertible, and in that case $l_{A^{-1},\mathbf{u}}^{\mathbf{v}} = (l_{A,\mathbf{v}}^{\mathbf{u}})^{-1}$.

Proof. 1. $\text{rank} A = m \stackrel{\text{Lemma 298}}{\Leftrightarrow} \dim \text{Im } l_{A,\mathbf{v}}^{\mathbf{u}} = m \Leftrightarrow l_{A,\mathbf{v}}^{\mathbf{u}}$ onto;

2. $\text{rank} A = n \stackrel{(1)}{\Leftrightarrow} \dim \ker l_{A,\mathbf{v}}^{\mathbf{u}} = 0 \stackrel{\text{Proposition 272}}{\Leftrightarrow} l_{A,\mathbf{v}}^{\mathbf{u}}$ one-to-one,

where (1) the first equivalence follows from the fact that $n = \dim \ker l_{A,\mathbf{v}}^{\mathbf{u}} + \dim \text{Im } l_{A,\mathbf{v}}^{\mathbf{u}}$, and Lemma 298.

3. First statement: A invertible $\stackrel{\text{Prop. 239}}{\Leftrightarrow} \text{rank} A = m = n \stackrel{1 \text{ and } 2 \text{ above}}{\Leftrightarrow} l_{A,\mathbf{v}}^{\mathbf{u}}$ invertible.

Second statement: Since $l_{A,\mathbf{v}}^{\mathbf{u}}$ invertible, there exists $(l_{A,\mathbf{v}}^{\mathbf{u}})^{-1} : U \rightarrow V$ such that

$$id_V = (l_{A,\mathbf{v}}^{\mathbf{u}})^{-1} \circ l_{A,\mathbf{v}}^{\mathbf{u}}.$$

Then

$$I = \varphi_{\mathbf{v}}^{\mathbf{v}}(id_V) \stackrel{\text{Prop. 292}}{=} \varphi_{\mathbf{u}}^{\mathbf{v}}\left((l_{A,\mathbf{v}}^{\mathbf{u}})^{-1}\right) \cdot \varphi_{\mathbf{v}}^{\mathbf{u}}(l_{A,\mathbf{v}}^{\mathbf{u}}) \stackrel{\text{Rmk. 296}}{=} \varphi_{\mathbf{u}}^{\mathbf{v}}\left((l_{A,\mathbf{v}}^{\mathbf{u}})^{-1}\right) \cdot A.$$

Then, by definition of inverse matrix,

$$A^{-1} = \varphi_{\mathbf{u}}^{\mathbf{v}}\left((l_{A,\mathbf{v}}^{\mathbf{u}})^{-1}\right)$$

and

$$\psi_{\mathbf{u}}^{\mathbf{v}}(A^{-1}) = (\psi_{\mathbf{u}}^{\mathbf{v}} \circ \varphi_{\mathbf{u}}^{\mathbf{v}})\left((l_{A,\mathbf{v}}^{\mathbf{u}})^{-1}\right) = id_{\mathcal{L}(U,U)}\left((l_{A,\mathbf{v}}^{\mathbf{u}})^{-1}\right) = (l_{A,\mathbf{v}}^{\mathbf{u}})^{-1}.$$

Finally, from the definition of $\psi_{\mathbf{u}}^{\mathbf{v}}$, we have

$$\psi_{\mathbf{u}}^{\mathbf{v}}(A^{-1}) = l_{A^{-1},\mathbf{u}}^{\mathbf{v}},$$

as desired. ■

Remark 300 Consider $A \in \mathbb{M}(n, n)$. Then from Proposition 299,

$$A \text{ invertible} \Leftrightarrow l_{A,\mathbf{v}}^{\mathbf{u}} \text{ invertible};$$

from Proposition 239,

$$A \text{ invertible} \Leftrightarrow A \text{ nonsingular};$$

from Proposition 275,

$$l_{A,\mathbf{v}}^{\mathbf{u}} \text{ invertible} \Leftrightarrow l_{A,\mathbf{v}}^{\mathbf{u}} \text{ nonsingular}.$$

Therefore,

$$A \text{ nonsingular} \Leftrightarrow l_{A,\mathbf{v}}^{\mathbf{u}} \text{ nonsingular}.$$

Symmetrically,

$$[l]_{\mathbf{v}}^{\mathbf{u}} \text{ invertible} \Leftrightarrow [l]_{\mathbf{v}}^{\mathbf{u}} \text{ nonsingular} \Leftrightarrow l \text{ invertible} \Leftrightarrow l \text{ nonsingular}.$$

Proposition 301 Let $l \in \mathcal{L}(V, U)$ be given. Then there exists a ordered basis \mathcal{V} of V and a ordered basis \mathcal{U} of U such that

$$[l]_{\mathcal{V}}^{\mathcal{U}} = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix},$$

where I is the r -square identity matrix and $r = \text{rank}[l]_{\mathcal{V}}^{\mathcal{U}}$.

Proof. Suppose $\dim V = m$ and $\dim U = n$. Let $W = \ker l$ and $U' = \text{Im } l$. By assumption, $\text{rank}[l]_{\mathcal{V}}^{\mathcal{U}} = r$. Then, by Lemma 294,

$$\dim \text{Im } l = \text{rank}[l]_{\mathcal{V}}^{\mathcal{U}} = r.$$

Therefore, from the Dimension Theorem

$$\dim \ker = \dim V - \dim \text{Im } l = m - r.$$

Let $\mathcal{W} = \{w^1, \dots, w^{m-r}\}$ be a ordered basis of W . Then, from the Completion Lemma, there exists $\{v^1, \dots, v^r\} \in V$ such that $\mathcal{V} := \{v^1, \dots, v^r, w^1, \dots, w^{m-r}\}$ is a ordered basis of V . For any $i \in \{1, \dots, r\}$, set $u^i = l(v^i)$. Then, $\mathcal{U}'^1, \dots, u^r\}$ is a ordered basis of U' . Then, again from the Completion Lemma, there exists $\{u^{r+1}, \dots, u^n\} \in U$ such that $\mathcal{U} := \{u^1, \dots, u^r, u^{r+1}, \dots, u^n\}$ is a ordered basis of U . Then,

$$\begin{aligned} l(v^1) &= u^1 = 1u^1 + 0u^2 + \dots + 0u^r + 0u^{r+1} + \dots + 0u^n, \\ l(v^2) &= u^2 = 0u^1 + 1u^2 + \dots + 0u^r + 0u^{r+1} + \dots + 0u^n, \\ &\dots\dots\dots \\ l(v^r) &= u^r = 0u^1 + 0u^2 + \dots + 1u^r + 0u^{r+1} + \dots + 0u^n, \\ l(w^1) &= 0 = 0u^1 + 0u^2 + \dots + 0u^r + 0u^{r+1} + \dots + 0u^n, \\ &\dots\dots\dots \\ l(w^{m-r}) &= 0 = 0u^1 + 0u^2 + \dots + 0u^r + 0u^{r+1} + \dots + 0u^n, \end{aligned}$$

i.e.,

$$[l]_{\mathcal{V}}^{\mathcal{U}} = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}.$$

■

7.5 Some facts on $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$

In this Section, we specialize (basically repeat) the content of the previous Section in the important case in which

$$\begin{aligned} V &= \mathbb{R}^n, & \mathbf{v} &= (e_n^j)_{j=1}^n := \mathbf{e}_n \\ U &= \mathbb{R}^m & \mathbf{u} &= (e_m^i)_{i=1}^m := \mathbf{e}_m \\ v &= x \end{aligned} \tag{7.5}$$

and therefore

$$l \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m).$$

From $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ to $\mathbb{M}(m, n)$.

From Definition 280, we have

$$\begin{aligned} [l]_{\mathbf{e}_n}^{\mathbf{e}_m} &= \begin{bmatrix} [l(e_n^1)]_{\mathbf{e}_m} & \dots & [l(e_n^j)]_{\mathbf{e}_m} & \dots & [l(e_n^n)]_{\mathbf{e}_m} \end{bmatrix} = \\ &= \begin{bmatrix} l(e_n^1) & \dots & l(e_n^j) & \dots & l(e_n^n) \end{bmatrix} := [l]; \end{aligned} \tag{7.6}$$

from Definition 282,

$$\varphi := \varphi_{\mathbf{e}_n}^{\mathbf{e}_m} : \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m) \rightarrow \mathbb{M}(m, n), \quad l \mapsto [l];$$

from Proposition 284,

$$[l] \cdot x = l(x). \tag{7.7}$$

From $\mathbb{M}(m, n)$ to $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$.

From Definition 286,

$$l_A := l_{A, \mathbf{e}_m}^{\mathbf{e}_m} : \mathbb{R}^n \rightarrow \mathbb{R}^m, \tag{7.8}$$

$$\begin{aligned} l_A(x) &= \sum_{i=1}^m (R^i(A) \cdot [x]_{\mathbf{e}_n}) \cdot e_m^i = \\ &= \begin{bmatrix} R^1(A) \cdot x \\ \dots \\ R^i(A) \cdot x \\ \dots \\ R^m(A) \cdot x \end{bmatrix} = Ax. \end{aligned} \tag{7.9}$$

From Definition 289,

$$\psi := \psi_{\mathbf{e}_n}^{\mathbf{e}_m} : \mathbb{M}(m, n) \rightarrow \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m) \quad : A \mapsto l_A. \quad .$$

From Proposition 291,

$\mathbb{M}(m, n)$ and $\mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ are isomorphic.

From Proposition 292, if $l_1 \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ and $l_2 \in \mathcal{L}(\mathbb{R}^m, \mathbb{R}^p)$, then

$$[l_2 \circ l_1] = [l_2] \cdot [l_1]. \quad (7.10)$$

Some related properties.

From Proposition 295, given $l \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$,

1. l onto $\Leftrightarrow \text{rank } [l] = m$;
2. l one-to-one $\Leftrightarrow \text{rank } [l] = n$;
3. l invertible $\Leftrightarrow [l]$ invertible, and in that case $[l^{-1}] = [l]^{-1}$.

From Remark 296,

1.

$$l = \psi(\varphi(l)) = \psi([l]) = l_{[l]}.$$

2.

$$A = \varphi(\psi(A)) = \varphi(l_A) = [l_A].$$

From Proposition 299, given $A \in \mathbb{M}(m, n)$,

1. $\text{rank } A = m \Leftrightarrow l_A$ onto;
2. $\text{rank } A = n \Leftrightarrow l_A$ one-to-one;
3. A invertible $\Leftrightarrow l_A$ invertible, and in that case $l_{A^{-1}} = (l_A)^{-1}$.

Remark 302 From (7.7) and Remark 152,

$$\text{Im } l := \{y \in \mathbb{R}^m : \exists x \in \mathbb{R}^n \text{ such that } y = [l] \cdot x\} = \text{colspan } [l]. \quad (7.11)$$

Then, from the above and Remark 244,

$$\dim \text{Im } l = \text{rank } [l] = \max \# \text{ linearly independent columns of } [l]. \quad (7.12)$$

Similarly, from (7.9) and Remark 244, we get

$$\text{Im } l_A := \{y \in \mathbb{R}^m : \exists x \in \mathbb{R}^n \text{ such that } y = l_A \cdot x = Ax\} = \text{colspan } A,$$

and

$$\dim \text{Im } l_A = \text{rank } A = \max \# \text{ linearly independent columns of } A.$$

Remark 303 Assume that $l \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$. The above Remark gives a way of finding a ordered basis of $\text{Im } l$: it is enough to consider a number equal to $\text{rank } [l]$ of linearly independent vectors among the column vectors of $[l]$. In a more detailed way, we have what follows.

1. Compute $[l]$.
2. Compute $\dim \text{Im } l = \text{rank } [l] := r$.
3. To find a ordered basis of $\text{Im } l$, we have to find r vectors which are a. linearly independent, and b. elements of $\text{Im } l$. Indeed, it is enough to take r linearly independent columns of $[l]$. Observe that for any $i \in \{1, \dots, m\}$, $C^i([l]) \in \text{Im } l = \text{colspan } [l]$.

To get a “simpler” basis, you can make elementary operations on those column. Recall that elementary operations on linearly independent vectors lead to linearly independent vectors, and that elementary operations on vectors lead to vector belonging to the span of the set of starting vectors.

Example 304 Given $n \in \mathbb{N}$ with $n \geq 3$, and the linear function

$$l : \mathbb{R}^n \rightarrow \mathbb{R}^3, x := (x_i)_{i=1}^n \mapsto \begin{cases} \sum_{i=1}^n x_i \\ x_2 \\ x_2 + x_3 \end{cases}.$$

find a basis for $\text{Im } l$.

1.

$$[l] = \begin{bmatrix} 1 & 1 & 1 & 1 & \dots & 1 \\ 0 & 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 1 & 0 & \dots & 0 \end{bmatrix}$$

2.

$$\text{rank} \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix} = \text{rank} \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = 3$$

3. A basis of $\text{Im } l_a$ is given by the column vectors of

$$\begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Remark 305 From (7.7), we have that

$$\ker l = \{x \in \mathbb{R}^n : [l]x = 0\},$$

i.e., $\ker l$ is the set, in fact the vector space, of solution to the systems $[l]x = 0$. In Remark 337, we will describe an algorithm to find a basis of the kernel of an arbitrary linear function.

Remark 306 From Remark 302 and Proposition 257 (the Dimension Theorem), given $l \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$,

$$\dim \mathbb{R}^n = \dim \ker l + \text{rank } [l],$$

and given $A \in \mathbb{M}(m, n)$,

$$\dim \mathbb{R}^n = \dim \ker l_A + \text{rank } A.$$

7.6 Examples of computation of $[l]_{\mathbf{V}}^{\mathbf{U}}$

1. $id \in \mathcal{L}(V, V)$.

$$\begin{aligned} [id]_{\mathbf{V}}^{\mathbf{V}} &= [[id(v^1)]_{\mathbf{V}}, \dots, [id(v^j)]_{\mathbf{V}}, \dots, [id(v^n)]_{\mathbf{V}}] = \\ &= [[v^1]_{\mathbf{V}}, \dots, [v^j]_{\mathbf{V}}, \dots, [v^n]_{\mathbf{V}}] = [e_n^1, \dots, e_n^j, \dots, e_n^n] = I_n. \end{aligned}$$

2. $0 \in \mathcal{L}(V, U)$.

$$[0]_{\mathbf{V}}^{\mathbf{U}} = [[0]_{\mathbf{V}}, \dots, [0]_{\mathbf{V}}, \dots, [0]_{\mathbf{V}}] = 0 \in \mathbb{M}(m, n).$$

3. $l_{\alpha} \in \mathcal{L}(V, V)$, with $\alpha \in F$.

$$\begin{aligned} [l_{\alpha}]_{\mathbf{V}}^{\mathbf{V}} &= [[\alpha \cdot v^1]_{\mathbf{V}}, \dots, [\alpha \cdot v^j]_{\mathbf{V}}, \dots, [\alpha \cdot v^n]_{\mathbf{V}}] = [\alpha \cdot [v^1]_{\mathbf{V}}, \dots, \alpha \cdot [v^j]_{\mathbf{V}}, \dots, \alpha \cdot [v^n]_{\mathbf{V}}] = \\ &= [\alpha \cdot e_n^1, \dots, \alpha \cdot e_n^j, \dots, \alpha \cdot e_n^n] = \alpha \cdot I_n. \end{aligned}$$

4. $l_A \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$, with $A \in \mathbb{M}(m, n)$.

$$[l_A] = [A \cdot e_n^1, \dots, A \cdot e_n^j, \dots, A \cdot e_n^n] = A \cdot [e_n^1, \dots, e_n^j, \dots, e_n^n] = A \cdot I_n = A.$$

5. (projection function) $proj_{n+k, n} \in \mathcal{L}(\mathbb{R}^{n+k}, \mathbb{R}^n)$, $proj_{n+k, n} : (x_i)_{i=1}^{n+k} \mapsto (x_i)_{i=1}^n$. Defined $proj_{n+k, n} := p$, we have

$$[p] = [p(e_{n+k}^1), \dots, p(e_{n+k}^n), p(e_{n+k}^{n+1}), \dots, p(e_{n+k}^{n+k})] = [e_n^1, \dots, e_n^n, 0, \dots, 0] = [I_n | 0],$$

where $0 \in \mathbb{M}(n, k)$.

6. (immersion function) $i_{n, n+k} \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^{n+k})$, $i_{n, n+k} : (x_i)_{i=1}^n \mapsto ((x_i)_{i=1}^n, 0)$ with $0 \in \mathbb{R}^k$. Defined $i_{n, n+k} := i$, we have

$$[i] = [i(e_n^1), \dots, i(e_n^n)] = \begin{bmatrix} e_n^1 & \dots & e_n^n \\ 0 & \dots & 0 \end{bmatrix} = \begin{bmatrix} I_n \\ 0 \end{bmatrix},$$

where $0 \in \mathbb{M}(k, n)$.

Remark 307 Point 4. above implies that if $l : \mathbb{R}^n \rightarrow \mathbb{R}^m$, $x \mapsto Ax$, then $[l] = A$. In other words, to compute $[l]$ you do not have to take the image of each element in the canonical basis; the first line of $[l]$ is the vector of the coefficient of the first component function of l and so on. For example, if $l : \mathbb{R}^2 \rightarrow \mathbb{R}^3$,

$$x \mapsto \begin{cases} a_{11}x_1 + a_{12}x_2 \\ a_{21}x_1 + a_{22}x_2 \\ a_{31}x_1 + a_{32}x_2 \end{cases},$$

then

$$[l] = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}$$

7.7 Exercises

Problem sets: 15,16,17,18,19,21,22.

From Lipschutz (1991), starting from page 352:

10.1 \rightarrow 10.9, 10.29 \rightarrow 10.33.

7.8 Appendices.

7.8.1 The dual and double dual space of a vector space

The dual space of a vector space

Definition 308 The³ dual space of a vector space V is

$$V^* := \mathcal{L}(V, \mathbb{R}).$$

Remark 309 From Proposition 291, we have that if $\dim V$ is finite, then $\dim V^* = \dim V$.

Definition 310 The Kronecker delta is a function defined as follows

$$\delta : \mathbb{N}^2 \rightarrow \{0, 1\}, \quad (i, j) \mapsto \delta_{ij} := \begin{cases} 1 & \text{if } i = j \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 311 Let V be a finite dimensional vector space over a field F and let $\mathcal{V} = \{v^i\}_{i=1}^n$ be a basis for V . Then,

1. there exists a unique basis $\mathcal{V}^* = \{l_i\}_{i=1}^n$ for V^* such that $\forall i, j \in \{1, \dots, n\}$, $l_i(v^j) = \delta_{ij}$, i.e., $(l_i(v^j))_{i=1}^n = e_n^j$;
- 2.

$$\forall f \in V^*, \quad f = \sum_{i=1}^n f(v^i) \cdot l_i, \text{ i.e., } [f]_{\mathcal{V}^*} = (f(v^i))_{i=1}^n,$$

i.e., the coordinates of f with respect to \mathcal{V}^* are the images via f of the elements of the basis \mathcal{V} , and

3.

$$\forall v \in V, \quad v = \sum_{i=1}^n l_i(v) \cdot v^i, \text{ i.e., } [v]_{\mathcal{V}} = (l_i(v))_{i=1}^n,$$

i.e., the coordinates of v with respect to \mathcal{V} are the images of v via the elements of the basis \mathcal{V} .

Proof. 1. From Proposition 269, we have that given a basis $\mathcal{V} = \{v^i\}_{i=1}^n$ for V ,

$$\forall i \in \{1, \dots, n\}, \quad \exists ! l_i \in V^* \text{ such that } \forall j \in \{1, \dots, n\}, \quad l_i(v^j) = \delta_{ij}.$$

We are left with showing that $\{l_i\}_{i=1}^n$ is a linearly independent and the therefore a basis of the n dimensional vector space V^* . We want to show that if $\sum_{i=1}^n \alpha_i l_i = 0$, then $(\alpha_i)_{i=1}^n = 0$, i.e., if $\forall v \in V$, $\sum_{i=1}^n \alpha_i l_i(v) = 0$, then $(\alpha_i)_{i=1}^n = 0$. Then, $\forall j \in \{1, \dots, n\}$, $0 = \sum_{i=1}^n \alpha_i l_i(v^j) = \alpha_j$, as desired.

³Here I follow pages 98 and 99 in Hoffman and Kunze (1971).

2. Take $f \in V^*$; then, since $\mathcal{V}^* = \{l_i\}_{i=1}^n$ is a basis for V^* , $\exists (\beta_i)_{i=1}^n \in \mathbb{R}^n$ such that $f = \sum_{i=1}^n \beta_i l_i$. Then,

$$\forall v^j \in \mathcal{V}, \quad f(v^j) = \sum_{i=1}^n \beta_i l_i(v^j) = \beta_j,$$

as desired.

3. Take $v \in V$; then, since $\mathcal{V} = \{v^i\}_{i=1}^n$ is a basis for V , $\exists (\gamma_i)_{i=1}^n \in \mathbb{R}^n$ such that $v = \sum_{i=1}^n \gamma_i v^i$. Then,

$$\forall l^j \in \mathcal{V}^*, \quad l_j(v) = \sum_{i=1}^n \gamma_i l_j(v^i) = \gamma_j,$$

as desired. ■

Definition 312 Let a basis $\mathcal{V} = \{v^i\}_{i=1}^n$ for V be given. The unique basis $\mathcal{V}^* = \{l_i\}_{i=1}^n$ for V^* such that $\forall i, j \in \{1, \dots, n\}$, $l_i(v^j) = \delta_{ij}$ is called the dual basis of \mathcal{V} .

Corollary 313 If \mathcal{V}^* is the unique dual basis of $\mathcal{V} = \{v^i\}_{i=1}^n$, defined $l : V \rightarrow \mathbb{R}^n$, $v \mapsto (l_i(v))_{i=1}^n$, we have that

$$[l]_{\mathcal{V}}^{\mathcal{E}_n} = I \in \mathbb{M}(n, n).$$

Proof. $[l]_{\mathcal{V}}^{\mathcal{E}_n} := [l(v^1), \dots, l(v^n)] = I$. ■

The double dual space of a vector space

Question. Take⁴ a basis \mathcal{V}^* of V^* . Is \mathcal{V}^* the dual basis of some basis \mathcal{V} of V ? Corollary 319 below will answer positively to that question.

Definition 314 The double dual V^{**} of V is the dual space of V^* , i.e.,

$$V^{**} := (V^*)^* := \mathcal{L}(V^*, \mathbb{R}).$$

Remark 315 If V has finite dimension, then

$$\dim V = \dim V^* = \dim V^{**}.$$

Proposition 316 For any $v \in V$, the function

$$L_v : V^* \rightarrow \mathbb{R}, \quad f \mapsto f(v)$$

is linear, i.e., $L_v \in V^{**}$.

Proof. For any $\alpha, \beta \in F$, $f_1, f_2 \in V^*$,

$$L_v(\alpha f_1 + \beta f_2) = (\alpha f_1 + \beta f_2)(v) = \alpha f_1(v) + \beta f_2(v) = \alpha L_{f_1} + \beta L_{f_2}.$$

■

Proposition 317 Let V be a finite dimensional vector space. Then

$$\varphi : V \rightarrow V^{**}, \quad v \mapsto L_v$$

is an isomorphism.

Proof. Step 1. φ is linear.

For any $\alpha, \beta \in F$, $v_1, v_2 \in V$, by definition of φ ,

$$\varphi(\alpha v_1 + \beta v_2) = L_{\alpha v_1 + \beta v_2},$$

and

$$\alpha \varphi(v_1) + \beta \varphi(v_2) = \alpha L_{v_1} + \beta L_{v_2}.$$

⁴Here I follow pages 107 and 108 in Hoffman and Kunze (1971).

Moreover, $\forall f \in V^*$,

$$L_{\alpha v_1 + \beta v_2}(f) := f(\alpha v_1 + \beta v_2) \stackrel{f \in V^*}{=} \alpha f(v_1) + \beta f(v_2) \stackrel{\text{def. } L_v}{=} \alpha L_{v_1} + \beta L_{v_2}.$$

Step 2. φ is nonsingular.

We want to show that $\ker \varphi = \{0\}$, i.e., $\langle v \neq 0 \Rightarrow \varphi(v) := L_v \neq 0 \rangle$, i.e., $\forall v \neq 0, \exists f \in V^*$ such that $f(v) \neq 0$. Take a basis $\mathcal{V} = \{v, v^i\}_{i=2}^n$ of V . The existence of such a basis is insured by Proposition 201. Then, from Proposition 311, there exists a unique dual basis $\mathcal{V}^* = (l_i)_{i=1}^n$ associated with \mathcal{V} , and $l_1(v) = 1 \neq 0$.

Step 3. Desired result.

From Remark 309, $n = \dim V = \dim V^* = \dim V^{**}$. Then, from Proposition 281, my EUI notes, since φ is nonsingular, it is invertible and therefore an isomorphism. ■

Corollary 318 *Let V be a finite dimensional vector space over a field F . If $L \in V^{**}$, then $\exists ! v \in V$ such that $\forall f \in V^*, L(f) = f(v)$.*

Proof. Such v is just $\varphi(L)$. ■

Corollary 319 *Let V be a finite dimensional vector space over a field F . Any basis of V^* is the unique dual basis of some basis of V .*

Proof. Take a basis $\{l_i\}_{i=1}^n$ of V^* . From Proposition 311, there exists a unique basis $\{L_i\}_{i=1}^n$ of V^{**} such that

$$\forall i, j \in \{1, \dots, n\}, L_i(f_j) = \delta_{ij}. \quad (7.13)$$

Then, from the previous Corollary,

$$\forall i \in \{1, \dots, n\}, \exists ! v^i \in V \text{ such that } \forall f \in V^*, L_i(f) = f(v^i). \quad (7.14)$$

Now, from the definition of L_{v^i} , we have

$$\forall f \in V^*, L_{v^i}(f) = f(v^i). \quad (7.15)$$

Then, from (7.14) and (7.15), we get

$$L_i = L_{v^i}. \quad (7.16)$$

Then,

$$\varphi^{-1}(L_i) \stackrel{(7.16)}{=} \varphi^{-1}(L_{v^i}) = v^i,$$

where the last equality follows from Proposition 317. Since $\{L_i\}_{i=1}^n$ is a basis of V^{**} and φ is an isomorphism, then from Proposition 264, $\{v^i\}_{i=1}^n$ is a basis of V . Moreover, from (7.14), $f_i(v^j) = L_i(f_j) \stackrel{(7.13)}{=} \delta_{ij}$, and then, by definition, $\{l_i\}_{i=1}^n$ is the unique dual basis associated with $\{v_i\}_{i=1}^n$. ■

A sometimes useful result is presented below.

Proposition 320 *If V is a vector space of dimension bigger than n , for any $i \in \{1, \dots, n\}$, $l_i \in \mathcal{L}(V, \mathbb{R})$,⁵ and $\{l_i\}_{i=1}^n$ is a linearly independent set, then $l : V \rightarrow \mathbb{R}^n, v \mapsto (l_i(v))_{i=1}^n$ is onto.*

Proof. Take an n dimensional vector subspace V' of V . From Corollary 319, there exists a basis $\mathcal{V} = \{v^i\}_{i=1}^n$ of V' such that $\{l_i\}_{i=1}^n$ is its unique associated dual basis. Then, by definition of dual basis, for any $i \in \{1, \dots, n\}$, $l(v^i) = e_n^i$. Therefore, for any $x \in \mathbb{R}^n$,

$$l\left(\sum_{i=1}^n x_i v^i\right) = \sum_{i=1}^n x_i e_n^i = x,$$

as desired. ■

⁵ $\mathcal{L}(V, \mathbb{R})$ is the vector space of linear functions from V to \mathbb{R} .

7.8.2 Vector spaces as Images or Kernels of well chosen linear functions

This section is preliminary and incomplete.

Proposition 321 *If L is a vector subspace of \mathbb{R}^S of dimension A , then*

1. a. $\exists l_1 \in L(\mathbb{R}^S, \mathbb{R}^{S-A})$ such that $\ker l_1 = L$;
- b. $\exists \sigma \in \Sigma$ and $\exists E \in M(S-A, A)$ such that

$$[l_1] = [I_{S-A}|E] \cdot P_\sigma \in \mathbb{M}(S-A, S);$$

- c. $E = \psi_\sigma(L)$ where ψ_σ is a chart of an atlas of $G_{S,A}$.

2. a. $\exists l_2 \in L(\mathbb{R}^A, \mathbb{R}^S)$ such that $\text{Im} l_2 = L$;

b. ⁶

$$[l_2] = P_{\sigma^{-1}} \begin{bmatrix} -E \\ I_{A \times A} \end{bmatrix}$$

and, therefore, if $\sigma = \text{id}$,

$$[l_2] = \begin{bmatrix} -E \\ I_A \end{bmatrix}_{S \times A}$$

3.

$$L = [I_{S-A}|E] \cdot P_\sigma = \text{Im} P_{\sigma^{-1}} \begin{bmatrix} -E \\ I_{A \times A} \end{bmatrix}$$

Moreover,

4. Let $M, M' \in M^f(S-A, S)$ be given. Then

$$\ker M = \ker M' \Leftrightarrow \text{there exists } B \in \mathbb{M}^f(S-A, S-A) \text{ such that } M' = BM.$$

5. Let $Y, Y' \in M^f(S, A)$ be given. Then

$$\text{Im} Y = \text{Im} Y' \Leftrightarrow \text{there exists a unique } C \in \mathbb{M}^f(A, A) \text{ such that } Y' = YC.$$

Proof. 1.

Take a basis $\{l^1, \dots, l^A\} \subseteq \mathbb{R}^S$ of L . Define

$$C = \begin{bmatrix} l^1 \\ \dots \\ l^A \end{bmatrix}_{S \times A},$$

where the vectors are taken to be row vectors. By definition of basis, $\text{rank } C = A$. From the Dimension Theorem, $\dim \mathbb{R}^S = \dim \text{Im} l_C + \dim \ker l_C$ and $S = A + \dim \ker l_C$, or $\dim \ker l_C = S - A$. Let $\{k^1, \dots, k^{S-A}\} \subseteq \mathbb{R}^S$ be a basis of $\ker l_C$. Then, $\forall s \in \{A, \dots, S-A\}$, $Ck^s = 0$, and

$$\forall a \in 1, \dots, A, \forall s \in \{1, \dots, S-A\}, l^a \cdot k^s = 0,$$

i.e.,⁷

$$L = (\ker l_C)^\perp. \quad (7.17)$$

Define

$$M = \begin{bmatrix} k^1 \\ \dots \\ k^{S-A} \end{bmatrix}_{(S-A) \times S}.$$

⁶Recall that $\forall \sigma \in \Sigma$,

$$P_{\sigma^{-1}} = P_\sigma^{-1} = P_\sigma^T.$$

⁷We are using the obvious fact that:

Two vector spaces V and W are orthogonal iff each element in a basis of V is orthogonal to each element in a basis of W .

By definition of basis, $\text{rank } M = S - A$. Moreover, from the fact that $\{k^1, \dots, k^{S-A}\} \subseteq \mathbb{R}^S$ be a basis of $\ker l_C$ and from (7.17), we have that

$$L = \{x \in \mathbb{R}^S : \forall s \in \{1, \dots, S - A\}, k^s \cdot x = 0\} = \{x \in \mathbb{R}^S : M \cdot x = 0\} = \ker l_M.$$

2.

a. Let $\{v^a\}_{a=1}^A$ be a basis of $L \subseteq \mathbb{R}^S$. Take $l_2 \in \mathcal{L}(\mathbb{R}^A, \mathbb{R}^S)$ such that

$$\forall a \in \{1, \dots, A\}, l_2(e_A^a) = v^a,$$

where e_A^a is the a -th element in the canonical basis in \mathbb{R}^A . From a standard Proposition in linear algebra⁸, such function does exists and, in fact, it is unique. Then, from the (linear algebra) dimension theorem

$$\dim \text{Im} l_2 = A - \dim \ker l_2 \leq A.$$

Moreover, $L = \text{span } \{v^a\}_{a=1}^A \subseteq \text{Im} l_2$ and $\dim L = A$. Therefore $\dim \text{Im} l_2 = A$, and $L = \text{Im} l_2$, as desired.

b. From Step 2a above, and by definition of $[l_2]$, we have that

$$[l_2] = [v^1, \dots, v^A] \in \mathbb{M}(S, A),$$

where $\{v^a\}_{a=1}^A$ is a basis of L and the vectors v^a are written as column vectors. Therefore, we are left with finding a basis of $L = \ker [I_{S-A}|E] \cdot P_\sigma$, which we claim is given by the A column vectors of

$$P_{\sigma^{-1}} \begin{bmatrix} -E \\ I_{A \times A} \end{bmatrix},$$

a fact which is proved below:

i. The vectors belong to $\ker [I_{S-A}|E] \cdot P_\sigma$:

$$[I|E] \cdot P_\sigma \cdot P_{\sigma^{-1}} \begin{bmatrix} -E \\ I \end{bmatrix} = [I|E] \cdot \begin{bmatrix} -E \\ I \end{bmatrix} = 0.$$

ii. the vectors are linearly independent:

$$\text{rank} P_{\sigma^{-1}} \begin{bmatrix} -E \\ I \end{bmatrix} \stackrel{(1)}{=} \text{rank} \begin{bmatrix} -E \\ I \end{bmatrix} = A,$$

where (1) follows from Corollary 7, page 10 in the 4-author book.

3.

It is an immediate consequence of 2. and 3. above.

4.

See Villanacci and others (2002), Proposition 39, page 388, or observe what follows

\Rightarrow

From, say, Theorem on page 41, in Ostaszewski (1990)⁹, we have that

$$(\ker Y)^\perp = \text{Im} Y^T,$$

(this result is proved, along other things, in my handwritten file some-facts-on-Stiefel-manifolds-2011-10-12.pdf

and therefore, using the assumption

$$\ker Y^T = (\text{Im} Y)^\perp = (\text{Im} Y')^\perp = \ker (Y')^T.$$

Then, from part 3 in the present Proposition,

$$\text{there exists } \widehat{B} \in \mathbb{M}^f(A, A) \text{ such that } Y'^T = \widehat{B} Y^T.$$

⁸Let V and U be finite dimensional vectors spaces such that $S = \{v^1, \dots, v^n\}$ is a basis of V and $\{u^1, \dots, u^n\}$ is a set of arbitrary vectors in U . Then there exists a unique linear function $l : V \rightarrow U$ such that $\forall i \in \{1, \dots, n\}$, $l(v^i) = u^i$ - see my math class notes, Proposition 273, page 82.

⁹Ostaszewski, A., (1990), *Advanced mathematical methods*, London School of Economics Mathematics Series, Cambridge University Press, Cambridge, UK

and

$$Y' = Y\widehat{B}^T,$$

and it is then enough to take $C = \widehat{B}^T$.

[\Leftarrow]

$\text{Im}Y \subseteq \text{Im}Y' : w \in \text{Im}Y \Rightarrow \exists z \in \mathbb{R}^A$ such that $w = Yz = Y'Cz \Rightarrow \exists z' = Cz \in \mathbb{R}^A$ such that $w = Y'z' \Leftrightarrow w \in \text{Im}Y'$.

$\text{Im}Y' \subseteq \text{Im}Y : w' \in \text{Im}Y' \Rightarrow \exists z' \in \mathbb{R}^A$ such that $w' = Y'z' = Y'CC^{-1}z' = YC^{-1}z' \Rightarrow \exists z = C^{-1}z' \in \mathbb{R}^A$ such that $w' = Yz \Leftrightarrow w' \in \text{Im}Y$.

Uniqueness. Since $Y \in \mathbb{M}^f(S, A)$, there exists $\sigma \in \Sigma(S)$ such that $P_\sigma Y = \begin{bmatrix} Y^* \\ \widehat{Y} \end{bmatrix}$, with $Y^* \in \mathbb{M}^f(A, A)$. Define also $P_\sigma Y' = \begin{bmatrix} Y'^* \\ \widehat{Y}' \end{bmatrix}$. Then, premultiplying $Y' = YC$ by P_σ , we get $Y'^* = Y^*C$ and $C = Y'^{-1}Y'^*$. ■

Chapter 8

Solutions to systems of linear equations

8.1 Some preliminary basic facts

Let's recall some basic definition from Section 1.6.

Definition 322 Consider the following linear system with m equations and n unknowns

$$\begin{cases} a_{11}x_1 + \cdots + a_{1n}x_n = b_1 \\ \vdots \\ a_{m1}x_1 + \cdots + a_{mn}x_n = b_m \end{cases}$$

which can be rewritten as

$$Ax = b$$

$A_{m \times n}$ is called matrix of the coefficients (or coefficient matrix) associated with the system and $M_{m \times (n+1)} = \begin{bmatrix} A & b \end{bmatrix}$ is called augmented matrix associated with the system.

Recall the following definition.

Definition 323 Two linear system are said to be equivalent if they have the same solutions.

Let's recall some basic facts we discussed in previous chapters.

Remark 324 It is well known that the following operations applied to a system of linear equations lead to an equivalent system:

- I) interchange two equations;
- II) multiply both sides of an equation by a nonzero real number;
- III) add left and right hand side of an equation to the left and right hand side of another equation;
- IV) change the place of the unknowns.

The transformations I), II), III) and IV) are said elementary transformations, and, as it is well known, they do not change the solution set of the system they are applied to.

Those transformations correspond to elementary operations on rows of M or columns of A in the way described below

- I) interchange two rows of M ;
- II) multiply a row of M by a nonzero real number;
- III) sum a row of M to another row of M ;
- IV) interchange two columns of A .

The above described operations do not change the rank of A and they do not change the rank of M - see Proposition 231.

Homogenous linear system.

Definition 325 A linear system for which $b = 0$, i.e., of the type

$$Ax = 0$$

with $A \in \mathbb{M}(m, n)$, is called homogenous system.

Remark 326 Obviously, 0 is a solution of the homogenous system. The set of solution of a homogenous system is $\ker l_A$. From Remark 306,

$$\dim \ker l_A = n - \text{rank } A.$$

8.2 A solution method: Rouché-Capelli's and Cramer's theorems

The solution method presented in this section is based on two basic theorems.

1. Rouché-Capelli's Theorem, which gives necessary and sufficient condition for the existence of solutions;
2. Cramer's Theorem, which gives a method to compute solutions - if they exist.

Theorem 327 (Rouché – Capelli) A system with m equations and n unknowns

$$A_{m \times n} x = b \tag{8.1}$$

has solutions

\Leftrightarrow

$$\text{rank } A = \text{rank} \begin{bmatrix} A & | & b \end{bmatrix}$$

Proof. $[\Rightarrow]$

Let x^* be a solution to 8.1. Then, b is a linear combination, via the solution x^* of the columns of A . Then, from Proposition 231,

$$\text{rank} \begin{bmatrix} A & | & b \end{bmatrix} = \text{rank} \begin{bmatrix} A & | & 0 \end{bmatrix} = \text{rank} \begin{bmatrix} A \end{bmatrix}.$$

$[\Leftarrow]$

1st proof.

We want to show that

$$\exists x \in \mathbb{R}^n \text{ such that } Ax^* = b, \text{ i.e., } b = \sum_{j=1}^n x_j^* \cdot C^j(A).$$

By assumption, $\text{rank } A = \text{rank} \begin{bmatrix} A & | & b \end{bmatrix} := r$. Since $\text{rank } A = r$, there are r linearly independent column vectors of A , say $\{C^j(A)\}_{j \in R}$, where $R \subseteq \{1, \dots, n\}$ and $\#R = r$.

Since $\text{rank} \begin{bmatrix} A & | & b \end{bmatrix} = r$, $\{C^j(A)\}_{j \in R} \cup \{b\}$ is a linearly dependent set and from Lemma 209, b is a linear combinations of the vectors in $\{C^j(A)\}_{j \in R}$, i.e., $\exists (x_j)_{j \in R}$ such that $b = \sum_{j \in R} x_j \cdot C^j(A)$ and

$$b = \sum_{j \in R} x_j \cdot C^j(A) + \sum_{j' \in \{1, \dots, n\} \setminus R} 0 \cdot C^{j'}(A).$$

Then, $x^* = (x_j^*)_{j=1}^n$ such that

$$x_j^* = \begin{cases} x_j & \text{if } j \in R \\ 0 & \text{if } j' \in \{1, \dots, n\} \setminus R \end{cases}$$

is a solution to $Ax = b$.

Second proof.

Since $\text{rank } A := r \leq \min \{m, n\}$, by definition of rank, there exists a rank r square submatrix A^* of A . From Remark 324, reordering columns of A and rows of $\begin{bmatrix} A & b \end{bmatrix}$ does not change the rank of A or $\begin{bmatrix} A & b \end{bmatrix}$ and leads to the following system, which is equivalent to $Ax = b$:

$$\begin{bmatrix} A^* & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} x^1 \\ x^2 \end{bmatrix} = \begin{bmatrix} b' \\ b'' \end{bmatrix},$$

where $A_{12} \in \mathbb{M}(r, n-r)$, $A_{21} \in \mathbb{M}(m-r, r)$, $A_{22} \in \mathbb{M}(m-r, n-r)$, $x^1 \in \mathbb{R}^r$, $x^2 \in \mathbb{R}^{n-r}$, $b' \in \mathbb{R}^r$, $b'' \in \mathbb{R}^{m-r}$,

(x^1, x^2) has been obtained from x performing on it the same permutations performed on the columns of A ,

(b', b'') has been obtained from b performing on it the same permutations performed on the rows of A .

Since

$$\text{rank} \begin{bmatrix} A^* & A_{12} & b' \end{bmatrix} = \text{rank } A^* = r,$$

the r rows of $\begin{bmatrix} A^* & A_{12} & b' \end{bmatrix}$ are linearly independent. Since

$$\text{rank} \begin{bmatrix} A^* & A_{12} & b' \\ A_{21} & A_{22} & b'' \end{bmatrix} = r,$$

the rows of that matrix are linearly dependent and from Lemma 209, the last $m-r$ rows of $[A|b]$ are linear combinations of the first r rows. Therefore, using again Remark 324, we have that $Ax = b$ is equivalent to

$$\begin{bmatrix} A^* & A_{12} \end{bmatrix} \begin{bmatrix} x^1 \\ x^2 \end{bmatrix} = b'$$

or, using Remark 74.2,

$$A^* x^1 + A_{12} x^2 = b'$$

and

$$x^1 = (A^*)^{-1} (b' - A_{12} x^2) \in \mathbb{R}^r$$

while x_2 can be chosen arbitrarily; more precisely

$$\left\{ (x^1, x^2) \in \mathbb{R}^n : x_1 = (A^*)^{-1} (b' - A_{12} x^2) \in \mathbb{R}^r \text{ and } x_2 \in \mathbb{R}^{n-r} \right\}$$

is the nonempty set of solution to the system $Ax = b$. ■

Theorem 328 (Cramer) *A system with n equations and n unknowns*

$$A_{n \times n} x = b$$

with $\det A \neq 0$, has a unique solution $x = (x_1, \dots, x_i, \dots, x_n)$ where for $i \in \{1, \dots, n\}$,

$$x_i = \frac{\det A_i}{\det A}$$

and A_i is the matrix obtained from A substituting the column vector b in the place of the i -th column.

Proof. since $\det A \neq 0$, A^{-1} exists and it is unique. Moreover, from $Ax = b$, we get $A^{-1}Ax = A^{-1}b$ and

$$x = A^{-1}b$$

Moreover

$$A^{-1}b = \frac{1}{\det A} \text{Adj } A \cdot b$$

It is then enough to verify that

$$\text{Adj } A \cdot b = \begin{bmatrix} \det A_1 \\ \dots \\ \det A_i \\ \dots \\ \det A_n \end{bmatrix}$$

which we omit (see Exercise 7.34, page 268, in Lipschutz (1991)). ■

The combinations of Rouché-Capelli and Cramer's Theorem allow to give a method to solve any linear system - apart from computational difficulties.

Remark 329 Rouché'-Capelli and Cramer's Theorem based method.

Let the following system with m equations and n unknowns be given:

$$A_{m \times n} x = b.$$

0. Simplify the system using elementary row operations on $[A \mid b]$ and elementary column operations on A . Those operation do not change rank A , rank B , set of solution to system $Ax = b$.

1. Compute rank A and rank $[A \mid b]$.

i. If

$$\text{rank } A \neq \text{rank } [A \mid b],$$

then the system has no solution.

ii. If

$$\text{rank } A = \text{rank } [A \mid b] := r,$$

then the system has solutions which can be computed as follows.

2. Extract a square r -dimensional invertible submatrix A_r from A .

i. Discard the equations, if any, whose corresponding rows are not part of A_r .

ii. In the remaining equations, bring on the right hand side the terms containing unknowns whose coefficients are not part of the matrix A_r , if any.

iii. You then get a system to which Cramer's Theorem can be applied, treating as constant the expressions on the right hand side and which contain $n - r$ unknowns. Those unknowns can be chosen arbitrarily. Sometimes it is said that then the system has " ∞^{n-r} " solutions or that the system admits $n - r$ degrees of freedom. More formally, we can say what follows.

Definition 330 Given $S, T \subseteq \mathbb{R}^n$, we define the sum of the sets S and T , denoted by $S + T$, as follows

$$\{x \in \mathbb{R}^n : \exists s \in S, \exists t \in T \text{ such that } x = s + t\}.$$

Proposition 331 Assume that the set S of solutions to the system $Ax = b$ is nonempty and let $x^* \in S$. Then

$$S = \{x^*\} + \ker l_A := \{x \in \mathbb{R}^n : \exists x^0 \in \ker l_A \text{ such that } x = x^* + x^0\}$$

Proof. $[\subseteq]$

Take $x \in S$. We want to find $x^0 \in \ker A$ such that $x = x^* + x^0$. Take $x^0 = x - x^*$. Clearly $x = x^* + (x - x^*)$. Moreover,

$$Ax_0 = A(x - x^*) \stackrel{(1)}{=} b - b = 0 \quad (8.2)$$

where (1) follows from the fact that $x, x^* \in S$.

$[\supseteq]$

Take $x = x^* + x^0$ with $x^* \in S$ and $x^0 \in \ker A$. Then

$$Ax = Ax^* + Ax_0 = b + 0 = b.$$

■

Remark 332 The above proposition implies that a linear system either has no solutions, or has a unique solution, or has infinite solutions.

Definition 333 V is an affine subspace of \mathbb{R}^n if there exists a vector subspace W of \mathbb{R}^n and a vector $x \in \mathbb{R}^n$ such that

$$V = \{x\} + W$$

We say that the¹ dimension of the affine subspace V is $\dim W$.

¹If W' and W'' are vector subspaces of \mathbb{R}^n , $x \in \mathbb{R}^n$ and $V := \{x\} + W' = \{x\} + W''$, then $W' = W''$.

Take $w \in W'$, then $x + w \in V = \{x\} + W''$. Then there exists $x \in \{x\}$ and $\hat{w} \in W''$ such that $x + w = x + \hat{w}$. Then $w = \hat{w} \in W''$, and $W' \subseteq W''$. Similar proof applies for the opposite inclusion.

Remark 334 Let $a \in \mathbb{R}$ be given. First of all observe that

$$\{(x, y) \in \mathbb{R}^2 : y = ax\} = \ker l,$$

where $l \in \mathcal{L}(\mathbb{R}^2, \mathbb{R})$ and $l(x, y) = ax - y$. Let's present a geometric description of Proposition 331. We want to verify that the following two sets are equal.

$$S := \{(x_0, y_0)\} + \{(x, y) \in \mathbb{R}^2 : y = ax\},$$

$$T := \{(x, y) \in \mathbb{R}^2 : y = a(x - x_0) + y_0\}.$$

In words, we want to verify that the affine space “ $\{(x_0, y_0)\}$ plus $\ker l$ ” is nothing but the set of points belonging to the line with slope a and going through the point (x_0, y_0) .

$S \subseteq T$. Take $(x', y') \in S$; then $\exists x'' \in \mathbb{R}$ such that $x' = x_0 + x''$ and $y' = y_0 + ax''$. We have to check that $y' = a(x' - x_0) + y_0$. Indeed, $a(x' - x_0) + y_0 = a(x_0 + x'' - x_0) + y_0 = ax'' + y_0$.

$T \subseteq S$. Take (x', y') such that $y' = a(x' - x_0) + y_0$. Take $x'' = x' - x_0$. Then $x' = x_0 + x''$ and $y' = y_0 + ax''$, as desired.

Remark 335 Since $\dim \ker l_A = n - \text{rank } A$, the above Proposition and Definition say that if a nonhomogeneous system has solutions, then the set of solutions is an affine space of dimension $n - \text{rank } A$.

Exercise 336 Apply the algorithm described in Remark 329 to solve the following linear system.

$$\begin{cases} x_1 + 2x_2 + 3x_3 = 1 \\ 4x_1 + 5x_2 + 6x_3 = 2 \\ 5x_1 + 7x_2 + 9x_3 = 3 \end{cases}$$

The associated matrix $[A \mid b]$ is

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 1 \\ 4 & 5 & 6 & 2 \\ 5 & 7 & 9 & 3 \end{array} \right]$$

1. Since the third row of the matrix $[A \mid b]$ is equal to the sum of the first two rows, and since

$$\det \begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix} = 5 - 8 = -3,$$

we have that

$$\text{rank } A = \text{rank } [A \mid b] = 2,$$

and the system has solutions.

2. Define

$$A_2 = \begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix}$$

i. Discarding the equations, whose corresponding rows are not part of A_2 and, in the remaining equations, bringing on the right hand side the terms containing unknowns whose coefficients are not part of the matrix A_2 , we get

$$\begin{cases} x_1 + 2x_2 = 1 - 3x_3 \\ 4x_1 + 5x_2 = 2 - 6x_3 \\ 5x_1 + 7x_2 = 3 - 9x_3 \end{cases}$$

iii. Then, using Cramer's Theorem, recalling that $\det A_2 = -3$, we get

$$x_1 = \frac{\det \begin{bmatrix} 1 - 3x_3 & 2 \\ 2 - 6x_3 & 5 \end{bmatrix}}{-3} = x_3 - \frac{1}{3},$$

$$x_2 = \frac{\det \begin{bmatrix} 1 & 1 - 3x_3 \\ 4 & 2 - 6x_3 \end{bmatrix}}{-3} = \frac{2}{3} - 2x_3,$$

and the solution set is

$$\left\{ (x_1, x_2, x_3) \in \mathbb{R}^3 : x_1 = x_3 - \frac{1}{3}, x_2 = \frac{2}{3} - 2x_3 \right\}.$$

Remark 337 *How to find a basis of \ker .*

Let $A \in \mathbb{M}(m, n)$ be given and $\text{rank} A = r \leq \min\{m, n\}$. Then, from the second proof of Rouche'-Capelli's Theorem, we have that system

$$Ax = 0$$

admits the following set of solutions

$$\left\{ (x^1, x^2) \in \mathbb{R}^r \times \mathbb{R}^{n-r} : x^1 = (A^*)^{-1} (-A_{12} \cdot x^2) \right\} \quad (8.3)$$

Observe that $\dim \ker l_A = n - r := p$. Then, a basis of $\ker l_A$ is

$$\mathcal{B} = \left\{ \begin{bmatrix} -[A^*]^{-1} A_{12} \cdot e_p^1 \\ e_p^1 \end{bmatrix}, \dots, \begin{bmatrix} -[A^*]^{-1} A_{12} \cdot e_p^p \\ e_p^p \end{bmatrix} \right\}.$$

To check the above statement, we check that 1. $\mathcal{B} \subseteq \ker l_A$, and 2. \mathcal{B} is linearly independent².

1. It follows from (8.3);

2. It follows from the fact that $\det [e_p^1, \dots, e_p^p] = \det I = 1$.

Example 338

$$\begin{cases} x_1 + x_2 + x_3 + 2x_4 = 0 \\ x_1 - x_2 + x_3 + 2x_4 = 0 \end{cases}$$

Defined

$$A^* = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}, \quad A_{12} = \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix}$$

the starting system can be rewritten as

$$\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = - \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} x_3 \\ x_4 \end{bmatrix}.$$

Then a basis of \ker is

$$\begin{aligned} \mathcal{B} &= \left\{ \begin{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \end{bmatrix} \\ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \end{bmatrix}, \begin{bmatrix} - \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 2 \\ 1 & 2 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} \\ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{bmatrix} \right\} = \\ &= \left\{ \begin{bmatrix} -1 \\ 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -2 \\ 0 \\ 0 \\ 1 \end{bmatrix} \right\} \end{aligned}$$

Example 339 *Discuss the following system (i.e., say if admits solutions).*

$$\begin{cases} x_1 + x_2 + x_3 = 4 \\ x_1 + x_2 + 2x_3 = 8 \\ 2x_1 + 2x_2 + 3x_3 = 12 \end{cases}$$

The augmented matrix $[A|b]$ is:

$$\left[\begin{array}{ccc|c} 1 & 1 & 1 & 4 \\ 1 & 1 & 2 & 8 \\ 2 & 2 & 3 & 12 \end{array} \right]$$

$$\text{rank } [A|b] = \text{rank} \left[\begin{array}{ccc|c} 1 & 1 & 1 & 4 \\ 1 & 1 & 2 & 8 \\ 0 & 0 & 0 & 0 \end{array} \right] = 2 = \text{rank } A$$

²Observe that \mathcal{B} is made up by p vectors and $\dim \ker l_A = p$.

From Step 2 of Rouché'-Capelli and Cramer's method, we can consider the system

$$\begin{cases} x_2 + x_3 = 4 - x_1 \\ x_2 + 2x_3 = 8 - x_1 \end{cases}$$

Therefore, x_1 can be chosen arbitrarily and since $\det \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} = 1$,

$$x_2 = \det \begin{bmatrix} 4 - x_1 & 1 \\ 8 - x_1 & 2 \end{bmatrix} = -x_1$$

$$x_3 = \det \begin{bmatrix} 1 & 4 - x_1 \\ 1 & 8 - x_1 \end{bmatrix} = 4$$

Therefore, the set of solution is

$$\{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_2 = -x_1, x_3 = 4\}$$

Example 340 Discuss the following system

$$\begin{cases} x_1 + x_2 = 2 \\ x_1 - x_2 = 0 \end{cases}$$

The augmented matrix $[A|b]$ is:

$$\left[\begin{array}{cc|c} 1 & 1 & 2 \\ 1 & -1 & 0 \end{array} \right]$$

Since $\det A = -1 - 1 = -2 \neq 0$,

$$\text{rank}[A|b] = \text{rank}A = 2$$

and the system has a unique solution:

$$x_1 = \frac{\det \begin{bmatrix} 2 & 1 \\ 0 & -1 \end{bmatrix}}{-2} = \frac{-2}{-2} = 1$$

$$x_2 = \frac{\det \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}}{-2} = \frac{-2}{-2} = 1$$

Therefore, the set of solution is

$$\{(1, 1)\}$$

Example 341 Discuss the following system

$$\begin{cases} x_1 + x_2 = 2 \\ x_1 + x_2 = 0 \end{cases}$$

The augmented matrix $[A|b]$ is:

$$\left\{ \begin{array}{cc|c} 1 & 1 & 2 \\ 1 & 1 & 0 \end{array} \right\}$$

Since $\det A = 1 - 1 = 0$, and $\det \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix} = -2 \neq 0$, we have that

$$\text{rank}[A|b] = 2 \neq 1 = \text{rank}A$$

and the system has no solutions. Therefore, the set of solution is \emptyset .

Example 342 Discuss the following system

$$\begin{cases} x_1 + x_2 = 2 \\ 2x_1 + 2x_2 = 4 \end{cases}$$

The augmented matrix $[A|b]$ is:

$$\left[\begin{array}{cc|c} 1 & 1 & 2 \\ 2 & 2 & 4 \end{array} \right]$$

From rank properties,

$$\begin{aligned} \text{rank} \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix} &= \text{rank} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} = 1 \\ \text{rank} \begin{bmatrix} 1 & 1 & 2 \\ 2 & 2 & 4 \end{bmatrix} &= \text{rank} \begin{bmatrix} 1 & 1 & 2 \\ 0 & 0 & 0 \end{bmatrix} = 1 \end{aligned}$$

Recall that elementary operations of **rows** on the augmented matrix do not change the rank of either the augmented or coefficient matrices.

Therefore

$$\text{rank}[A|b] = 1 = \text{rank}A$$

and the system has infinite solutions. More precisely, the set of solutions is

$$\{(x_1, x_2) \in \mathbb{R}^2 : x_1 = 2 - x_2\}$$

Example 343 Say for which value of the parameter $k \in \mathbb{R}$ the following system has one, infinite or no solutions:

$$\begin{cases} (k-1)x + (k+2)y = 1 \\ -x + ky = 1 \\ x - 2y = 1 \end{cases}$$

$$[A|b] = \left[\begin{array}{cc|c} k-1 & k+2 & 1 \\ -1 & k & 1 \\ 1 & -2 & 1 \end{array} \right]$$

$$\det[A|b] = \det \begin{bmatrix} k-1 & k+2 & 1 \\ -1 & k & 1 \\ 1 & -2 & 1 \end{bmatrix} = \det \begin{bmatrix} -1 & k \\ 1 & -2 \end{bmatrix} - \det \begin{bmatrix} k-1 & k+2 \\ 1 & -2 \end{bmatrix} + \det \begin{bmatrix} k-1 & k+2 \\ -1 & k \end{bmatrix} =$$

$$(2-k) - (-2k+2-k-2) + (k^2-k+k+2) = 2-k+2k+k+k^2+2 = 2k+k^2+4$$

$\Delta = -1 - 17 < 0$. Therefore, the determinant is never equal to zero and $\text{rank}[A|b] = 3$. Since $\text{rank}A_{3 \times 2} \leq 2$, the solution set of the system is empty of each value of k .

Remark 344 To solve a parametric linear system $Ax = b$, where $A \in \mathbb{M}(m, n) \setminus \{0\}$, it is convenient to proceed as follows.

1. Perform “easy” row operations on $[A|b]$;
2. Compute $\min\{m, n+1\} := k$ and consider the $k \times k$ submatrices of the matrix $[A|b]$.

There are two possibilities.

Case 1. There exists a $k \times k$ submatrix whose determinant is different from zero for some values of the parameters;

Case 2. All $k \times k$ submatrices have zero determinant for each value of the parameters.

If Case 2 occurs, then at least one row of $[A|b]$ is a linear combinations of other rows; therefore you can eliminate it. We can therefore assume that we are in Case 1. In that case, proceed as described below.

3. among those $k \times k$ submatrices, choose a matrix A^* which is a submatrix of A , if possible; if you have more than one matrix to choose among, choose “the easiest one” from a computational viewpoint, i.e., that one with highest number of zeros, the lowest number of times a parameters appear, ... ;

4. compute $\det A^*$, a function of the parameter;

5. analyze the cases $\det A^* \neq 0$ and possibly $\det A^* = 0$.

Example 345 Say for which value of the parameter $a \in \mathbb{R}$ the following system has one, infinite or no solutions:

$$\begin{cases} ax_1 + x_2 + x_3 &= 2 \\ x_1 - ax_2 &= 0 \end{cases}$$

Example 346 Say for which value of the parameter $a \in \mathbb{R}$ the following system has one, infinite or no solutions:

$$\begin{cases} ax_1 + x_2 + x_3 &= 2 \\ x_1 - ax_2 &= 0 \\ 2ax_1 + ax_2 &= 4 \end{cases}$$

$$[A|b] = \begin{bmatrix} a & 1 & 1 & 2 \\ 1 & -a & 0 & 0 \\ 2a & a & 0 & 4 \end{bmatrix}$$

$$\det \begin{bmatrix} a & 1 & 1 \\ 1 & -a & 0 \\ 2a & a & 0 \end{bmatrix} = a + 2a^2 = 0 \quad \text{if} \quad a = 0, -\frac{1}{2}.$$

Therefore, if $a \neq 0, -\frac{1}{2}$,

$$\text{rank}[A|b] = \text{rank} A = 3$$

and the system has a unique solution.

If $a = 0$,

$$[A|b] = \begin{bmatrix} 0 & 1 & 1 & 2 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 4 \end{bmatrix}$$

Since $\det \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = -1$, $\text{rank} A = 2$. On the other hand,

$$\det \begin{bmatrix} 0 & 1 & 2 \\ 1 & 0 & 0 \\ 0 & 0 & 4 \end{bmatrix} = -1 \det \begin{bmatrix} 1 & 2 \\ 0 & 4 \end{bmatrix} = -4 \neq 0$$

and therefore $\text{rank}[A|b] = 3$, and

$$\text{rank}[A|b] = 3 \neq 2 = \text{rank} A$$

and the system has no solutions.

If $a = -\frac{1}{2}$,

$$[A|b] = \begin{bmatrix} -\frac{1}{2} & 1 & 1 & 2 \\ 1 & \frac{1}{2} & 0 & 0 \\ -1 & -\frac{1}{2} & 0 & 4 \end{bmatrix}$$

Since

$$\det \begin{bmatrix} 1 & 1 \\ \frac{1}{2} & 0 \end{bmatrix} = -\frac{1}{2},$$

$$\text{rank} A = 2.$$

Since

$$\det \begin{bmatrix} -\frac{1}{2} & 1 & 2 \\ 1 & 0 & 0 \\ 1 & 0 & 4 \end{bmatrix} = -\det \begin{bmatrix} 1 & 2 \\ 0 & 4 \end{bmatrix} = -4$$

and therefore

$$\text{rank}[A|b] = 3,$$

$$\text{rank}[A|b] = 3 \neq 2 = \text{rank} A$$

and the system has no solutions.

Example 347 Say for which value of the parameter $a \in \mathbb{R}$ the following system has one, infinite or no solutions:

$$\begin{cases} (a+1)x_1 + (-2)x_2 + 2ax_3 = a \\ ax_1 + (-a)x_2 + x_3 = -2 \\ 2 + (2a-4)x_2 + (4a-2)x_3 = 2a+4 \end{cases}$$

Then,

$$[A|b] = \left[\begin{array}{ccc|c} a+1 & -2 & 2a & a \\ a & -a & 1 & -2 \\ 2 & 2a-4 & 4a-2 & 2a+4 \end{array} \right].$$

It is easy to see that we are in Case 2 described in Remark 344: all 3×3 submatrices of $[A|b]$ have determinant equal to zero - indeed, the last row is equal to 2 times the first row plus (-2) times the second row. We can then erase the third equation/row to get the following system and matrix.

$$\begin{cases} (a+1)x_1 + (-2)x_2 + 2ax_3 = a \\ ax_1 + (-a)x_2 + x_3 = -2 \end{cases}$$

$$[A|b] = \left[\begin{array}{ccc|c} a+1 & -2 & 2a & a \\ a & -a & 1 & -2 \end{array} \right]$$

$$\det \begin{bmatrix} -2 & 2a \\ -a & 1 \end{bmatrix} = 2a^2 - 2 = 0,$$

whose solutions are $-1, 1$. Therefore, if $a \in \mathbb{R} \setminus \{-1, 1\}$,

$$\text{rank } [A|b] = 2 = \text{rank } A$$

and the system has infinite solutions. Let's study the system for $a \in \{-1, 1\}$.

If $a = -1$, we get

$$[A|b] = \left[\begin{array}{ccc|c} 0 & -2 & -2 & -1 \\ -1 & 1 & 1 & -2 \end{array} \right]$$

and since

$$\det \begin{bmatrix} 0 & -2 \\ -1 & 1 \end{bmatrix} = 2 \neq 0$$

we have again

$$\text{rank } [A|b] = 2 = \text{rank } A$$

If $a = 1$, we have

$$[A|b] = \left[\begin{array}{ccc|c} 2 & -2 & 2 & 1 \\ 1 & -1 & 1 & -2 \end{array} \right]$$

and

$$\text{rank } [A|b] = 2 > 1 = \text{rank } A$$

and the system has no solution..

Example 348 Say for which value of the parameter $a \in \mathbb{R}$ the following system has one, infinite or no solutions:

$$\begin{cases} ax_1 + x_2 = 1 \\ x_1 + x_2 = a \\ 2x_1 + x_2 = 3a \\ 3x_1 + 2x_2 = a \end{cases}$$

$$[A|b] = \left[\begin{array}{cc|c} a & 1 & 1 \\ 1 & 1 & a \\ 2 & 1 & 3a \\ 3 & 2 & a \end{array} \right]$$

Observe that

$$\text{rank } A_{4 \times 2} \leq 2.$$

$$\det \begin{bmatrix} 1 & 1 & a \\ 2 & 1 & 3a \\ 3 & 2 & a \end{bmatrix} = 3a = 0 \quad \text{if} \quad a = 0.$$

Therefore, if $a \in \mathbb{R} \setminus \{0\}$,

$$\text{rank}[A|b] = 3 > 2 \geq \text{rank } A_{4 \times 2}$$

If $a = 0$,

$$[A|b] = \left[\begin{array}{cc|c} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 2 & 1 & 0 \\ 3 & 2 & 0 \end{array} \right]$$

and since

$$\det \begin{bmatrix} 0 & 1 & 1 \\ 1 & 1 & 0 \\ 2 & 1 & 0 \end{bmatrix} = -1$$

the system has no solution for $a = 0$.

Summarizing, $\forall a \in \mathbb{R}$, the system has no solutions.

8.3 Exercises

Problem sets: 20,23,24.

From Lipschutz (1991), page 179: 5.56; starting from page 263, 7.17 \rightarrow 7.20.

Part II

Some topology in metric spaces

Chapter 9

Metric spaces

9.1 Definitions and examples

Definition 349 Let X be a nonempty set. A metric or distance on X is a function $d : X \times X \rightarrow \mathbb{R}$ such that $\forall x, y, z \in X$

1. (a.) $d(x, y) \geq 0$, and (b.) $d(x, y) = 0 \Leftrightarrow x = y$,
 2. $d(x, y) = d(y, x)$,
 3. $d(x, z) \leq d(x, y) + d(y, z)$ (Triangle inequality).
- (X, d) is called a metric space.

Remark 350 Observe that the definition requires that $\forall x, y \in X$, it must be the case that $d(x, y) \in \mathbb{R}$.

Example 351 n -dimensional Euclidean space with Euclidean metric.
Given $n \in \mathbb{N}$, take $X = \mathbb{R}^n$, and

$$d_{2,n} : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}, \quad (x, y) \mapsto \left(\sum_{i=1}^n (x_i - y_i)^2 \right)^{\frac{1}{2}}.$$

$(X, d_{2,n})$ was shown to be a metric space in Proposition 58, Section 2.3. $d_{2,n}$ is called the Euclidean distance in \mathbb{R}^n . In what follows, unless needed, we write simply d_2 in the place of $d_{2,n}$.

Proposition 352 (Discrete metric space) Given a nonempty set X and the function

$$d : X^2 \rightarrow \mathbb{R}, \quad d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y, \end{cases}$$

(X, d) is a metric space, called discrete metric space.

Proof. 1a. $0, 1 \geq 0$.

1b. From the definition, $d(x, y) = 0 \Leftrightarrow x = y$.

2. It follows from the fact that $x = y \Leftrightarrow y = x$ and $x \neq y \Leftrightarrow y \neq x$.

3. If $x = z$, the result follows. If $x \neq z$, then it cannot be $x = y$ and $y = z$, and again the result follows. ■

Proposition 353 Given $n \in \mathbb{N}, p \in [1, +\infty)$, $X = \mathbb{R}^n$,

$$d : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}, \quad (x, y) \mapsto \left(\sum_{i=1}^n |x_i - y_i|^p \right)^{\frac{1}{p}},$$

(X, d) is a metric space.

Proof. 1a. It follows from the definition of absolute value.

1b. [\Leftarrow] Obvious.

[\Rightarrow] $(\sum_{i=1}^n |x_i - y_i|^p)^{\frac{1}{p}} = 0 \Leftrightarrow \sum_{i=1}^n |x_i - y_i|^p = 0 \Rightarrow \text{for any } i, |x_i - y_i| = 0 \Rightarrow \text{for any } i, x_i - y_i = 0.$

2. It follows from the fact $|x_i - y_i| = |y_i - x_i|$.

3. First of all observe that

$$d(x, z) = \left(\sum_{i=1}^n |(x_i - y_i) + (y_i - z_i)|^p \right)^{\frac{1}{p}}.$$

Then, it is enough to show that

$$\left(\sum_{i=1}^n |(x_i - y_i) + (y_i - z_i)|^p \right)^{\frac{1}{p}} \leq \left(\sum_{i=1}^n |x_i - y_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |y_i - z_i|^p \right)^{\frac{1}{p}}$$

which is a consequence of Proposition 354 below. ■

Proposition 354 Taken $n \in \mathbb{N}, p \in [1, +\infty), X = \mathbb{R}^n, a, b \in \mathbb{R}^n$

$$\left(\sum_{i=1}^n |a_i + b_i|^p \right)^{\frac{1}{p}} \leq \left(\sum_{i=1}^n |a_i|^p \right)^{\frac{1}{p}} + \left(\sum_{i=1}^n |b_i|^p \right)^{\frac{1}{p}}$$

Proof. It follows from the proof of the Proposition 357 below. ■

Definition 355 Let \mathbb{R}^∞ be the set of sequences in \mathbb{R} .

Definition 356 For any $p \in [1, +\infty)$, define¹

$$l^p = \left\{ (x_n)_{n \in \mathbb{N}} \in \mathbb{R}^\infty : \sum_{n=1}^{+\infty} |x_n|^p < +\infty \right\},$$

i.e., roughly speaking, l^p is the set of sequences whose associated series are absolutely convergent.

Proposition 357 (Minkowski inequality). $\forall (x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \in l^p, \forall p \in [1, +\infty)$,

$$\left(\sum_{n=1}^{+\infty} |x_n + y_n|^p \right)^{\frac{1}{p}} \leq \left(\sum_{n=1}^{+\infty} |x_n|^p \right)^{\frac{1}{p}} + \left(\sum_{n=1}^{+\infty} |y_n|^p \right)^{\frac{1}{p}}. \quad (9.1)$$

Proof. If either $(x_n)_{n \in \mathbb{N}}$ or $(y_n)_{n \in \mathbb{N}}$ are such that $\forall n \in \mathbb{N}, x_n = 0$ or $\forall n \in \mathbb{N}, y_n = 0$, i.e., if either sequence is the constant sequence of zeros, then (9.1) is trivially true.

Then, we can consider the case in which

$$\exists \alpha, \beta \in \mathbb{R}_{++} \text{ such that } \left(\sum_{n=1}^{+\infty} |x_n|^p \right)^{\frac{1}{p}} = \alpha \text{ and } \left(\sum_{n=1}^{+\infty} |y_n|^p \right)^{\frac{1}{p}} = \beta. \quad (9.2)$$

Define

$$\forall n \in \mathbb{N}, \quad \hat{x}_n = \left(\frac{|x_n|}{\alpha} \right)^p \text{ and } \hat{y}_n = \left(\frac{|y_n|}{\beta} \right)^p. \quad (9.3)$$

Then

$$\sum_{n=1}^{+\infty} \hat{x}_n = \sum_{n=1}^{+\infty} \hat{y}_n = 1. \quad (9.4)$$

For any $n \in \mathbb{N}$, from the triangle inequality for the absolute value, we have

$$|x_n + y_n| \leq |x_n| + |y_n|;$$

¹For basic results on series, see, for example, Section 10.5 in Apostol (1967).

since $\forall p \in [1, +\infty)$, $f_p : \mathbb{R}_+ \rightarrow \mathbb{R}$, $f_p(t) = t^p$ is an increasing function, we have

$$|x_n + y_n|^p \leq (|x_n| + |y_n|)^p. \quad (9.5)$$

Moreover, from (9.3),

$$(|x_n| + |y_n|)^p = \left(\alpha |\widehat{x}_n|^{\frac{1}{p}} + \beta |\widehat{y}_n|^{\frac{1}{p}} \right)^p = (\alpha + \beta)^p \left(\left(\frac{\alpha}{\alpha + \beta} |\widehat{x}_n|^{\frac{1}{p}} + \frac{\beta}{\alpha + \beta} |\widehat{y}_n|^{\frac{1}{p}} \right)^p \right). \quad (9.6)$$

Since $\forall p \in [1, +\infty)$, f_p is convex (just observe that $f_p''(t) = p(p-1)t^{p-2} \geq 0$), we get

$$\left(\frac{\alpha}{\alpha + \beta} |\widehat{x}_n| + \frac{\beta}{\alpha + \beta} |\widehat{y}_n| \right)^p \leq \frac{\alpha}{\alpha + \beta} |\widehat{x}_n| + \frac{\beta}{\alpha + \beta} |\widehat{y}_n| \quad (9.7)$$

From (9.5), (9.6) and (9.7), we get

$$|x_n + y_n|^p \leq (\alpha + \beta) \cdot \left(\frac{\alpha}{\alpha + \beta} |\widehat{x}_n|^p + \frac{\beta}{\alpha + \beta} |\widehat{y}_n|^p \right).$$

From the above inequalities and basic properties of the series, we then get

$$\sum_{n=1}^{+\infty} |x_n + y_n|^p \leq (\alpha + \beta)^p \left(\frac{\alpha}{\alpha + \beta} \sum_{n=1}^{+\infty} |\widehat{x}_n|^p + \frac{\beta}{\alpha + \beta} \sum_{n=1}^{+\infty} |\widehat{y}_n|^p \right) \stackrel{(9.4)}{=} (\alpha + \beta)^p \left(\frac{\alpha}{\alpha + \beta} + \frac{\beta}{\alpha + \beta} \right) = (\alpha + \beta)^p.$$

Therefore, using (9.2), we get

$$\sum_{n=1}^{+\infty} |x_n + y_n|^p \leq \left(\left(\sum_{n=1}^{+\infty} |x_n|^p \right)^{\frac{1}{p}} + \left(\sum_{n=1}^{+\infty} |y_n|^p \right)^{\frac{1}{p}} \right)^p,$$

and therefore the desired result. ■

Proposition 358 (l^p, d_p) with

$$d_p : l^p \times l^p \rightarrow \mathbb{R}, \quad d_p((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) = \left(\sum_{n=1}^{+\infty} |x_n - y_n|^p \right)^{\frac{1}{p}}.$$

is a metric space.

Proof. We first of all have to check that $d_p(x, y) \in \mathbb{R}$, i.e., that $\left(\sum_{n=1}^{+\infty} |x_n - y_n|^p \right)^{\frac{1}{p}}$ converges.

$$\left(\sum_{n=1}^{+\infty} |x_n - y_n|^p \right)^{\frac{1}{p}} = \left(\sum_{n=1}^{+\infty} |x_n + (-y_n)|^p \right)^{\frac{1}{p}} \leq \left(\sum_{n=1}^{+\infty} |x_n|^p \right)^{\frac{1}{p}} + \left(\sum_{n=1}^{+\infty} |y_n|^p \right)^{\frac{1}{p}} < +\infty,$$

where the first inequality follows from Minkowski inequality and the second inequality from the assumption that we are considering sequences in l^p .

Properties 1 and 2 of the distance follow easily from the definition. Property 3 is again a consequence of Minkowski inequality:

$$d_p(x, z) = \left(\sum_{n=1}^{+\infty} |(x_n - y_n) + (y_n - z_n)|^p \right)^{\frac{1}{p}} \leq \left(\sum_{n=1}^{+\infty} |x_n - y_n|^p \right)^{\frac{1}{p}} + \left(\sum_{n=1}^{+\infty} |y_n - z_n|^p \right)^{\frac{1}{p}} := d_p(x, y) + d_p(y, z).$$

■

Definition 359 Let T be a non empty set. $\mathcal{B}(T)$ is the set of all bounded real functions defined on T , i.e.,

$$\mathcal{B}(T) := \{f : T \rightarrow \mathbb{R} \quad : \quad \sup \{|f(x)| : x \in T\} < +\infty\},$$

and²

$$d_\infty : \mathcal{B}(T) \times \mathcal{B}(T) \rightarrow \mathbb{R}, \quad d_\infty(f, g) = \sup \{|f(x) - g(x)| : x \in T\}$$

Definition 361

$$l^\infty = \{(x_n)_{n \in \mathbb{N}} \in \mathbb{R}^\infty : \sup \{|x_n| : n \in \mathbb{N}\} < +\infty\}$$

is called the set of bounded real sequences, and, still using the symbol of the previous definition,

$$d_\infty : l^\infty \times l^\infty \rightarrow \mathbb{R}, \quad d_\infty((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) = \sup \{|x_n - y_n| : n \in \mathbb{N}\}$$

Proposition 362 $(\mathcal{B}(T), d_\infty)$ and (l^∞, d_∞) are metric spaces, and d_∞ is called the sup metric.

Proof. We show that $(\mathcal{B}(T), d_\infty)$ is a metric space. As usual, the difficult part is to show property 3 of d_∞ , which is done below.

$$\forall f, g, h \in \mathcal{B}(T), \forall x \in T,$$

$$\begin{aligned} |f(x) - g(x)| &\leq |f(x) - h(x)| + |h(x) - g(x)| \leq \\ &\leq \sup \{|f(x) - h(x)| : x \in T\} + \sup \{|h(x) - g(x)| : x \in T\} = \\ &= d_\infty(f, h) + d_\infty(h, g). \end{aligned}$$

Then, $\forall x \in T$,

$$d_\infty(f, g) := \sup |f(x) - g(x)| \leq d_\infty(f, h) + d_\infty(h, g).$$

■

Exercise 363 If (X, d) is a metric space, then

$$\left(X, \frac{d}{1+d}\right)$$

is a metric space.

Proposition 364 Given a metric space (X, d) and a set Y such that $\emptyset \neq Y \subseteq X$, then $(Y, d|_{Y \times Y})$ is a metric space.

Proof. By definition. ■

Definition 365 Given a metric space (X, d) and a set Y such that $\emptyset \neq Y \subseteq X$, then $(Y, d|_{Y \times Y})$, or simply, (Y, d) is called a metric subspace of X .

Example 366 1. Given \mathbb{R} with the (Euclidean) distance $d_{2,1}$, $([0, 1], d_{2,1})$ is a metric subspace of $(\mathbb{R}, d_{2,1})$.

2. Given \mathbb{R}^2 with the (Euclidean) distance $d_{2,2}$, $(\{0\} \times \mathbb{R}, d_{2,2})$ is a metric subspace of $(\mathbb{R}^2, d_{2,2})$.

Exercise 367 Let $C([0, 1])$ be the set of continuous functions from $[0, 1]$ to \mathbb{R} . Show that a metric on that set is defined by

$$d(f, g) = \int_0^1 |f(x) - g(x)| \, dx,$$

where $f, g \in C([0, 1])$.

Example 368 Let X be the set of continuous functions from \mathbb{R} to \mathbb{R} , and consider $d(f, g) = \sup_{x \in \mathbb{R}} |f(x) - g(x)|$. (X, d) is **not** a metric space because d is not a function from X^2 to \mathbb{R} : it can be $\sup_{x \in \mathbb{R}} |f(x) - g(x)| = +\infty$.

2

Definition 360 Observe that $d_\infty(f, g) \in \mathbb{R}$:

$$d_\infty(f, g) := \sup \{|f(x) - g(x)| : x \in T\} \leq \sup \{|f(x)| : x \in T\} + \sup \{|g(x)| : x \in T\} < +\infty.$$

Example 369 Let $X = \{a, b, c\}$ and $d : X^2 \rightarrow \mathbb{R}$ such that

$$d(a, b) = d(b, a) = 2$$

$$d(a, c) = d(c, a) = 0$$

$$d(b, c) = d(c, b) = 1.$$

Since $d(a, b) = 2 > 0 + 1 = d(a, c) + d(b, c)$, then (X, d) is **not** a metric space.

Example 370 Given $n \in \mathbb{N}, p \in (0, 1)$, $X = \mathbb{R}^2$, define

$$d : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}, \quad (x, y) \mapsto \left(\sum_{i=1}^2 |x_i - y_i|^p \right)^{\frac{1}{p}},$$

(X, d) is **not** a metric space, as shown below. Take $x = (0, 1)$, $y = (1, 0)$ and $z = (0, 0)$. Then

$$d(x, y) = (1^p + 1^p)^{\frac{1}{p}} = 2^{\frac{1}{p}},$$

$$d(x, z) = (0^p + 1^p)^{\frac{1}{p}} = 1$$

$$d(z, y) = 1.$$

Then, $d(x, y) - (d(x, z) + d(z, y)) = 2^{\frac{1}{p}} - 2 > 0$.

9.2 Open and closed sets

Definition 371 Let (X, d) be a metric space. $\forall x_0 \in X$ and $\forall r \in \mathbb{R}_{++}$, the open r -ball of x_0 in (X, d) is the set

$$B_{(X, d)}(x_0, r) = \{x \in X : d(x, x_0) < r\}.$$

If there is no ambiguity about the metric space (X, d) we are considering, we use the lighter notation $B(x_0, r)$ in the place of $B_{(X, d)}(x_0, r)$.

Example 372 1.

$$B_{(\mathbb{R}, d_2)}(x_0, r) = (x_0 - r, x_0 + r)$$

is the open interval of radius r centered in x_0 .

2.

$$B_{(\mathbb{R}^2, d_2)}(x_0, r) = \left\{ (x_1, x_2) \in \mathbb{R}^2 : \sqrt{(x_1 - x_{01})^2 + (x_2 - x_{02})^2} < r \right\}$$

is the open disk of radius r centered in x_0 .

3. In \mathbb{R}^2 with the metric d given by

$$d((x_1, x_2), (y_1, y_2)) = \max\{|x_1 - y_1|, |x_2 - y_2|\}$$

the open ball $B(0, 1)$ can be pictured as done below:

a square around zero.

Definition 373 Let (X, d) be a metric space. x is an interior point of $S \subseteq X$ if

there exists an open ball centered in x and contained in S , i.e.,

$\exists r \in \mathbb{R}_{++}$ such that $B(x, r) \subseteq S$.

Definition 374 The set of all interior points of S is called the Interior of S and it is denoted by $\text{Int}_{(X, d)} S$ or simply by $\text{Int } S$.

Remark 375 *Int $S \subseteq S$, simply because $x \in \text{Int } S \Rightarrow x \in B(x, r) \subseteq S$, where the first inclusion follows from the definition of open ball and the second one from the definition of Interior. In other words, to find interior points of S , we can limit our search to points belonging to S .*

It is not true that $\forall S \subseteq X, S \subseteq \text{Int } S$, as shown below. We want to prove that

$$\neg(\forall S \subseteq X, \forall x \in S, \quad x \in S \Rightarrow x \in \text{Int } S),$$

i.e.,

$$(\exists S \subseteq X \text{ and } x \in S \text{ such that } x \notin \text{Int } S).$$

Take $(X, d) = (\mathbb{R}, d_2)$, $S = \{1\}$ and $x = 1$. Then, clearly $1 \in \{1\}$, but $1 \notin \text{Int } S : \forall r \in \mathbb{R}_{++}, (1 - r, 1 + r) \not\subseteq \{1\}$.

Remark 376 *To understand the following example, recall that $\forall a, b \in \mathbb{R}$ such that $a < b$, $\exists c \in \mathbb{Q}$ and $d \in \mathbb{R} \setminus \mathbb{Q}$ such that $c, d \in (a, b)$ - see, for example, Apostol (1967).*

Example 377 *Let (\mathbb{R}, d_2) be given.*

1. *Int $\mathbb{N} = \text{Int } \mathbb{Q} = \emptyset$.*
2. *$\forall a, b \in \mathbb{R}, a < b$, $\text{Int } [a, b] = \text{Int } [a, b) = \text{Int } (a, b] = \text{Int } (a, b) = (a, b)$.*
3. *Int $\mathbb{R} = \mathbb{R}$.*
4. *Int $\emptyset = \emptyset$.*

Definition 378 *Let (X, d) be a metric space. A set $S \subseteq X$ is open in (X, d) , or (X, d) -open, or open with respect to the metric space (X, d) , if $S \subseteq \text{Int } S$, i.e., $S = \text{Int } S$, i.e.,*

$$\forall x \in S, \exists r \in \mathbb{R}_{++} \text{ such that } B_{(X, d)}(x, r) := \{y \in X : d(y, x) < r\} \subseteq S.$$

Remark 379 *Let (\mathbb{R}, d_2) be given. From Example 377, it follows that*

$\mathbb{N}, \mathbb{Q}, [a, b], [a, b), (a, b]$ are not open sets, and $(a, b), \mathbb{R}$ and \emptyset are open sets. In particular, open interval are open sets, but there are open sets which are not open interval. Take for example $S = (0, 1) \cup (2, 3)$.

Exercise 380 *$\forall n \in \mathbb{N}, \forall i \in \{1, \dots, n\}, \forall a_i, b_i \in \mathbb{R}$ with $a_i < b_i$,*

$$\times_{i=1}^n (a_i, b_i)$$

is (\mathbb{R}^n, d_2) open.

Proposition 381 *Let (X, d) be a metric space. An open ball is an open set.*

Proof. Take $y \in B(x_0, r)$. Define

$$\delta = r - d(x_0, y). \quad (9.8)$$

First of all, observe that, since $y \in B(x_0, r)$, $d(x_0, y) < r$ and then $\delta \in \mathbb{R}_{++}$. It is then enough to show that $B(y, \delta) \subseteq B(x_0, r)$, i.e., we assume that

$$d(z, y) < \delta \quad (9.9)$$

and we want to show that $d(z, x_0) < r$. From the triangle inequality

$$d(z, x_0) \leq d(z, y) + d(y, x_0) \stackrel{(9.9), (9.8)}{<} \delta + (r - \delta) = r,$$

as desired. ■

Example 382 *In a discrete metric space (X, d) , $\forall x \in X, \forall r \in (0, 1]$, $B(x, r) := \{y \in X : d(x, y) < r\} = \{x\}$ and $\forall r > 1$, $B(x, r) := \{y \in X : d(x, y) < r\} = X$. Then, it is easy to show that any subset of a discrete metric space is open, as verified below. Let (X, d) be a discrete metric space and $S \subseteq X$. For any $x \in S$, take $\varepsilon = \frac{1}{2}$; then $B(x, \frac{1}{2}) = \{x\} \subseteq S$.*

Definition 383 Let a metric space (X, d) be given. A set $T \subseteq X$ is (X, d) closed or closed in (X, d) if its complement in X , i.e., $X \setminus T$ is open in (X, d) .

If no ambiguity arises, we simply say that T is closed in X , or even, T is closed; we also write T^C in the place of $X \setminus T$.

Remark 384 S is open $\Leftrightarrow S^C$ is closed, simply because S^C closed $\Leftrightarrow (S^C)^C = S$ is open.

Example 385 The following sets are closed in (\mathbb{R}, d_2) : $\mathbb{R}; \mathbb{N}; \emptyset; \forall a, b \in \mathbb{R}, a < b, \{a\}$ and $[a, b]$.

Remark 386 It is false that:

$$S \text{ is not open} \Rightarrow S \text{ is closed}$$

(and therefore that S is not closed $\Rightarrow S$ is open), i.e., there exist sets which are not open and not closed, for example $(0, 1]$ in (\mathbb{R}, d_2) . There are also two sets which are both open and closed: \emptyset and \mathbb{R}^n in (\mathbb{R}^n, d_2) .

Proposition 387 Let a metric space (X, d) be given.

1. \emptyset and X are open sets.
2. The union of any (finite or infinite) collection of open sets is an open set.
3. The intersection of any finite collection of open sets is an open set.

Proof. 1.

$\forall x \in X, \forall r \in \mathbb{R}_{++}, B(x, r) \subseteq X$. \emptyset is open because it contains no elements.

2.

Let \mathcal{I} be a collection of open sets and $S = \cup_{A \in \mathcal{I}} A$. Assume that $x \in S$. Then there exists $A \in \mathcal{I}$ such that $x \in A$. Then, for some $r \in \mathbb{R}_{++}$

$$x \in B(x, r) \subseteq A \subseteq S$$

where the first inclusion follows from fact that A is open and the second one from the definition of S .

3.

Let \mathcal{F} be a collection of open sets, i.e., $\mathcal{F} = \{A_n\}_{n \in N}$, where $N \subseteq \mathbb{N}$, $\#N$ is finite and $\forall n \in N$, A_n is an open set. Take $S = \cap_{n \in N} A_n$. If $S = \emptyset$, we are done. Assume that $S \neq \emptyset$ and that $x \in S$. Then from the fact that each set A is open and from the definition of S as the intersection of sets

$$\forall n \in N, \exists r_n \in \mathbb{R}_{++} \text{ such that } x \in B(x, r_n) \subseteq A_n$$

Since N is a finite set, there exists a positive $r^* = \min \{r_n : n \in N\} > 0$. Then

$$\forall n \in N, x \in B(x, r^*) \subseteq B(x, r_n) \subseteq A_n$$

and from the very definition of intersections

$$x \in B(x, r^*) \subseteq \cap_{n \in N} B(x, r_n) \subseteq \cap_{n \in N} A_n = S.$$

■

Remark 388 The assumption that $\#N$ is finite cannot be dispensed with:

$$\cap_{n=1}^{+\infty} B\left(0, \frac{1}{n}\right) = \cap_{n=1}^{+\infty} \left(-\frac{1}{n}, \frac{1}{n}\right) = \{0\}$$

is not open.

Remark 389 A generalization of metric spaces is the concept of topological spaces. In fact, we have the following definition which “assumes the previous Proposition”.

Let X be a nonempty set. A collection \mathcal{T} of subsets of X is said to be a topology on X if

1. \emptyset and X belong to \mathcal{T} ,
2. The union of any (finite or infinite) collection of sets in \mathcal{T} belongs to \mathcal{T} ,
3. The intersection of any finite collection of sets in \mathcal{T} belongs to \mathcal{T} .

(X, \mathcal{T}) is called a topological space.

The members of \mathcal{T} are said to be open set with respect to the topology \mathcal{T} , or (X, \mathcal{T}) open.

Proposition 390 *Let a metric space (X, d) be given.*

1. \emptyset and X are closed sets.
2. The intersection of any (finite or infinite) collection of closed sets is a closed set.
3. The union of any finite collection of closed sets is a closed set.

Proof. 1

It follows from the definition of closed set, the fact that $\emptyset^C = X$, $X^C = \emptyset$ and Proposition 387.

2.

Let \mathcal{I} be a collection of closed sets and $S = \cap_{B \in \mathcal{I}} B$. Then, from de Morgan's laws,

$$S^C = (\cap_{B \in \mathcal{I}} B)^C = \cup_{B \in \mathcal{I}} B^C$$

Then from Remark 384, $\forall B \in \mathcal{I}$, B^C is open and from Proposition 387.1, $\cup_{B \in \mathcal{I}} B^C$ is open as well.

2.

Let \mathcal{F} be a collection of closed sets, i.e., $\mathcal{F} = \{B_n\}_{n \in N}$, where $N \subseteq \mathbb{N}$, $\#N$ is finite and $\forall n \in N$, B_n is an open set. Take $S = \cup_{n \in N} B_n$. Then, from de Morgan's laws,

$$S^C = (\cup_{n \in N} B_n)^C = \cap_{n \in N} B_n^C$$

Then from Remark 384, $\forall n \in N$, B_n^C is open and from Proposition 387.2, $\cap_{n \in N} B_n^C$ is open as well.

■

Remark 391 *The assumption that $\#N$ is finite cannot be dispensed with:*

$$\left(\cap_{n=1}^{+\infty} B \left(0, \frac{1}{n} \right) \right)^C = \cup_{n=1}^{+\infty} B \left(0, \frac{1}{n} \right)^C = \cup_{n=1}^{+\infty} \left(\left(-\infty, -\frac{1}{n} \right] \cup \left[\frac{1}{n}, +\infty \right) \right) = \mathbb{R} \setminus \{0\}.$$

is not closed.

Definition 392 *If S is both closed and open in (X, d) , S is called clopen in (X, d) .*

Remark 393 *In any metric space (X, d) , X and \emptyset are clopen.*

Proposition 394 *In any metric space (X, d) , $\{x\}$ is closed.*

Proof. We want to show that $X \setminus \{x\}$ is open. If $X = \{x\}$, then $X \setminus \{x\} = \emptyset$, and we are done. If $X \neq \{x\}$, take $y \in X$, where $y \neq x$. Taken

$$r = d(y, x) \tag{9.10}$$

with $r > 0$, because $x \neq y$. We are left with showing that $B(y, r) \subseteq X \setminus \{x\}$, which is true because of the following argument. Suppose otherwise; then $x \in B(y, r)$, i.e., $r \stackrel{(9.10)}{=} d(y, x) < r$, a contradiction. ■

Remark 395 *From Example 382, any set in any discrete metric space is open. Therefore, the complement of each set is open, and therefore each set is then clopen.*

Definition 396 *Let a metric space (X, d) and a set $S \subseteq X$ be given. x is an boundary point of S if*

any open ball centered in x intersects both S and its complement in X , i.e.,

$$\forall r \in \mathbb{R}_{++}, \quad B(x, r) \cap S \neq \emptyset \quad \wedge \quad B(x, r) \cap S^C \neq \emptyset.$$

Definition 397 *The set of all boundary points of S is called the Boundary of S and it is denoted by $\mathcal{F}(S)$.*

Exercise 398 $\mathcal{F}(S) = \mathcal{F}(S^C)$.

Exercise 399 $\mathcal{F}(S)$ is a closed set.

Definition 400 The closure of S , denoted by $\text{Cl}(S)$ is the intersection of all closed sets containing S , i.e., $\text{Cl}(S) = \bigcap_{S' \in \mathcal{S}} S'$ where $\mathcal{S} := \{S' \subseteq X : S' \text{ is closed and } S \subseteq S'\}$.

Proposition 401 1. $\text{Cl}(S)$ is a closed set;
2. S is closed $\Leftrightarrow S = \text{Cl}(S)$.

Proof. 1.

It follows from the definition and Proposition 390.

2.

$[\Leftarrow]$

It follows from 1. above.

$[\Rightarrow]$

Since S is closed, then $S \in \mathcal{S}$. Therefore, $\text{Cl}(S) = S \cap (\bigcap_{S' \in \mathcal{S}} S') = S$. ■

Definition 402 $x \in X$ is an accumulation point for $S \subseteq X$ if any open ball centered at x contains points of S different from x , i.e., if

$$\forall r \in \mathbb{R}_{++}, (S \setminus \{x\}) \cap B(x, r) \neq \emptyset$$

The set of accumulation points of S is denoted by $D(S)$ and it is called the Derived set of S .

Definition 403 $x \in X$ is an isolated point for $S \subseteq X$ if $x \in S$ and it is not an accumulation point for S , i.e.,

$$x \in S \text{ and } \exists r \in \mathbb{R}_{++} \text{ such that } (S \setminus \{x\}) \cap B(x, r) = \emptyset,$$

or

$$\exists r \in \mathbb{R}_{++} \text{ such that } S \cap B(x, r) = \{x\}.$$

The set of isolated points of S is denoted by $Is(S)$.

Proposition 404 $D(S) = \{x \in \mathbb{R}^n : \forall r \in \mathbb{R}_{++}, S \cap B(x, r) \text{ has an infinite cardinality}\}$.

Proof. $[\subseteq]$

Suppose otherwise, i.e., x is an accumulation point of S and $\exists r \in \mathbb{R}_{++}$ such that $S \cap B(x, r) = \{x_1, \dots, x_n\}$. Then defined $\delta := \min \{d(x, x_i) : i \in \{1, \dots, n\}\}$, $(S \setminus \{x\}) \cap B(x, \frac{\delta}{2}) = \emptyset$, a contradiction.

$[\supseteq]$

Since $S \cap B(x, r)$ has an infinite cardinality, then $(S \setminus \{x\}) \cap B(x, r) \neq \emptyset$. ■

9.2.1 Sets which are open or closed in metric subspaces.

Remark 405 1. $[0, 1)$ is $([0, 1), d_2)$ open.

2. $[0, 1)$ is not (\mathbb{R}, d_2) open. We want to show

$$\neg \langle \forall x_0 \in [0, 1), \exists r \in \mathbb{R}_{++} \text{ such that } B_{(\mathbb{R}, d_2)}(x_0, r) = (x_0 - r, x_0 + r) \subseteq [0, 1) \rangle, \quad (9.11)$$

i.e.,

$$\exists x_0 \in [0, 1) \text{ such that } \forall r \in \mathbb{R}_{++}, \exists x' \in \mathbb{R} \text{ such that } x' \in (x_0 - r, x_0 + r) \text{ and } x' \notin [0, 1).$$

It is enough to take $x_0 = 0$ and $x' = -\frac{r}{2}$.

3. Let $([0, +\infty), d_2)$ be given. $[0, 1)$ is open, as shown below. By definition of open set, - go back to Definition 378 and read it again - we have that, given the metric space $([0, +\infty), d_2)$, $[0, 1)$ is open if

$$\forall x_0 \in [0, 1), \exists r \in \mathbb{R}_{++} \text{ such that } B_{([0, +\infty), d_2)}(x_0, r) := \{x \in [0, +\infty) : d(x_0, x) < r\} = [0, r) \subseteq [0, 1).$$

If $x_0 \in (0, 1)$, then take $r = \min \{x_0, 1 - x_0\} > 0$.

If $x_0 = 0$, then take $r = \frac{1}{2}$. Therefore, we have $B_{(\mathbb{R}_+, d_2)}(0, \frac{1}{2}) = \{x \in \mathbb{R}_+ : |x - 0| < \frac{1}{2}\} = [0, \frac{1}{2}) \subseteq [0, 1)$.

Remark 406 1. $(0, 1)$ is $((0, 1), d_2)$ closed.

2. $(0, 1]$ is $((0, +\infty), d_2)$ closed, simply because $(1, +\infty)$ is open.

Proposition 407 Let a metric space (X, d) , a metric subspace (Y, d) of (X, d) and a set $S \subseteq Y$ be given.

S is open in $(Y, d) \Leftrightarrow$ there exists a set O open in (X, d) such that $S = Y \cap O$.

Proof. Preliminary remark.

$$\forall x_0 \in Y, \forall r \in \mathbb{R}_{++},$$

$$B_{(Y,d)}(x_0, r) := \{x \in Y : d(x_0, x) < r\} = Y \cap \{x \in X : d(x_0, x) < r\} = Y \cap B_{(X,d)}(x_0, r). \quad (9.12)$$

[\Rightarrow]

Taken $x_0 \in S$, by assumption $\exists r_{x_0} \in \mathbb{R}_{++}$ such that $B_{(Y,d)}(x_0, r) \subseteq S \subseteq Y$. Then

$$S = \cup_{x_0 \in S} B_{(Y,d)}(x_0, r) \stackrel{(9.12)}{=} \cup_{x_0 \in S} (Y \cap B_{(X,d)}(x_0, r)) \stackrel{\text{distributive laws}}{=} Y \cap \left(\cup_{x_0 \in S} B_{(X,d)}(x_0, r) \right),$$

and the it is enough to take $O = \cup_{x_0 \in S} B_{(X,d)}(x_0, r)$ to get the desired result.

[\Leftarrow]

Take $x_0 \in S$. then, $x_0 \in O$, and, since, by assumption, O is open in (X, d) , $\exists r \in \mathbb{R}_{++}$ such that $B_{(X,d)}(x_0, r) \subseteq O$. Then

$$B_{(Y,d)}(x_0, r) \stackrel{(9.12)}{=} Y \cap B_{(X,d)}(x_0, r) \subseteq O \cap Y = S,$$

where the last equality follows from the assumption. Summarizing, $\forall x_0 \in S, \exists r \in \mathbb{R}_{++}$ such that $B_{(Y,d)}(x_0, r) \subseteq S$, as desired. ■

Corollary 408 Let a metric space (X, d) , a metric subspace (Y, d) of (X, d) and a set $S \subseteq Y$ be given.

1.

$$\langle S \text{ closed in } (Y, d) \rangle \Leftrightarrow \langle \text{there exists a set } C \text{ closed in } (X, d) \text{ such that } S = Y \cap C. \rangle.$$

2.

$$\langle S \text{ open (respectively, closed) in } (X, d) \rangle \stackrel{\Rightarrow}{\Leftarrow} \langle S \text{ open (respectively, closed) in } (Y, d) \rangle.$$

3. If Y is open (respectively, closed) in X ,

$$\langle S \text{ open (respectively, closed) in } (X, d) \rangle \Leftarrow \langle S \text{ open (respectively, closed) in } (Y, d) \rangle.$$

i.e., “the implication \Leftarrow in the above statement 2. does hold true”.

Proof. 1.

$$\begin{aligned} \langle S \text{ closed in } (Y, d) \rangle &\stackrel{\text{def.}}{\Leftrightarrow} \langle Y \setminus S \text{ open in } (Y, d) \rangle \stackrel{\text{Prop. 407}}{\Leftrightarrow} \\ &\Leftrightarrow \langle \text{there exists an open set } S'' \text{ in } (X, d) \text{ such that } Y \setminus S = S'' \cap Y \rangle \Leftrightarrow \\ &\Leftrightarrow \langle \text{there exists a closed set } S' \text{ in } (X, d) \text{ such that } S = S' \cap Y \rangle, \end{aligned}$$

where the last equivalence is proved below;

[\Leftarrow]

Take $S'' = X \setminus S'$, open in (X, d) by definition. We want to show that if $S'' = X \setminus S$, $S = S' \cap Y$ and $Y \subseteq X$, then $Y \setminus S = S'' \cap Y$:

$$\begin{aligned} x \in Y \setminus S \quad \text{iff} \quad &x \in Y \quad \wedge \quad x \notin S \\ &x \in Y \quad \wedge \quad (x \notin S' \cap Y) \\ &x \in Y \quad \wedge \quad (\neg (x \in S' \cap Y)) \\ &x \in Y \quad \wedge \quad (\neg (x \in S' \wedge x \in Y)) \\ &x \in Y \quad \wedge \quad ((x \notin S' \vee x \notin Y)) \\ &(x \in Y \quad \wedge \quad x \notin S') \quad \vee \quad ((x \in Y \quad \wedge \quad x \notin Y)) \\ &x \in Y \quad \wedge \quad x \notin S' \end{aligned}$$

$$\begin{aligned}
x \in S'' \cap Y \quad \text{iff} \quad & x \in Y \quad \wedge \quad x \in S'' \\
& x \in Y \quad \wedge \quad (x \in X \wedge x \notin S') \\
& (x \in Y \quad \wedge \quad x \in X) \wedge x \notin S' \\
& x \in Y \wedge x \notin S'
\end{aligned}$$

[\Rightarrow]

Take $S' = X \setminus S$. Then S' is closed in (X, d) . We want to show that

if $T' = X \setminus S''$, $Y \setminus S = S'' \cap Y$, $Y \subseteq X$, then $S = S' \cap Y$.

Observe that we want to show that $Y \setminus S = Y \setminus (S' \cap Y)$, or from the assumptions, we want to show that

$$\begin{aligned}
S'' \cap Y &= Y \setminus ((X \setminus S'') \cap Y). \\
x \in Y \setminus ((X \setminus S'') \cap Y) \quad \text{iff} \quad & x \in Y \quad \wedge \quad (\neg(x \in X \setminus S'' \wedge x \in Y)) \\
& x \in Y \quad \wedge \quad (x \notin X \setminus S'' \vee x \notin Y) \\
& x \in Y \quad \wedge \quad (x \in S'' \vee x \notin Y) \\
& (x \in Y \wedge x \in S'') \quad \vee \quad (x \in Y \wedge x \notin Y) \\
& x \in Y \wedge x \in S'' \\
& x \in S'' \cap Y
\end{aligned}$$

2. and 3.

Exercises. ■

9.3 Sequences

Unless otherwise specified, up to the end of the chapter, we assume that

X is a metric space with metric d ,

and

\mathbb{R}^n is the metric space with Euclidean metric.

Definition 409 A sequence in X is a function $x : \mathbb{N} \rightarrow X$.

Usually, for any $n \in \mathbb{N}$, the value $x(n)$ is denoted by x_n , which is called the n -th term of the sequence; the sequence is denoted by $(x_n)_{n \in \mathbb{N}}$.

Definition 410 Given a nonempty set X , X^∞ is the set of sequences $(x_n)_{n \in \mathbb{N}}$ such that $\forall n \in \mathbb{N}$, $x_n \in X$.

Definition 411 A strictly increasing sequence of natural numbers is a sequence $(k_n)_{n \in \mathbb{N}}$ in \mathbb{N} such

$$1 \leq k_1 < k_2 < \dots < k_n < \dots$$

Definition 412 A subsequence of a sequence $(x_n)_{n \in \mathbb{N}}$ is a sequence $(y_n)_{n \in \mathbb{N}}$ such that there exists a strictly increasing sequence $(k_n)_{n \in \mathbb{N}}$ of natural numbers such that $\forall n \in \mathbb{N}$, $y_n = x_{k_n}$.

Definition 413 A sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$ is said to be (X, d) convergent to $x_0 \in X$ (or convergent to $x_0 \in X$ with respect to the metric space (X, d)) if

$$\forall \varepsilon > 0, \exists n_0 \in \mathbb{N} \text{ such that } \forall n > n_0, \quad d(x_n, x_0) < \varepsilon \quad (9.13)$$

x_0 is called the limit of the sequence $(x_n)_{n \in \mathbb{N}}$ and we write

$$\lim_{n \rightarrow +\infty} x_n = x_0, \text{ or } x_n \xrightarrow{n} x_0. \quad (9.14)$$

$(x_n)_{n \in \mathbb{N}}$ in a metric space (X, d) is convergent if there exist $x_0 \in X$ such that (9.13) holds. In that case, we say that the sequence converges to x_0 and x_0 is the limit of the sequence.³

Remark 414 A more precise, and heavy, notation for (9.14) would be

$$\lim_{\substack{n \rightarrow +\infty \\ (X, d)}} x_n = x_0 \quad \text{or} \quad x_n \xrightarrow[(X, d)]{n} x_0$$

³For the last sentence in the Definition, see, for example, Morris (2007), page 121.

Remark 415 Observe that $(\frac{1}{n})_{n \in \mathbb{N}_+}$ converges with respect to (\mathbb{R}, d_2) and it does not converge with respect to (\mathbb{R}_{++}, d_2) .

Proposition 416 $\lim_{n \rightarrow +\infty} x_n = x_0 \Leftrightarrow \lim_{n \rightarrow +\infty} d(x_n, x_0) = 0$.

Proof. Observe that we can define the sequence $(d(x_n, x_0))_{n \in \mathbb{N}}$ in \mathbb{R} . Then from definition 413, we have that $\lim_{n \rightarrow +\infty} d(x_n, x_0) = 0$ means that

$$\forall \varepsilon > 0, \exists n_0 \in \mathbb{N} \text{ such that } \forall n > n_0, |d(x_n, x_0) - 0| < \varepsilon.$$

■

Remark 417 Since $(d(x_n, x_0))_{n \in \mathbb{N}}$ is a sequence in \mathbb{R} , all well known results hold for that sequence. Some of those results are listed below.

Proposition 418 (Some properties of sequences in \mathbb{R}).

All the following statements concern sequences in \mathbb{R} .

1. Every convergent sequence is bounded.
2. Every increasing (decreasing) sequence that is bounded above (below) converges to its sup (inf).
3. Every sequence has a monotone subsequence.
4. (Bolzano-Weierstrass 1) Every bounded sequence has a convergent subsequence.
5. (Bolzano-Weierstrass 2) Every sequence contained in a closed and bounded set has a convergent subsequence in the set.

Moreover, suppose that $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ are sequences in \mathbb{R} and $\lim_{n \rightarrow \infty} x_n = x_0$ and $\lim_{n \rightarrow +\infty} y_n = y_0$. Then

6. $\lim_{n \rightarrow +\infty} (x_n + y_n) = x_0 + y_0$;
7. $\lim_{n \rightarrow +\infty} x_n \cdot y_n = x_0 \cdot y_0$;
8. if $\forall n \in \mathbb{N}, x_n \neq 0$ and $x_0 \neq 0$, $\lim_{n \rightarrow +\infty} \frac{1}{x_n} = \frac{1}{x_0}$;
9. if $\forall n \in \mathbb{N}, x_n \leq y_n$, then $x_0 \leq y_0$;
10. Let $(z_n)_{n \in \mathbb{N}}$ be a sequence such that $\forall n \in \mathbb{N}, x_n \leq z_n \leq y_n$, and assume that $x_0 = y_0$. Then $\lim_{n \rightarrow +\infty} z_n = x_0$.

Proof. See Villanacci, (2015), Basic Facts on sequences and series in \mathbb{R} , mimeo. ■

Proposition 419 If $(x_n)_{n \in \mathbb{N}}$ converges to x_0 and $(y_n)_{n \in \mathbb{N}}$ is a subsequence of $(x_n)_{n \in \mathbb{N}}$, then $(y_n)_{n \in \mathbb{N}}$ converges to x_0 .

Proof. By definition of subsequence, there exists a strictly increasing sequence $(k_n)_{n \in \mathbb{N}}$ of natural numbers, i.e., $1 < k_1 < k_2 < \dots < k_n < \dots$, such that $\forall n \in \mathbb{N}, y_n = x_{k_n}$.

If $n \rightarrow +\infty$, then $k_n \rightarrow +\infty$. Moreover, $\forall n, \exists k_n$ such that

$$d(x_0, x_{k_n}) = d(x_0, y_n)$$

Taking limits of both sides for $n \rightarrow +\infty$, we get the desired result. ■

Proposition 420 A sequence in (X, d) converges at most to one element in X .

Proof. Assume that $x_n \xrightarrow{n} p$ and $x_n \xrightarrow{n} q$; we want to show that $p = q$. From the Triangle inequality,

$$\forall n \in \mathbb{N}, \quad 0 \leq d(p, q) \leq d(p, x_n) + d(x_n, q) \quad (9.15)$$

Since $d(p, x_n) \rightarrow 0$ and $d(x_n, q) \rightarrow 0$, Proposition 418.10 and (9.15) imply that $d(p, q) = 0$ and therefore $p = q$. ■

Proposition 421 Given a sequence $(x_n)_{n \in \mathbb{N}} = \left((x_n^i)_{i=1}^k \right)_{n \in \mathbb{N}}$ in \mathbb{R}^k ,

$$\langle (x_n)_{n \in \mathbb{N}} \text{ converges to } x \rangle \Leftrightarrow \langle \forall i \in \{1, \dots, k\}, (x_n^i)_{n \in \mathbb{N}} \text{ converges to } x^i \rangle,$$

and

$$\lim_{n \rightarrow +\infty} x_n = \left(\lim_{n \rightarrow +\infty} x_n^i \right)_{i=1}^k.$$

Proof. $[\Rightarrow]$

Observe that

$$|x_n^i - x^i| = \sqrt{(x_n^i - x^i)^2} \leq d(x_n x).$$

Then, the result follows.

$[\Leftarrow]$

By assumption, $\forall \varepsilon > 0$ and $\forall i \in \{1, \dots, k\}$, there exists n_0 such that $\forall n > n_0$, we have $|x_n^i - x^i| < \frac{\varepsilon}{\sqrt{k}}$. Then $\forall n > n_0$,

$$d(x_n x) = \left(\sum_{i=1}^k |x_n^i - x^i|^2 \right)^{\frac{1}{2}} < \left(\sum_{i=1}^k \left| \frac{\varepsilon}{\sqrt{k}} \right|^2 \right)^{\frac{1}{2}} = \left(\varepsilon^2 \sum_{i=1}^k \frac{1}{k} \right)^{\frac{1}{2}} = \varepsilon.$$

■

Proposition 422 Suppose that $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ are sequences in \mathbb{R}^k and $\lim_{n \rightarrow \infty} x_n = x_0$ and $\lim_{n \rightarrow +\infty} y_n = y_0$. Then

1. $\lim_{n \rightarrow +\infty} (x_n + y_n) = x_0 + y_0$;
2. $\forall c \in \mathbb{R}, \lim_{n \rightarrow +\infty} c \cdot x_n = c \cdot x_0$;
3. $\lim_{n \rightarrow +\infty} x_n \cdot y_n = x_0 \cdot y_0$.

Proof. It follows from Propositions 418 and 421. ■

Example 423 In Proposition 362, we have seen that $(\mathcal{B}([0, 1]), d_\infty)$ is a metric space. Observe that defined $\forall n \in \mathbb{N}$,


$$f_n : [0, 1] \rightarrow \mathbb{R}, \quad t \mapsto t^n,$$

we have that $(f_n)_n \in \mathcal{B}([0, 1])^\infty$. Moreover, $\forall \bar{t} \in [0, 1]$, $(f_n(\bar{t}))_{n \in \mathbb{N}} \in \mathbb{R}^\infty$ and it converges in (\mathbb{R}, d_2) . In fact,

$$\lim_{n \rightarrow +\infty} \bar{t}^n = \begin{cases} 0 & \text{if } \bar{t} \in [0, 1) \\ 1 & \text{if } \bar{t} = 1. \end{cases}$$

Define

$$f : [0, 1] \rightarrow \mathbb{R}, \quad t \mapsto \begin{cases} 0 & \text{if } t \in [0, 1) \\ 1 & \text{if } t = 1. \end{cases}.$$



We want to check that **it is false that**

$$f_m \xrightarrow{(\mathcal{B}([0,1]), d_\infty)} f,$$

i.e., it is false that $d_\infty(f_m, f) \xrightarrow{m} 0$. Then, we have to check

$$\neg \left\langle \forall \varepsilon > 0 \quad \exists N_\varepsilon \in \mathbb{N} \text{ such that } \forall n > N_\varepsilon, d_\infty(f_n, f) < \varepsilon \right\rangle,$$

i.e.,

$$\exists \varepsilon > 0 \quad \text{such that } \forall N_\varepsilon \in \mathbb{N}, \exists n > N_\varepsilon \text{ such that } d_\infty(f_n, f) \geq \varepsilon.$$

Then, taken $\varepsilon = \frac{1}{4}$, it suffice to show that

$$\langle \forall m \in \mathbb{N}, \exists \bar{t} \in (0, 1) \text{ such that } |f_m(\bar{t}) - f(\bar{t})| \geq \frac{1}{4} \rangle.$$

It is then enough to take $\bar{t} = (\frac{1}{2})^m$.

Exercise 424 For any metric space (X, d) and $(x_n)_{n \in \mathbb{N}} \in X^\infty$,

$$\left\langle x_n \xrightarrow{(X, d)} x \right\rangle \Leftrightarrow \left\langle x_n \xrightarrow{(X, \frac{1}{1+d})} x \right\rangle.$$

9.4 Sequential characterization of closed sets

Proposition 425 Let (X, d) be a metric space and $S \subseteq X$.⁴

$$\langle S \text{ is closed} \rangle \Leftrightarrow \langle \text{any } (X, d) \text{ convergent sequence } (x_n)_{n \in \mathbb{N}} \in S^\infty \text{ converges to an element of } S \rangle.$$

Proof. We want to show that

$$S \text{ is closed} \Leftrightarrow$$

$$\Leftrightarrow \left\langle \left\langle (x_n)_{n \in \mathbb{N}} \text{ is such that} \begin{array}{l} 1. \forall n \in \mathbb{N}, x_n \in S, \quad \text{and} \\ 2. x_n \rightarrow x_0 \end{array} \right\rangle \Rightarrow 3. x_0 \in S \right\rangle.$$

[\Rightarrow]

We are going to show that if S is closed, 2. and not 3. hold, then 1. does not hold. Taken a sequence converging to $x_0 \in X \setminus S$. Since S is closed, $X \setminus S$ is open and therefore $\exists r \in \mathbb{R}_{++}$ such that $B(x_0, r) \subseteq X \setminus S$. Since $x_n \rightarrow x_0$, $\exists M \in \mathbb{N}$ such that $\forall n > M$, $x_n \in B(x_0, r) \subseteq X \setminus S$, contradicting Assumption 1 above.

[\Leftarrow]

Suppose otherwise, i.e., S is not closed. Then, $X \setminus S$ is not open. Then, $\exists \bar{x} \in X \setminus S$ such that $\forall n \in \mathbb{N}$, $\exists x_n \in X$ such that $x_n \in B(\bar{x}, \frac{1}{n}) \cap S$, i.e.,

i. $\bar{x} \in X \setminus S$

ii. $\forall n \in \mathbb{N}$, $x_n \in S$,

iii. $d(x_n, \bar{x}) < \frac{1}{n}$, and therefore $x_n \rightarrow \bar{x}$,

and i., ii. and iii. contradict the assumption. ■

Remark 426 The Appendix to this chapter contains some other characterizations of closed sets and summarizes all the presented characterizations of open and closed sets.

9.5 Compactness

Definition 427 Let (X, d) be a metric space, S a subset of X , and Γ be a set of arbitrary cardinality. A family $\mathcal{S} = \{S_\gamma\}_{\gamma \in \Gamma}$ such that $\forall \gamma \in \Gamma$, S_γ is (X, d) open, is said to be an open cover of S if $S \subseteq \cup_{\gamma \in \Gamma} S_\gamma$.

A subfamily \mathcal{S}' of \mathcal{S} is called a subcover of S if $S \subseteq \cup_{S' \in \mathcal{S}'} S'$.

Definition 428 A metric space (X, d) is compact if every open cover of X has a finite subcover.

A set $S \subseteq X$ is compact in X if every open cover of S has a finite subcover of S .

Example 429 Any finite set in any metric space is compact.

Take $S = \{x_i\}_{i=1}^n$ in (X, d) and an open cover \mathcal{S} of S . For any $i \in \{1, \dots, n\}$, take an open set in \mathcal{S} which contains x_i ; call it S_i . Then $\mathcal{S}' = \{S_i : i \in \{1, \dots, n\}\}$ is the desired open subcover of \mathcal{S} .

⁴Proposition 488 in Appendix 9.8.1 presents a different proof of the result below.

Example 430 1. $(0, 1)$ is not compact in (\mathbb{R}, d_2) .

We want to show that the following statement is true:

$$\neg \langle \forall \mathcal{S} \text{ such that } \cup_{S \in \mathcal{S}} S \supseteq (0, 1), \exists \mathcal{S}' \subseteq \mathcal{S} \text{ such that } \#\mathcal{S}' \text{ is finite and } \cup_{S \in \mathcal{S}'} S \supseteq (0, 1) \rangle,$$

i.e.,

$$\exists \mathcal{S} \text{ such that } \cup_{S \in \mathcal{S}} S \supseteq (0, 1) \text{ and } \forall \mathcal{S}' \subseteq \mathcal{S} \text{ either } \#\mathcal{S}' \text{ is infinite or } \cup_{S \in \mathcal{S}'} S \not\supseteq (0, 1).$$

Take $\mathcal{S} = \left(\left(\frac{1}{n}, 1 \right) \right)_{n \in \mathbb{N} \setminus \{0, 1\}}$ and \mathcal{S}' any finite subcover of \mathcal{S} . Then there exists a finite set N such that $\mathcal{S}' = \left(\left(\frac{1}{n}, 1 \right) \right)_{n \in N}$. Take $n^* = \max \{n \in N\}$. Then, $\cup_{S \in \mathcal{S}'} S = \cup_{n \in N} \left(\frac{1}{n}, 1 \right) = \left(\frac{1}{n^*}, 1 \right)$ and $\left(\frac{1}{n^*}, 1 \right) \not\supseteq (0, 1)$.

2. $(0, 1]$ is not compact in $((0, +\infty), d_2)$. Take $\mathcal{S} = \left(\left(\frac{1}{n}, 2 \right) \right)_{n \in \mathbb{N}}$ and \mathcal{S}' any finite subcover of \mathcal{S} . Then there exists a finite set N such that $\mathcal{S}' = \left(\left(\frac{1}{n}, 2 \right) \right)_{n \in N}$. Take $n^* = \max \{n \in N\}$. Then, $\cup_{S \in \mathcal{S}'} S = \cup_{n \in N} \left(\frac{1}{n}, 2 \right) = \left(\frac{1}{n^*}, 2 \right)$ and $\left(\frac{1}{n^*}, 2 \right) \not\supseteq (0, 1]$.

Proposition 431 Let (X, d) be a metric space.

$$X \text{ compact and } C \subseteq X \text{ closed} \Rightarrow \langle C \text{ compact} \rangle.$$

Proof. Take an open cover \mathcal{S} of C . Then $\mathcal{S} \cup \{X \setminus C\}$ is an open cover of X . Since X is compact, then there exists an open covers \mathcal{S}' of $\mathcal{S} \cup \{X \setminus C\}$ which cover X . Then $\mathcal{S}' \setminus \{X \setminus C\}$ is a finite subcover of \mathcal{S} which covers C . ■

9.5.1 Compactness and bounded, closed sets

Definition 432 Let (X, d) be a metric space and a nonempty subset S of X . S is bounded in (X, d) if $\exists r \in \mathbb{R}_{++}$ such that $\forall x, y \in S$, $d(x, y) < r$.

Proposition 433 Given a metric space (X, d) and a nonempty subset S of X , then

$$S \text{ is bounded} \Leftrightarrow \exists r^* \in \mathbb{R}_{++} \text{ and } \exists \bar{z} \in X \text{ such that } S \subseteq B(\bar{z}, r^*).$$

Proof. $[\Rightarrow]$ Take $r^* = r$ and an arbitrary point \bar{z} in $S \subseteq X$.

$[\Rightarrow]$ Take $x, y \in S$. Then

$$d(x, y) \leq d(x, \bar{z}) + d(\bar{z}, y) < 2r.$$

Then it is enough to take $r^* = 2r$. ■

Proposition 434 The finite union of bounded set is bounded.

Proof. Take $n \in \mathbb{N}$ and $\{S_i\}_{i=1}^n$ such that $\forall i \in \{1, \dots, n\}$, S_i is bounded. Then, $\forall i \in \{1, \dots, n\}$, $\exists r_i \in \mathbb{R}_{++}$ such that $S_i \subseteq B(\bar{z}, r_i)$. Take $r = \max \{r_i\}_{i=1}^n$. Then $\cup_{i=1}^n S_i \subseteq \cup_{i=1}^n B(\bar{z}, r_i) \subseteq B(\bar{z}, r)$. ■

Proposition 435 Let (X, d) be a metric space and S a subset of X .

$$S \text{ compact} \Rightarrow S \text{ bounded}.$$

Proof. If $S = \emptyset$, we are done. Assume then that $S \neq \emptyset$, and take $\bar{x} \in S$ and $\mathcal{B} = \{B(\bar{x}, n)\}_{n \in \mathbb{N}}$. \mathcal{B} is an open cover of X and therefore of S . Then, there exists $\mathcal{B}' \subseteq \mathcal{B}$ such that

$$\mathcal{B}' = \{B(\bar{x}, n_i)\}_{i \in N},$$

where N is a finite set and \mathcal{B}' covers S .

Then takes $n^* = \max_{i \in N} n_i$, we get $S \subseteq B(\bar{x}, n^*)$ as desired. ■

Proposition 436 Let (X, d) be a metric space and S a subset of X .

$$S \text{ compact} \Rightarrow S \text{ closed}.$$

Proof. If $S = X$, we are done by Proposition 390. Assume that $S \neq X$: we want to show that $X \setminus S$ is open. Take $y \in S$ and $x \in X \setminus S$. Then, taken $r_y \in \mathbb{R}$ such that

$$0 < r_y < \frac{1}{2}d(x, y),$$

we have

$$B(y, r_y) \cap B(x, r_y) = \emptyset.$$

Now, $\mathcal{S} = \{B(y, r_y) : y \in S\}$ is an open cover of S , and since S is compact, there exists a finite subcover \mathcal{S}' of \mathcal{S} which covers S , say

$$\mathcal{S}' = \{B(y_n, r_n)\}_{n \in N},$$

such that N is a finite set. Take

$$r^* = \min_{n \in N} r_n,$$

and therefore $r^* > 0$. Then $\forall n \in N$,

$$B(y_n, r_n) \cap B(x, r_n) = \emptyset,$$

$$B(y_n, r_n) \cap B(x, r^*) = \emptyset,$$

and

$$(\cup_{n \in N} B(y_n, r_n)) \cap B(x, r^*) = \emptyset.$$

Since $\{B(y_n, r_n)\}_{n \in N}$ covers S , we then have

$$S \cap B(x, r^*) = \emptyset,$$

or

$$B(x, r^*) \subseteq X \setminus S.$$

Therefore, we have shown that

$$\forall x \in X \setminus S, \exists r^* \in \mathbb{R}_{++} \text{ such that } B(x, r^*) \subseteq X \setminus S,$$

i.e., $X \setminus S$ is open and S is closed. ■

Remark 437 Summarizing, we have seen that in any metric space

$$S \text{ compact} \Rightarrow S \text{ bounded and closed.}$$

The opposite implication is false. In fact, the following sets are bounded, closed and not compact.

1. Let the metric space $((0, +\infty), d_2)$. $(0, 1]$ is closed from Remark 406, it is clearly bounded and it is not compact from Example 430.2.

2. (X, d) where X is an infinite set and d is the discrete metric.

X is closed, from Remark 395.

X is bounded: take $x \in X$ and $r = 2$.

X is not compact. Take $\mathcal{S} = \{B(x, 1)\}_{x \in X}$. Then $\forall x \in X$ there exists a unique element S_x in \mathcal{S} such that $x \in S_x$.⁵

Remark 438 In next section we are going to show that if (X, d) is an Euclidean space with the Euclidean distance and $S \subseteq X$, then

$$S \text{ compact} \Leftrightarrow S \text{ bounded and closed.}$$

⁵For other examples, see among others, page 155, Ok (2007).

9.5.2 Sequential compactness

Definition 439 Let a metric space (X, d) be given. $S \subseteq X$ is sequentially compact if every sequence of elements of S has a subsequence which converges to an element of S , i.e.,

$$\langle (x_n)_{n \in \mathbb{N}} \text{ is a sequence in } S \rangle \Rightarrow \langle \exists \text{ a subsequence } (y_n)_{n \in \mathbb{N}} \text{ of } (x_n)_{n \in \mathbb{N}} \text{ such that } y_n \rightarrow x \in S \rangle.$$

In what follows, we want to prove that in metric spaces, compactness is equivalent to sequential compactness. To do that requires some work and the introduction of some, useful in itself, concepts.

Proposition 440 (Nested intervals) For every $n \in \mathbb{N}$, define $I_n = [a_n, b_n] \subseteq \mathbb{R}$ such that $I_{n+1} \subseteq I_n$. Then $\cap_{n \in \mathbb{N}} I_n \neq \emptyset$.

Proof. By assumption,

$$a_1 \leq a_2 \leq \dots \leq a_n \leq \dots \quad (9.16)$$

and

$$\dots b_n \leq b_{n-1} \leq \dots \leq b_1 \quad (9.17)$$

Then,

$$\forall m, n \in \mathbb{N}, \quad a_m < b_n$$

simply because, if $m > n$, then $a_m < b_m \leq b_n$, where the first inequality follows from the definition of interval I_m and the second one from (9.17), and if $m \leq n$, then $a_m \leq a_n \leq b_n$, where the first inequality follows from (9.16) and the second one from the definition of interval I_n .

Then $A := \{a_n : n \in \mathbb{N}\}$ is nonempty and bounded above by b_n for any n . Then $\sup A := s$ exists. Since $\forall n \in \mathbb{N}$, b_n is an upper bound for A ,

$$\forall n \in \mathbb{N}, \quad s \leq b_n$$

and from the definition of sup,

$$\forall n \in \mathbb{N}, \quad a_n \leq s$$

Then

$$\forall n \in \mathbb{N}, \quad a_n \leq s \leq b_n$$

and

$$\forall n \in \mathbb{N}, \quad I_n \neq \emptyset.$$

■

Remark 441 The statement in the above Proposition is false if instead of taking closed bounded intervals we take either open or unbounded intervals. To see that consider $I_n = (0, \frac{1}{n})$ and $I_n = [n, +\infty]$.

Proposition 442 (Bolzano- Weirstrass) If $S \subseteq \mathbb{R}^n$ has infinite cardinality and is bounded, then S admits at least an accumulation point, i.e., $D(S) \neq \emptyset$.

Proof. Step 1. $n = 1$.

Since S is bounded, $\exists a_0, b_0 \in \mathbb{R}$ such that $S \subseteq [a_0, b_0] := B_0$. Divide B_0 in two subinterval of equal length:

$$\left[a_0, \frac{a_0 + b_0}{2} \right] \text{ and } \left[\frac{a_0 + b_0}{2}, b_0 \right]$$

Choose an interval which contains an infinite number of points in S . Call $B_1 = [a_1, b_1]$ that interval. Proceed as above for B_1 . We therefore obtain a family of intervals

$$B_0 \supseteq B_1 \supseteq \dots \supseteq B_n \supseteq \dots$$

Observe that $\text{lenght } B_0 := b_0 - a_0$ and

$$\forall n \in \mathbb{N}, \text{lenght } B_n = \frac{b_0 - a_0}{2^n}.$$

Therefore, $\forall \varepsilon > 0, \exists N \in \mathbb{N}$ such that $\forall n > N, \text{length } B_n < \varepsilon$.

From Proposition 440, it follows that

$$\exists x \in \bigcap_{n=0}^{+\infty} B_n$$

We are now left with showing that x is an accumulation point for S

$$\forall r \in \mathbb{R}_{++}, B(x, r) \text{ contains an infinite number of points.}$$

By construction, $\forall n \in \mathbb{N}, B_n$ contains an infinite number of points; it is therefore enough to show that

$$\forall r \in \mathbb{R}_{++}, \exists n \in \mathbb{N} \text{ such that } B(x, r) \supseteq B_n.$$

Observe that

$$B(x, r) \supseteq B_n \Leftrightarrow (x - r, x + r) \supseteq [a_n, b_n] \Leftrightarrow x - r < a_n < b_n < x + r \Leftrightarrow \max\{x - a_n, b_n - x\} < r$$

Moreover, since $x \in [a_n, b_n]$,

$$\max\{x - a_n, b_n - x\} < b_n - a_n = \text{length } B_n = \frac{b_0 - a_0}{2^n}$$

Therefore, it suffices to show that

$$\forall r \in \mathbb{R}_{++}, \exists n \in \mathbb{N} \text{ such that } \frac{b_0 - a_0}{2^n} < r$$

i.e., $n \in \mathbb{N}$ and $n > \log_2(b_0 - a_0)$.

Step 2. Omitted (See Ok (2007)). ■

Remark 443 The above Proposition does not say that there exists an accumulation point which belongs to S . To see that, consider $S = \{\frac{1}{n} : n \in \mathbb{N}\}$.

Proposition 444 Let a metric space (X, d) be given and consider the following statements.

1. S is compact set;
2. Every infinite subset of S has an accumulation point which belongs to S , i.e.,

$$\langle T \subseteq S \wedge \#T \text{ is infinite} \rangle \Rightarrow \langle D(T) \cap S \neq \emptyset \rangle,$$

3. S is sequentially compact
4. S is closed and bounded.

Then

$$1. \Leftrightarrow 2. \Leftrightarrow 3 \Rightarrow 4.$$

If $X = \mathbb{R}^n, d = d_2$, then we also have that

$$3 \Leftarrow 4.$$

Proof. (1) \Rightarrow (2)⁶

Take an infinite subset $T \subseteq S$ and suppose otherwise. Then, no point in S is an accumulation point of T , i.e., $\forall x \in S \exists r_x > 0$ such that

$$B(x, r_x) \cap T \setminus \{x\} = \emptyset.$$

Then⁷

$$B(x, r_x) \cap T \subseteq \{x\}. \tag{9.19}$$

⁶Proofs of $1 \Rightarrow 2$ and $2 \Rightarrow 3$ are taken from Aliprantis and Burkinshaw (1990), pages 38-39.

⁷In general,

$$A \setminus B = C \Rightarrow A \subseteq C \cup B, \tag{9.18}$$

as shown below.

Since

$$S \subseteq \cup_{x \in S} B(x, r_x)$$

and S is compact, $\exists x_1, \dots, x_n$ such that

$$S \subseteq \cup_{i=1}^n B(x_i, r_i)$$

Then, since $T \subseteq S$,

$$T = S \cap T \subseteq (\cup_{i=1}^n B(x_i, r_i)) \cap T = \cup_{i=1}^n (B(x_i, r_i) \cap T) \subseteq \{x_1, \dots, x_n\}$$

where the last inclusion follows from (9.19). But then $\#T \leq n$, a contradiction.

(2) \Rightarrow (3)

Take a sequence $(x_n)_{n \in \mathbb{N}}$ of elements in S .

If $\#\{x_n : n \in \mathbb{N}\}$ is finite, then $\exists x_{n^*}$ such that $x_j = x_{n^*}$ for j in an infinite subset of \mathbb{N} , and $(x_{n^*}, \dots, x_{n^*}, \dots)$ is the required convergent subsequence - converging to $x_{n^*} \in S$.

If $\#\{x_n : n \in \mathbb{N}\}$ is infinite, then there exists a subsequence $(y_n)_{n \in \mathbb{N}}$ of $(x_n)_{n \in \mathbb{N}}$ with an infinite amount distinct values, i.e., such that $\forall n, m \in \mathbb{N}, n \neq m$, we have $y_n \neq y_m$. To construct the subsequence $(y_n)_{n \in \mathbb{N}}$, proceed as follows.

$$y_1 = x_1 := x_{k_1},$$

$$y_2 = x_{k_2} \notin \{x_{k_1}\},$$

$$y_3 = x_{k_3} \notin \{x_{k_1}, x_{k_2}\},$$

...

$$y_n = x_{k_n} \notin \{x_{k_1}, x_{k_2}, \dots, x_{k_{n-1}}\},$$

...

Since $T := \{y_n : n \in \mathbb{N}\}$ is an infinite subset of S , by assumption it does have an accumulation point x in S ; moreover, we can redefine $(y_n)_{n \in \mathbb{N}}$ in order to have $\forall n \in \mathbb{N}, y_n \neq x$ ⁸, as follows. If $\exists k$ such that $y_k = x$, take the (sub)sequence $(y_{k+1}, y_{k+2}, \dots) = (y_{k+n})_{n \in \mathbb{N}}$. With some abuse of notation, call still $(y_n)_{n \in \mathbb{N}}$ the sequence so obtained. Now take a further subsequence as follows, using the fact that x is an accumulation point of $\{y_n : n \in \mathbb{N}\} := T$,

$$y_{m_1} \in T \text{ such that } d(y_{m_1}, x) < \frac{1}{1},$$

$$y_{m_2} \in T \text{ such that } d(y_{m_2}, x) < \min \left\{ \frac{1}{2}, (d(y_m, x))_{m \leq m_1} \right\},$$

$$y_{m_3} \in T \text{ such that } d(y_{m_3}, x) < \min \left\{ \frac{1}{3}, (d(y_m, x))_{m \leq m_2} \right\},$$

...

$$y_{m_n} \in T \text{ such that } d(y_{m_n}, x) < \min \left\{ \frac{1}{n}, (d(y_m, x))_{m \leq m_{n-1}} \right\},$$

Observe that since $\forall n, d(y_{m_n}, x) < \min \left\{ (d(y_m, x))_{m \leq m_{n-1}} \right\}$, we have that $\forall n, m_n > m_{n-1}$ and therefore $(y_{m_n})_{n \in \mathbb{N}}$ is a subsequence of $(y_n)_{n \in \mathbb{N}}$ and therefore of $(x_n)_{n \in \mathbb{N}}$. Finally, since

$$\lim_{n \rightarrow +\infty} d(y_{m_n}, x) < \lim_{n \rightarrow +\infty} \frac{1}{n} = 0$$

we also have that

$$\lim_{n \rightarrow +\infty} y_{m_n} = x$$

as desired.

Since $A \setminus B = A \cap B^C$, by assumption, we have

$$(A \cap B^C) \cup B = C \cup B$$

Moreover,

$$(A \cap B^C) \cup B = (A \cup B) \cap (B^C \cup B) = A \cup B \supseteq A$$

Observe that the inclusion in (9.18) can be strict, i.e., it can be

$$A \setminus B = C \wedge A \subset C \cup B;$$

just take $A = \{1\}, B = \{2\}$ and $C = \{1\}$:

$$A \setminus B = \{1\} = C \wedge A = \{1\} \subset C \cup B = \{1, 2\} .;$$

⁸Below we need to have $d(y_n, x) > 0$.

(3) \Rightarrow (1)

It is the content of Proposition 453 below.

(1) \Rightarrow (4)

It is the content of Remark 437.

If $X = \mathbb{R}^n$, (4) \Rightarrow (2)

Take an infinite subset $T \subseteq S$. Since S is bounded T is bounded as well. Then from Bolzano-Weierstrass theorem, i.e., Proposition 442, $D(T) \neq \emptyset$. Since $T \subseteq S$, from Proposition 479, $D(T) \subseteq D(S)$ and since S is closed, $D(S) \subseteq S$. Then, summarizing $\emptyset \neq D(T) \subseteq S$ and therefore $D(T) \cap S = D(T) \neq \emptyset$, as desired. ■

To complete the proof of the above Theorem it suffices to show sequential compactness implies compactness, which is done below, and it requires some preliminary results.

Definition 445 Let (X, d) be a metric space and S a subset of X . S is totally bounded if $\forall \varepsilon > 0, \exists$ a finite set $T \subseteq S$ such that $S \subseteq \cup_{x \in T} B(x, \varepsilon)$.

Proposition 446 Let (X, d) be a metric space and S a subset of X .

$$S \text{ totally bounded} \Rightarrow S \text{ bounded.}$$

Proof. It follows from the definition of totally bounded sets and from Proposition 434. ■

Remark 447 In the previous Proposition, the opposite implication does not hold true.

Example 448 Take (X, d) where X is an infinite set and d is the discrete metric. Then, if $\varepsilon = \frac{1}{2}$, a ball is needed to “take care of each element in X ”. Similar situation arises in the following probably more interesting example.

Example 449 Consider the metric space (l^2, d_2) - see Proposition 358. Recall that

$$l^2 = \left\{ (x_n)_{n \in \mathbb{N}} \in \mathbb{R}^\infty : \sum_{n=1}^{+\infty} |x_n|^2 < +\infty \right\}$$

and

$$d_2 : l^2 \times l^2 \rightarrow \mathbb{R}_+, \quad ((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) \mapsto \left(\sum_{n=1}^{+\infty} |x_n - y_n|^2 \right)^{\frac{1}{2}}.$$

Define $e_m = (e_{m,n})_{n \in \mathbb{N}}$ such that

$$e_{m,n} = \begin{cases} 1 & \text{if } n = m \\ 0 & \text{if } n \neq m, \end{cases}$$

and $S = \{e_m : m \in \mathbb{N}\}$. In other words, $S =$

$$\{(1, 0, 0, \dots, 0, \dots), (0, 1, 0, \dots, 0, \dots), (0, 0, 1, \dots, 0, \dots), \dots\}.$$

Observe that $\forall m \in \mathbb{N}, \sum_{n=1}^{+\infty} |e_{m,n}|^2 = 1$ and therefore $S \subseteq l^2$. We now want to check that S is bounded, but not totally bounded. The main ingredient of the argument below is that

$$\forall m, p \in \mathbb{N} \text{ such that } m \neq p, \quad d(e_m, e_p) = \sqrt{2}. \quad (9.20)$$

1. S is bounded. For any $m \in \mathbb{N}$, $d(e_1, e_m) = \left(\sum_{n=1}^{+\infty} |e_{1,n} - e_{m,n}|^2 \right)^{\frac{1}{2}} = 2^{\frac{1}{2}}$.

2. S is not totally bounded. We want to show that $\exists \varepsilon > 0$ such that for any finite subset T of S there exists $x \in S$ such that $x \notin \cup_{x \in T} B(x, \varepsilon)$. Take $\varepsilon = 1$ and let $T = \{e_k : k \in N\}$ with N arbitrary finite subset of \mathbb{N} . Then, for $k' \in \mathbb{N} \setminus N$, $e_{k'} \in S$ and from (9.20), for any $k' \in \mathbb{N} \setminus N$ and $k \in N$, $d(e_k, e_{k'}) = \sqrt{2} > 1$. Therefore, for $k' \in \mathbb{N} \setminus N$, $e_{k'} \notin \cup_{k \in N} B(e_k, 1)$.

Remark 450 In (\mathbb{R}^n, d_2) , S bounded $\Rightarrow S$ totally bounded.

Lemma 451 *Let (X, d) be a metric space and S a subset of X .*

$$S \text{ sequentially compact} \Rightarrow S \text{ totally bounded.}$$

Proof. Suppose otherwise, i.e., $\exists \varepsilon > 0$ such that for any finite set $T \subseteq S$, $S \not\subseteq \bigcup_{x \in T} B(x, \varepsilon)$. We are now going to construct a sequence in S which does not admit any convergent subsequence, contradicting sequential compactness.

Take an arbitrary

$$x_1 \in S.$$

Then, by assumption $S \not\subseteq B(x_1, \varepsilon)$. Then take $x_2 \in S \setminus B(x_1, \varepsilon)$, i.e.,

$$x_2 \in S \text{ and } d(x_1, x_2) > \varepsilon.$$

By assumption, $S \not\subseteq B(x_1, \varepsilon) \cup B(x_2, \varepsilon)$. Then, take $x_3 \in S \setminus (B(x_1, \varepsilon) \cup B(x_2, \varepsilon))$, i.e.,

$$x_3 \in S \text{ and for } i \in \{1, 2\}, \quad d(x_3, x_i) > \varepsilon.$$

By the axiom of choice, we get that

$$\forall n \in \mathbb{N}, \quad x_n \in S \text{ and for } i \in \{1, \dots, n-1\}, \quad d(x_n, x_i) > \varepsilon.$$

Therefore, we have constructed a sequence $(x_n)_{n \in \mathbb{N}} \in S^\infty$ such that

$$\forall i, j \in \mathbb{N}, \text{ if } i \neq j, \text{ then } d(x_i, x_j) > \varepsilon. \quad (9.21)$$

But, then it is easy to check that $(x_n)_{n \in \mathbb{N}}$ does not have any convergent subsequence in S , as verified below. Suppose otherwise, then $(x_n)_{n \in \mathbb{N}}$ would admit a subsequence $(x_m)_{m \in \mathbb{N}} \in S^\infty$ such that $x_m \rightarrow x \in S$. But, by definition of convergence, $\exists N \in \mathbb{N}$ such that $\forall m > N, d(x_m, x) < \frac{\varepsilon}{2}$, and therefore

$$d(x_m, x_{m+1}) \leq d(x_m, x) + d(x_{m+1}, x) < \varepsilon,$$

contradicting (9.21). ■

Lemma 452 *Let (X, d) be a metric space and S a subset of X .*

$$\left\langle \begin{array}{l} S \text{ sequentially compact} \\ S \text{ is an open cover of } S \end{array} \right\rangle \Rightarrow \left\langle \exists \varepsilon > 0 \text{ such that } \forall x \in S, \exists O_x \in \mathcal{S} \text{ such that } B(x, \varepsilon) \subseteq O_x \right\rangle.$$

Proof. Suppose otherwise; then

$$\forall n \in \mathbb{N}_+, \exists x_n \in S \text{ such that } \forall O \in \mathcal{S}, \quad B\left(x_n, \frac{1}{n}\right) \not\subseteq O. \quad (9.22)$$

By sequential compactness, the sequence $(x_n)_{n \in \mathbb{N}} \in S^\infty$ admits a subsequence, without loss of generality the sequence itself, $(x_n)_{n \in \mathbb{N}} \in S^\infty$ such that $x_n \rightarrow x \in S$. Since \mathcal{S} is an open cover of S , $\exists O \in \mathcal{S}$ such that $x \in O$ and, since O is open, $\exists \varepsilon > 0$ such that

$$B(x, \varepsilon) \subseteq O. \quad (9.23)$$

Since $x_n \rightarrow x$, $\exists M \in \mathbb{N}$ such that $\{x_{M+i}, i \in \mathbb{N}\} \subseteq B(x, \frac{\varepsilon}{2})$. Now, take $n > \max\{M, \frac{2}{\varepsilon}\}$. Then,

$$B\left(x_n, \frac{1}{n}\right) \subseteq B(x, \varepsilon). \quad (9.24)$$

i.e., $d(y, x_n) < \frac{1}{n} \Rightarrow d(y, x) < \varepsilon$, as shown below.

$$d(y, x) \leq d(y, x_n) + d(x_n, x) < \frac{1}{n} + \frac{\varepsilon}{2} < \varepsilon.$$

From (9.23) and (9.24), we get $B(x_n, \frac{1}{n}) \subseteq O \in \mathcal{S}$, contradicting (9.22). ■

Proposition 453 *Let (X, d) be a metric space and S a subset of X .*

$$S \text{ sequentially compact} \Rightarrow S \text{ compact.}$$

Proof. Take an open cover \mathcal{S} of S . Since S is sequentially compact, from Lemma 452,

$$\exists \bar{\varepsilon} > 0 \text{ such that } \forall x \in S \exists O_x \in \mathcal{S} \text{ such that } B(x, \bar{\varepsilon}) \subseteq O_x.$$

Moreover, from Lemma 451 and the definition of total boundedness, there exists a finite set $T \subseteq S$ such that $S \subseteq \cup_{x \in T} B(x, \bar{\varepsilon}) \subseteq \cup_{x \in T} O_x$. But then $\{O_x : x \in T\}$ is the required subcover of \mathcal{S} which covers S . ■

We conclude our discussion on compactness with some results we hope will clarify the concept of compactness in \mathbb{R}^n .

Proposition 454 *Let X be a proper subset of \mathbb{R}^n , and C a subset of X .*

$$\begin{aligned} & C \text{ is bounded and } (\mathbb{R}^n, d_2) \text{ closed} \\ & (\Downarrow 1) \\ & C \text{ is } (\mathbb{R}^n, d_2) \text{ compact} \\ & (\Downarrow 2) \\ & C \text{ is } (X, d_2) \text{ compact} \\ & \Downarrow (\text{not } \Uparrow) 3 \\ & C \text{ is bounded and } (X, d_2) \text{ closed} \end{aligned}$$

Proof. [1 \Downarrow]

It is the content of (Propositions 435, 436 and last part of) Proposition 444.

To show the other result, observe preliminarily that

$$(X \cap S_\alpha) \cup (X \cap S_\beta) = X \cap (S_\alpha \cup S_\beta)$$

[2 \Downarrow]

Take $\mathcal{T} := \{T_\alpha\}_{\alpha \in A}$ such that $\forall \alpha \in A$, T_α is (X, d) open and $C \subseteq \cup_{\alpha \in A} T_\alpha$. From Proposition 407,

$$\forall \alpha \in A, \exists S_\alpha \text{ such that } S_\alpha \text{ is } (\mathbb{R}^n, d_2) \text{ open and } T_\alpha = X \cap S_\alpha.$$

Then

$$C \subseteq \cup_{\alpha \in A} T_\alpha \subseteq \cup_{\alpha \in A} (X \cap S_\alpha) = X \cap (\cup_{\alpha \in A} S_\alpha).$$

We then have that

$$C \subseteq \cup_{\alpha \in A} S_\alpha,$$

i.e., $\mathcal{S} := \{S_\alpha\}_{\alpha \in A}$ is a (\mathbb{R}^n, d_2) open cover of C and since C is (\mathbb{R}^n, d_2) compact, then there exists a finite subcover $\{S_i\}_{i \in N}$ of \mathcal{S} such that

$$C \subseteq \cup_{i \in N} S_i.$$

Since $C \subseteq X$, we then have

$$C \subseteq (\cup_{i \in N} S_i) \cap X = \cup_{i \in N} (S_i \cap X) = \cup_{i \in N} T_i,$$

i.e., $\{T_i\}_{i \in N}$ is a (X, d) open subcover of $\{T_\alpha\}_{\alpha \in A}$ which covers C , as required.

[2 \Uparrow]

Take $\mathcal{S} := \{S_\alpha\}_{\alpha \in A}$ such that $\forall \alpha \in A$, S_α is (\mathbb{R}^n, d_2) open and $C \subseteq \cup_{\alpha \in A} S_\alpha$. From Proposition 407,

$$\forall \alpha \in A, T_\alpha := X \cap S_\alpha \text{ is } (X, d) \text{ open.}$$

Since $C \subseteq X$, we then have

$$C \subseteq (\cup_{\alpha \in A} S_\alpha) \cap X = \cup_{\alpha \in A} (S_\alpha \cap X) = \cup_{\alpha \in A} T_\alpha.$$

Then, by assumption, there exists $\{T_i\}_{i \in N}$ is an open subcover of $\{T_\alpha\}_{\alpha \in A}$ which covers C , and therefore there exists a set N with finite cardinality such that

$$C \subseteq \cup_{i \in N} T_i = \cup_{i \in N} (S_i \cap X) = (\cup_{i \in N} S_i) \cap X \subseteq (\cup_{i \in N} S_i),$$

i.e., $\{S_i\}_{i \in N}$ is a (\mathbb{R}^n, d_2) open subcover of $\{S_\alpha\}_{\alpha \in A}$ which covers C , as required.

[3 \Downarrow]

It is the content of Propositions 435, 436.

[3 not \Uparrow]

See Remark 437.1. ■

Remark 455 The proof of part [2 ⇕] above can be used to show the following result.

Given a metric space (X, d) , a metric subspace (Y, d) a set $C \subseteq Y$, then

$$\begin{array}{c} C \text{ is } (Y, d) \text{ compact} \\ \Downarrow \\ C \text{ is } (X, d) \text{ compact} \end{array}$$

In other words, (X', d) compactness of $C \subseteq X' \subseteq X$ is an intrinsic property of C : it does not depend by the subspace X' you are considering. On the other hand, as we have seen, closedness and openness are **not** an intrinsic property of the set.

Remark 456 Observe also that to define “anyway” compact sets as closed and bounded sets would not be a good choice. The conclusion of the extreme value theorem (see Theorem 525) would not hold in that case. That theorem basically says that a continuous real valued function on a compact set admits a global maximum. It is not the case that a continuous real valued function on a closed and bounded set admits a global maximum: consider the continuous function

$$f : (0, 1] \rightarrow \mathbb{R}, \quad f(x) = \frac{1}{x}.$$

The set $(0, 1]$ is bounded and closed (in $((0, +\infty), d_2)$) and f has no maximum on $(0, 1]$.

9.6 Completeness

9.6.1 Cauchy sequences

Definition 457 Let (X, d) be a metric space. A sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$ is a Cauchy sequence if

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ such that } \forall l, m > N, \quad d(x_l, x_m) < \varepsilon.$$

Proposition 458 Let a metric space (X, d) and a sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$ be given.

1. $(x_n)_{n \in \mathbb{N}}$ is convergent $\Rightarrow (x_n)_{n \in \mathbb{N}}$ is Cauchy, but not vice-versa;
2. $(x_n)_{n \in \mathbb{N}}$ is Cauchy $\Rightarrow (x_n)_{n \in \mathbb{N}}$ is bounded;
3. $(x_n)_{n \in \mathbb{N}}$ is Cauchy and it has a subsequence converging to $x \in X \Rightarrow (x_n)_{n \in \mathbb{N}}$ is convergent to $x \in X$.

Proof. 1.

[\Rightarrow] Since $(x_n)_{n \in \mathbb{N}}$ is convergent, by definition, $\exists x \in X$ such that $x_n \rightarrow x$, $\exists N \in \mathbb{N}$ such that $\forall l, m > N, d(x_l, x_m) < \frac{\varepsilon}{2}$ and $d(x, x_m) < \frac{\varepsilon}{2}$. But then $d(x_l, x_m) \leq d(x_l, x) + d(x, x_m) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$.

[\Leftarrow]

Take $X = (0, 1)$, $d =$ absolute value, $(x_n)_{n \in \mathbb{N}} \in (0, 1)^\infty$ such that $\forall n \in \mathbb{N}, x_n = \frac{1}{n}$. $(x_n)_{n \in \mathbb{N}}$ is Cauchy:

$$\forall \varepsilon > 0, \quad d\left(\frac{1}{l}, \frac{1}{m}\right) = \left|\frac{1}{l} - \frac{1}{m}\right| < \left|\frac{1}{l}\right| + \left|\frac{1}{m}\right| = \frac{1}{l} + \frac{1}{m} < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon,$$

where the last inequality is true if $\frac{1}{l} < \frac{\varepsilon}{2}$ and $\frac{1}{m} < \frac{\varepsilon}{2}$, i.e., if $l > \frac{2}{\varepsilon}$ and $m > \frac{2}{\varepsilon}$. Then, it is enough to take $N > \frac{2}{\varepsilon}$ and $N \in \mathbb{N}$, to get the desired result.

$(x_n)_{n \in \mathbb{N}}$ is not convergent to any point in $(0, 1)$:

take any $\bar{x} \in (0, 1)$. We want to show that

$$\exists \varepsilon > 0 \text{ such that } \forall N \in \mathbb{N} \exists n > N \text{ such that } d(x_n, \bar{x}) > \varepsilon.$$

Take $\varepsilon = \frac{\bar{x}}{2} > 0$ and $\forall N \in \mathbb{N}$, take $n^* \in \mathbb{N}$ such that $\frac{1}{n^*} < \min\left\{\frac{1}{N}, \frac{\bar{x}}{2}\right\}$. Then, $n^* > N$, and

$$\left|\frac{1}{n^*} - \bar{x}\right| = \bar{x} - \frac{1}{n^*} > \bar{x} - \frac{\bar{x}}{2} = \frac{\bar{x}}{2} = \varepsilon.$$

2.

Take $\varepsilon = 1$. Then $\exists N \in \mathbb{N}$ such that $\forall l, m > N$, $d(x_l, x_m) < 1$. If $N = 1$, we are done. If $N \geq 1$, define

$$r = \max \{1, d(x_1, x_N), \dots, d(x_{N-1}, x_N)\}.$$

Then

$$\{x_n : n \in \mathbb{N}\} \subseteq B(x_N, r).$$

3.

Let $(x_{n_k})_{k \in \mathbb{N}}$ be a convergent subsequence to $x \in X$. Then,

$$d(x_n, x) \leq d(x_n, x_{n_k}) + d(x_{n_k}, x).$$

Since $d(x_n, x_{n_k}) \rightarrow 0$, because the sequence is Cauchy, and $d(x_{n_k}, x) \rightarrow 0$, because the subsequence is convergent, the desired result follows. ■

9.6.2 Complete metric spaces

Definition 459 A metric space (X, d) is complete if every Cauchy sequence is a convergent sequence.

Remark 460 If a metric space is complete, to show convergence you do not need to guess the limit of the sequence: it is enough to show that the sequence is Cauchy.

Example 461 $((0, 1), \text{absolute value})$ is not a complete metric space; it is enough to consider $(\frac{1}{n})_{n \in \mathbb{N}}$.

Example 462 Let (X, d) be a discrete metric space. Then, it is complete. Take a Cauchy sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$. Then, we claim that $\exists N \in \mathbb{N}$ and $\bar{x} \in X$ such that $\forall n > N$, $x_n = \bar{x}$. Suppose otherwise:

$$\forall N \in \mathbb{N}, \exists m, m' > N \text{ such that } x_m \neq x_{m'},$$

but then $d(x_m, x_{m'}) = 1$, contradicting the fact that the sequence is Cauchy.

Example 463 (\mathbb{Q}, d_2) is not a complete metric space. Since $\mathbb{R} \setminus \mathbb{Q}$ is dense in \mathbb{R} , $\forall x \in \mathbb{R} \setminus \mathbb{Q}$, we can find $(x_n)_{n \in \mathbb{N}} \in \mathbb{Q}^\infty$ such that $x_n \rightarrow x$.

Proposition 464 (\mathbb{R}^k, d_2) is complete.

Proof. 1. (\mathbb{R}, d_2) is complete.

Take a Cauchy sequence $(x_n)_{n \in \mathbb{N}} \in \mathbb{R}^\infty$. Then, from Proposition 458.2, it is bounded. Then from Bolzano-Weierstrass Theorem (i.e., Proposition 418.4), $(x_n)_{n \in \mathbb{N}}$ does have a convergent subsequence - i.e., $\exists (y_n)_{n \in \mathbb{N}} \in \mathbb{R}^\infty$ which is a subsequence of $(x_n)_{n \in \mathbb{N}}$ and such that $y_n \rightarrow a \in \mathbb{R}$. Then from Proposition 458.3.

2. For any $k \geq 2$, (\mathbb{R}^k, d_2) is complete.

Take a Cauchy sequence $(x_n)_{n \in \mathbb{N}} \in (\mathbb{R}^k)^\infty$. For $i \in \{1, \dots, k\}$, consider $(x_n^i)_{n \in \mathbb{N}} \in \mathbb{R}^\infty$. Then, for any $n, m \in \mathbb{N}$,

$$|x_n^i - x_m^i| < \|x_n - x_m\|.$$

Then, $\forall i \in \{1, \dots, k\}$, $(x_n^i)_{n \in \mathbb{N}} \in \mathbb{R}^\infty$ is Cauchy and therefore from 1. above, $\forall i \in \{1, \dots, k\}$, $(x_n^i)_{n \in \mathbb{N}}$ is convergent. Finally, from Proposition 421, the desired result follows. ■

Example 465 For any nonempty set T , $(\mathcal{B}(T), d_\infty)$ is a complete metric space.

Let $(f_n)_{n \in \mathbb{N}} \in (\mathcal{B}(T))^\infty$ be a Cauchy sequence. For any $\bar{x} \in T$, $(f_n(\bar{x}))_{n \in \mathbb{N}} \in \mathbb{R}^\infty$ is a Cauchy sequence, and since \mathbb{R} is complete, it has a convergent subsequence, without loss of generality, $(f_n(\bar{x}))_{n \in \mathbb{N}}$ itself converging say to $f_{\bar{x}} \in \mathbb{R}$. Define

$$f : T \rightarrow \mathbb{R}, \quad : \bar{x} \mapsto f_{\bar{x}}.$$

We are going to show that (i). $f \in \mathcal{B}(T)$, and (ii) $f_n \rightarrow f$.

(i). Since $(f_n)_n$ is Cauchy,

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ such that } \forall l, m > N, \quad d_\infty(f_l, f_m) := \sup_{x \in T} |f_l(x) - f_m(x)| < \varepsilon.$$

Then,

$$\forall x \in T, \quad |f_l(x) - f_m(x)| \leq \sup_{x \in T} |f_l(x) - f_m(x)| = d_\infty(f_l, f_m) < \varepsilon. \quad (9.25)$$

Taking limits of both sides of (9.25) for $l \rightarrow +\infty$, and using the continuity of the absolute value function, we have that

$$\forall x \in T, \quad \lim_{l \rightarrow +\infty} |f_l(x) - f_m(x)| = |f(x) - f_m(x)| < \varepsilon. \quad (9.26)$$

Since⁹

$$\forall x \in T, \quad ||f(x)| - |f_m(x)|| \leq |f(x) - f_m(x)| < \varepsilon,$$

and therefore,

$$\forall x \in T, |f(x)| \leq |f_m(x)| + \varepsilon.$$

Since $f_l \in \mathcal{B}(T)$, $f \in \mathcal{B}(T)$ as well.

(ii) From (9.26), we also have that

$$\forall x \in T, |f(x) - f_m(x)| < \varepsilon,$$

and by definition of \sup

$$d_\infty(f_m, f) := \sup_{x \in T} |f_m(x) - f(x)| < \varepsilon,$$

i.e., $d_\infty(f_m, f) \rightarrow 0$.

For future use, we also show the following result.

Proposition 466

$$\mathcal{BC}(X) := \{f : X \rightarrow \mathbb{R} : f \text{ is bounded and continuous}\}$$

endowed with the metric $d(f, g) = \sup_{x \in X} |f(x) - g(x)|$ is a complete metric space.

Proof. See Stokey and Lucas (1989), page 47. ■

9.6.3 Completeness and closedness

Proposition 467 Let a metric space (X, d) and a metric subspace (Y, d) of (X, d) be given.

1. Y complete $\Rightarrow Y$ closed;
2. Y complete $\Leftarrow Y$ closed and X complete.

Proof. 1.

Take $(x_n)_{n \in \mathbb{N}} \in Y^\infty$ such that $x_n \rightarrow x$. From Proposition 425, it is enough to show that $x \in Y$. Since $(x_n)_{n \in \mathbb{N}}$ is convergent in X , then it is Cauchy. Since Y is complete, by definition, $x_n \rightarrow x \in Y$.

2.

Take a Cauchy sequence $(x_n)_{n \in \mathbb{N}} \in Y^\infty$. We want to show that $x_n \rightarrow x \in Y$. Since $Y \subseteq X$, $(x_n)_{n \in \mathbb{N}}$ is Cauchy in X , and since X is complete, $x_n \rightarrow x \in X$. But since Y is closed, $x \in Y$. ■

Remark 468 An example of a metric subspace (Y, d) of (X, d) which is closed and not complete is the following one. $(X, d) = (\mathbb{R}_{++}, d_2)$, $(Y, d) = ((0, 1], d_2)$ and $(x_n)_{n \in \mathbb{N}} = (\frac{1}{n})_{n \in \mathbb{N}}$.

Corollary 469 Let a complete metric space (X, d) and a metric subspace (Y, d) of (X, d) be given. Then,

$$Y \text{ complete} \Leftrightarrow Y \text{ closed.}$$

⁹See, for example, page 37 in Ok (2007).

9.7 Fixed point theorem: contractions

Definition 470 Let (X, d) be a metric space. A function $\varphi : X \rightarrow X$ is said to be a contraction if

$$\exists k \in (0, 1) \text{ such that } \forall x, y \in X, \quad d(\varphi(x), \varphi(y)) \leq k \cdot d(x, y).$$

The inf of the set of k satisfying the above condition is called contraction coefficient of φ .

Example 471 1. Given (\mathbb{R}, d_2) ,

$$f_\alpha : \mathbb{R} \rightarrow \mathbb{R}, \quad x \mapsto \alpha x$$

is a contraction iff $|\alpha| < 1$; in that case $|\alpha|$ is the contraction coefficient of f_α .

2. Let S be a nonempty open subset of \mathbb{R} and $f : S \rightarrow S$ a differentiable function. If

$$\sup_{x \in S} |f'(x)| < 1,$$

then f is a contraction.

Definition 472 For any $f, g \in X \subseteq \mathcal{B}(T)$, we say that $f \leq g$ if $\forall x \in T, f(x) \leq g(x)$.

Proposition 473 (Blackwell) Let the following objects be given:

1. a nonempty set T ;
 2. X is a nonempty subset of the set $\mathcal{B}(T)$ such that $\forall f \in X, \forall \alpha \in \mathbb{R}_+, f + \alpha \in X$;
 3. $\phi : X \rightarrow X$ is increasing, i.e., $f \leq g \Rightarrow \phi(f) \leq \phi(g)$;
 4. $\exists \delta \in (0, 1)$ such that $\forall f \in X, \forall \alpha \in \mathbb{R}_+, \phi(f + \alpha) \leq \phi(f) + \delta\alpha$.
- Then ϕ is a contraction with contraction coefficient δ .

Proof. $\forall f, g \in X, \forall x \in T$

$$f(x) - g(x) \leq |f(x) - g(x)| \leq \sup_{x \in T} |f(x) - g(x)| = d_\infty(f, g).$$

Therefore, $f \leq g + d_\infty(f, g)$, and from Assumption 3,

$$\phi(f) \leq \phi(g + d_\infty(f, g)).$$

Then, from Assumption 4,

$$\exists \delta \in (0, 1) \text{ such that } \phi(g + d_\infty(f, g)) \leq \phi(g) + \delta d_\infty(f, g),$$

and therefore

$$\phi(f) \leq \phi(g) + \delta d_\infty(f, g). \quad (9.27)$$

Since the argument above is symmetric with respect to f and g , we also have

$$\phi(g) \leq \phi(f) + \delta d_\infty(f, g). \quad (9.28)$$

From (9.27) and (9.28) and the definition of absolute value, we have

$$|\phi(f) - \phi(g)| \leq \delta d_\infty(f, g),$$

as desired. ■

Proposition 474 (Banach fixed point theorem) Let (X, d) be a complete metric space. If $\phi : X \rightarrow X$ is a contraction with coefficient k , then

$$\exists! x^* \in X \text{ such that } x^* = \phi(x^*). \quad (9.29)$$

and

$$\forall x_0 \in X \text{ and } \forall n \in \mathbb{N}, \quad d(\phi^n(x_0), x^*) \leq k^n \cdot d(x_0, x^*), \quad (9.30)$$

where $\phi^n := (\phi \circ \phi \circ \dots \circ \phi)$.

Proof. (9.29) holds true.

Take any $x_0 \in X$ and define the sequence

$$(x_n)_{n \in \mathbb{N}} \in X^\infty, \text{ with } \forall n \in \mathbb{N}, x_{n+1} = \phi(x_n).$$

We want to show that 1. that $(x_n)_{n \in \mathbb{N}}$ is Cauchy, 2. its limit is a fixed point for ϕ , and 3. that fixed point is unique.

1. First of all observe that

$$\forall n \in \mathbb{N}, d(x_{n+1}, x_n) \leq k^n d(x_1, x_0), \quad (9.31)$$

where k is the contraction coefficient of ϕ , as shown by induction below.

Step 1: $\mathcal{P}(1)$ is true:

$$d(x_2, x_1) = d(\phi(x_1), \phi(x_0)) \leq kd(x_1, x_0)$$

from the definition of the chosen sequence and the assumption that ϕ is a contraction.

Step 2. $\mathcal{P}(n-1) \Rightarrow \mathcal{P}(n)$:

$$d(x_{n+1}, x_n) = d(\phi(x_n), \phi(x_{n-1})) \leq kd(x_n, x_{n-1}) \leq k^n d(x_1, x_0)$$

from the definition of the chosen sequence, the assumption that ϕ is a contraction and the assumption of the induction step.

Now, for any $m, l \in \mathbb{N}$ with $m > l$,

$$\begin{aligned} d(x_m, x_l) &\leq d(x_m, x_{m-1}) + d(x_{m-1}, x_{m-2}) + \dots + d(x_{l+1}, x_l) \leq \\ &\leq (k^{m-1} + k^{m-2} + \dots + k^l) d(x_1, x_0) \leq k^l \frac{1-k^{m-l}}{1-k} d(x_1, x_0), \end{aligned}$$

where the first inequality follows from the triangle inequality, the third one from the following computation¹⁰:

$$k^{m-1} + k^{m-2} + \dots + k^l = k^l (1 + k + \dots + k^{m-l-1}) = k^l \frac{1-k^{m-l}}{1-k}.$$

Finally, since $k \in (0, 1)$, we get

$$d(x_m, x_l) \leq \frac{k^l}{1-k} d(x_1, x_0). \quad (9.32)$$

If $x_1 = x_0$, then for any $m, l \in \mathbb{N}$ with $m > l$, $d(x_m, x_l) = 0$ and $\forall n \in \mathbb{N}$, $x_n = x_0$ and the sequence is converging and therefore it is Cauchy. Therefore, consider the case $x_1 \neq x_0$. From (9.32) it follows that $(x_n)_{n \in \mathbb{N}} \in X^\infty$ is Cauchy: $\forall \varepsilon > 0$ choose $N \in \mathbb{N}$ such that $\frac{k^N}{1-k} d(x_1, x_0) < \varepsilon$, i.e., $k^N < \frac{\varepsilon(1-k)}{d(x_1, x_0)}$ and $N > \frac{\log \frac{\varepsilon(1-k)}{d(x_1, x_0)}}{\log k}$.

2. Since (X, d) is a complete metric space, $(x_n)_{n \in \mathbb{N}} \in X^\infty$ does converge say to $x^* \in X$, and, in fact, we want to show that $\phi(x^*) = x^*$. Then, $\forall \varepsilon > 0$, $\exists N \in \mathbb{N}$ such that $\forall n > N$,

$$\begin{aligned} d(\phi(x^*), x^*) &\leq d(\phi(x^*), x_{n+1}) + d(x_{n+1}, x^*) \leq \\ &\leq d(\phi(x^*), \phi(x_n)) + d(x_{n+1}, x^*) \leq kd(x^*, x_n) + d(x_{n+1}, x^*) \leq \\ &\leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon, \end{aligned}$$

¹⁰We are also using the basic facat used to study geometrical series. Define

$$s_n \equiv 1 + a + a^2 + \dots + a^n;$$

:

Multiply both sides of the above equality by $(1-a)$:

$$(1-a)s_n \equiv (1-a)(1 + a + a^2 + \dots + a^n)$$

$$(1-a)s_n \equiv (1 + a + a^2 + \dots + a^n) - (a + a^2 + \dots + a^{n+1}) = 1 - a^{n+1}$$

Divide both sides by $(1-a)$:

$$s_n \equiv (1 + a + a^2 + \dots + a^n) - (a + a^2 + \dots + a^{n+1}) = \frac{1-a^{n+1}}{1-a} = \frac{1}{1-a} - \frac{a^{n+1}}{1-a}$$

where the first equality comes from the triangle inequality, the second one from the construction of the sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$, the third one from the assumption that ϕ is a contraction and the last one from the fact that $(x_n)_{n \in \mathbb{N}}$ converges to x^* . Since ε is arbitrary, $d(\phi(x^*), x^*) = 0$, as desired.

3. Suppose that \hat{x} is another fixed point for ϕ - beside x^* . Then,

$$d(\hat{x}, x^*) = d(\phi(\hat{x}), \phi(x^*)) \leq kd(\hat{x}, x^*)$$

and assuming $\hat{x} \neq x^*$ would imply $1 \leq k$, a contradiction of the fact that ϕ is a contraction with contraction coefficient k .

(9.30) holds true.

We show the claim by induction on $n \in \mathbb{N}$.

$\mathcal{P}(1)$ is true.

$$d(\phi(x_0), x^*) = d(\phi(x_0), \phi(x^*)) \leq k \cdot d(x_0, x^*),$$

where the equality follows from the fact that x^* is a fixed point for ϕ , and the inequality by the fact that ϕ is a contraction.

$\mathcal{P}(n-1)$ is true implies that $\mathcal{P}(n)$ is true.

$$\begin{aligned} d(\phi^n(x_0), x^*) &= d(\phi^n(x_0), \phi(x^*)) = d(\phi(\phi^{n-1}(x_0)), \phi(x^*)) \leq \\ &\leq k \cdot d(\phi^{n-1}(x_0), x^*) \leq k \cdot k^{n-1} \cdot d(x_0, x^*) = k^n \cdot d(x_0, x^*). \end{aligned}$$

■

9.8 Appendices.

9.8.1 Some characterizations of open and closed sets

Remark 475 From basic set theory, we have $A^C \cap B = \emptyset \Leftrightarrow B \subseteq A$, as verified below.

$$\begin{aligned} \neg \langle \exists x : x \in A^C \wedge x \in B \rangle &= \langle \forall x : x \in A \vee \neg(x \in B) \rangle = \\ &= \langle \forall x : \neg(x \in B) \vee x \in A \rangle \stackrel{(*)}{=} \langle \forall x : x \in B \Rightarrow x \in A \rangle, \end{aligned}$$

where $(*)$ follows from the fact that $\langle p \Rightarrow q \rangle = \langle (\neg p) \vee q \rangle$.

Proposition 476 S is open $\Leftrightarrow S \cap \mathcal{F}(S) = \emptyset$.

Proof. $[\Rightarrow]$

Suppose otherwise, i.e., $\exists x \in S \cap \mathcal{F}(S)$. Since $x \in \mathcal{F}(S)$, $\forall r \in \mathbb{R}_{++}$, $B(x, r) \cap S^C \neq \emptyset$. Then, from Remark 475, $\forall r \in \mathbb{R}_{++}$, it is false that $B(x, r) \subseteq S$, contradicting the assumption that S is open.

$[\Leftarrow]$

Suppose otherwise, i.e., $\exists x \in S$ such that

$$\forall r \in \mathbb{R}_{++}, B(x, r) \cap S^C \neq \emptyset \tag{9.33}$$

Moreover

$$x \in B(x, r) \cap S \neq \emptyset \tag{9.34}$$

But (9.33) and (9.34) imply $x \in \mathcal{F}(S)$. Since $x \in S$, we would have $S \cap \mathcal{F}(S) \neq \emptyset$, contradicting the assumption. ■

Proposition 477 S is closed $\Leftrightarrow \mathcal{F}(S) \subseteq S$.

Proof.

$$S \text{ closed} \Leftrightarrow S^C \text{ open} \stackrel{(1)}{\Leftrightarrow} S^C \cap \mathcal{F}(S^C) = \emptyset \stackrel{(2)}{\Leftrightarrow} S^C \cap \mathcal{F}(S) = \emptyset \stackrel{(3)}{\Leftrightarrow} \mathcal{F}(S) \subseteq S$$

where

(1) follows from Proposition 476;

(2) follows from Remark 398

(3) follows Remark 475. ■

Proposition 478 S is closed $\Leftrightarrow D(S) \subseteq S$.

Proof. We are going to use Proposition 477, i.e., S is closed $\Leftrightarrow \mathcal{F}(S) \subseteq S$.

[\Rightarrow]

Suppose otherwise, i.e.,

$$\exists x \notin S \text{ such that } \forall r \in \mathbb{R}_{++}, (S \setminus \{x\}) \cap B(x, r) \neq \emptyset$$

and since $x \notin S$, it is also true that

$$\forall r \in \mathbb{R}_{++}, S \cap B(x, r) \neq \emptyset \quad (9.35)$$

and

$$\forall r \in \mathbb{R}_{++}, S^C \cap B(x, r) \neq \emptyset \quad (9.36)$$

From (9.35) and (9.36), it follows that $x \in \mathcal{F}(S)$, while $x \notin S$, which from Proposition 477 contradicts the assumption that S is closed.

[\Leftarrow]

Suppose otherwise, i.e., using Proposition 477,

$$\exists x \in \mathcal{F}(S) \text{ such that } x \notin S$$

Then, by definition of $\mathcal{F}(S)$,

$$\forall r \in \mathbb{R}_{++}, B(x, r) \cap S \neq \emptyset.$$

Since $x \notin S$, we also have

$$\forall r \in \mathbb{R}_{++}, B(x, r) \cap (S \setminus \{x\}) \neq \emptyset,$$

i.e., $x \in D(S)$ and $x \notin S$, a contradiction. ■

Proposition 479 $\forall S, T \subseteq X, S \subseteq T \Rightarrow D(S) \subseteq D(T)$.

Proof. Take $x \in D(S)$. Then

$$\forall r \in \mathbb{R}_{++}, (S \setminus \{x\}) \cap B(x, r) \neq \emptyset. \quad (9.37)$$

Since $S \subseteq T$, we also have

$$(T \setminus \{x\}) \cap B(x, r) \supseteq (S \setminus \{x\}) \cap B(x, r). \quad (9.38)$$

From (9.37) and (9.38), we get $x \in D(T)$. ■

Proposition 480 $S \cup D(S)$ is a closed set.

Proof. Take $x \in (S \cup D(S))^C$ i.e., $x \notin S$ and $x \notin D(S)$. We want to show that

$$\exists r \in \mathbb{R}_{++} \text{ such that } B(x, r) \cap (S \cup D(S)) = \emptyset,$$

i.e.,

$$\exists r \in \mathbb{R}_{++} \text{ such that } (B(x, r) \cap S) \cup (B(x, r) \cap D(S)) = \emptyset,$$

Since $x \notin D(S)$, $\exists r \in \mathbb{R}_{++}$ such that $B(x, r) \cap (S \setminus \{x\}) = \emptyset$. Since $x \notin S$, we also have that

$$\exists r \in \mathbb{R}_{++} \text{ such that } S \cap B(x, r) = \emptyset. \quad (9.39)$$

We are then left with showing that $B(x, r) \cap D(S) = \emptyset$. If $y \in B(x, r)$, then from (9.39), $y \notin S$ and $B(x, r) \cap S \setminus \{y\} = \emptyset$, i.e., $y \notin D(S)$, i.e., $B(x, r) \cap D(S) = \emptyset$, as desired. ■

Proposition 481 $\text{Cl}(S) = S \cup D(S)$.

Proof. $[\supseteq]$

Since

$$S \subseteq \text{Cl } (S) \quad (9.40)$$

from Proposition 479,

$$D(S) \subseteq D(\text{Cl } (S)). \quad (9.41)$$

Since $\text{Cl } (S)$ is closed, from Proposition 478,

$$D(\text{Cl } (S)) \subseteq \text{Cl } (S) \quad (9.42)$$

From (9.40), and (9.41), (9.42), we get

$$S \cup D(S) \subseteq \text{Cl } (S)$$

$[\subseteq]$

Since, from Proposition 480, $S \cup D(S)$ is closed and contains S , then by definition of $\text{Cl } (S)$,

$$\text{Cl } (S) \subseteq S \cup D(S).$$

■

To proceed in our analysis, we need the following result.

Lemma 482 For any metric space (X, d) and any $S \subseteq X$, we have that

1. $X = \text{Int } S \cup \mathcal{F}(S) \cup \text{Int } S^C$, and
2. $(\text{Int } S \cup \mathcal{F}(S))^C = \text{Int } S^C$.

Proof. If either $S = \emptyset$ or $S = X$, the results are trivial. Otherwise, observe that either $x \in S$ or $x \in X \setminus S$.

1. If $x \in S$, then

either $\exists r \in \mathbb{R}_{++}$ such that $B(x, r) \subseteq S$ and then $x \in \text{Int } S$,

or $\forall r \in \mathbb{R}_{++}$, $B(x, r) \cap S^C \neq \emptyset$ and then $x \in \mathcal{F}(S)$.

Similarly, if $x \in X \setminus S$, then

either $\exists r' \in \mathbb{R}_{++}$ such that $B(x, r') \subseteq X \setminus S$ and then $x \in \text{Int } (X \setminus S)$,

or $\forall r' \in \mathbb{R}_{++}$, $B(x, r') \cap S \neq \emptyset$ and then $x \in \mathcal{F}(S)$.

2. By definition of Interior and Boundary of a set, $(\text{Int } S \cup \mathcal{F}(S)) \cap \text{Int } S^C = \emptyset$.

Now, for arbitrary sets $A, B \subseteq X$ such that $A \cup B = X$ and $A \cap B = \emptyset$, we have what follow:

$A \cup B = X \Leftrightarrow (A \cup B)^C = X^C \Leftrightarrow A^C \cap B^C = \emptyset$, and from Remark 475, $B^C \subseteq A$;

$A \cap B = \emptyset \Leftrightarrow A \cap (B^C)^C = \emptyset \Rightarrow A \subseteq B^C$.

Therefore we can the desired result. ■

Proposition 483 $\text{Cl } (S) = \text{Int } S \cup \mathcal{F}(S)$.

Proof. From Lemma 482, it is enough to show that

$$(\text{Cl } (S))^C = \text{Int } S^C.$$

$[\supseteq]$

Take $x \in \text{Int } S^C$. Then, $\exists r \in \mathbb{R}_{++}$ such that $B(x, r) \subseteq S^C$ and therefore $B(x, r) \cap S = \emptyset$ and, since $x \notin S$,

$$B(x, r) \cap (S \setminus \{x\}) = \emptyset.$$

Then $x \notin S$ and $x \notin D(S)$, i.e.,

$$x \notin S \cup D(S) = \text{Cl } (S)$$

where last equality follows from Proposition 481. In other words, $x \in (\text{Cl } (S))^C$.

$[\subseteq]$

Take $x \in (\text{Cl } (S))^C = (D(S) \cup S)^C$. Since $x \notin D(S)$,

$$\exists r \in \mathbb{R}_{++} \text{ such that } (S \setminus \{x\}) \cap B(x, r) = \emptyset \quad (9.43)$$

Since $x \notin S$,

$$\exists r \in \mathbb{R}_{++} \text{ such that } S \cap B(x, r) = \emptyset \quad (9.44)$$

i.e.,

$$\exists r \in \mathbb{R}_{++} \text{ such that } B(x, r) \subseteq S^C \quad (9.45)$$

and $x \in \text{Int } S^C$. ■

Definition 484 $x \in X$ is an adherent point for S if $\forall r \in \mathbb{R}_{++}, B(x, r) \cap S \neq \emptyset$ and

$$\text{Ad}(S) := \{x \in X : \forall r \in \mathbb{R}_{++}, B(x, r) \cap S \neq \emptyset\}$$

Corollary 485 1. $\text{Cl}(S) = \text{Ad}(S)$.

2. A set S is closed $\Leftrightarrow \text{Ad}(S) = S$.

Proof. 1.

[\subseteq]

$x \in \text{Cl}(S) \Rightarrow \langle x \in \text{Int} S \text{ or } \mathcal{F}(S) \rangle$ and in both cases the desired conclusion is insured.

[\supseteq]

If $x \in S$, then, by definition of closure, $x \in \text{Cl}(S)$. If $x \notin S$, then $S = S \setminus \{x\}$ and, from the assumption, $\forall r \in \mathbb{R}_{++}, B(x, r) \cap (S \setminus \{x\}) \neq \emptyset$, i.e., $x \in D(S)$ which is contained in $\text{Cl}(S)$ from Proposition 481.

2. It follows from 1. above and Proposition 401.2. ■

Proposition 486 Let $S \subseteq \mathbb{R}^n$ be given. Then, $\mathcal{F}(S) = \text{Cl}(S) \setminus \text{Int}(S)$.

Proof. We want to show

$$\langle \forall r \in \mathbb{R}_{++}, B(x, r) \cap S \neq \emptyset \text{ and } B(x, r) \cap S^c \neq \emptyset \rangle \Leftrightarrow \langle x \in \text{Cl}(S) \setminus \text{Int} S \rangle.$$

From Definition ?? and ??, $x \in \text{Cl}(S) \setminus \text{Int}(S)$ iff $\forall r \in \mathbb{R}_{++}, B(x, r) \cap S \neq \emptyset$ and $\neg(\exists \delta > 0 \text{ such that } B(x, \delta) \subseteq S)$, i.e., $\forall \delta > 0$,

$$B(x, \delta) \cap (\mathbb{R}^n \setminus S) = B(x, \delta) \cap \mathbb{R}^n \cap S^C = B(x, \delta) \cap S^C \neq \emptyset.$$

Therefore, $x \in \text{Cl}(S) \setminus \text{Int}(S)$ iff $\forall r \in \mathbb{R}_{++}, B(x, r) \cap S \neq \emptyset$ and $B(x, \delta) \cap S^C \neq \emptyset$. ■

Proposition 487 $x \in \text{Cl}(S) \Leftrightarrow \exists (x_n)_{n \in \mathbb{N}}$ in S converging to x .

Proof. [\Rightarrow]

From Corollary 485, if $x \in \text{Cl}(S)$ then $\forall n \in \mathbb{N}$, we can take $x_n \in B(x, \frac{1}{n}) \cap S$. Then $d(x, x_n) < \frac{1}{n}$ and $\lim_{n \rightarrow +\infty} d(x, x_n) = 0$.

[\Leftarrow]

By definition of convergence,

$$\forall \varepsilon > 0, \exists n_\varepsilon \in \mathbb{N} \text{ such that } \forall n > n_\varepsilon, d(x_n, x) < \varepsilon \text{ or } x_n \in B(x, \varepsilon)$$

or

$$\forall \varepsilon > 0, B(x, \varepsilon) \cap S \supseteq \{x_n : n > n_\varepsilon\}$$

and

$$\forall \varepsilon > 0, B(x, \varepsilon) \cap S \neq \emptyset$$

i.e., $x \in \text{Ad}(S)$, and from the Corollary 485.1, the desired result follows. ■

Proposition 488 S is closed \Leftrightarrow any convergent sequence $(x_n)_{n \in \mathbb{N}}$ with elements in S converges to an element of S .

Proof. We are going to show that S is closed using Proposition 478, i.e., S is closed $\Leftrightarrow D(S) \subseteq S$. We want to show that

$$\langle D(S) \subseteq S \rangle \Leftrightarrow \left\langle \left\langle (x_n)_{n \in \mathbb{N}} \text{ is such that } \begin{array}{l} 1. \forall n \in \mathbb{N}, x_n \in S, \quad \text{and} \\ 2. x_n \rightarrow x_0 \end{array} \right\rangle \Rightarrow x_0 \in S \right\rangle,$$

[\Rightarrow]

Suppose otherwise, i.e., there exists $(x_n)_{n \in \mathbb{N}}$ such that 1. $\forall n \in \mathbb{N}, x_n \in S$. and 2. $x_n \rightarrow x_0$, but $x_0 \notin S$.

By definition of convergent sequence, we have

$$\forall \varepsilon > 0, \exists n_0 \in \mathbb{N} \text{ such that } \forall n > n_0, \quad d(x_n, x_0) < \varepsilon$$

and, since $\forall n \in \mathbb{N}, x_n \in S$,

$$\{x_n : n > n_0\} \subseteq B(x_0, \varepsilon) \cap (S \setminus \{x_0\})$$

Then,

$$\forall \varepsilon > 0, B(x_0, \varepsilon) \cap (S \setminus \{x_0\}) \neq \emptyset$$

and therefore $x_0 \in D(S)$ while $x_0 \notin S$, contradicting the fact that S is closed.

[\Leftarrow]

Suppose otherwise, i.e., $\exists x_0 \in D(S)$ and $x_0 \notin S$. We are going to construct a convergent sequence $(x_n)_{n \in \mathbb{N}}$ with elements in S which converges to x_0 (a point not belonging to S).

From the definition of accumulation point,

$$\forall n \in \mathbb{N}, \quad (S \setminus \{x_0\}) \cap B\left(x, \frac{1}{n}\right) \neq \emptyset.$$

Then, we can take $x_n \in (S \setminus \{x_0\}) \cap B\left(x, \frac{1}{n}\right)$, and since $d(x_n, x_0) < \frac{1}{n}$, we have that $d(x_n, x_0) \rightarrow 0$. ■

Summarizing, the following statements are equivalent:

1. S is open (i.e., $S \subseteq \text{Int } S$)
2. S^C is closed,
3. $S \cap \mathcal{F}(S) = \emptyset$,

and the following statements are equivalent:

1. S is closed,
2. S^C is open,
3. $\mathcal{F}(S) \subseteq S$,
4. $S = \text{Cl}(S)$.
5. $D(S) \subseteq S$,
6. $\text{Ad}(S) = S$,
7. any convergent sequence $(x_n)_{n \in \mathbb{N}}$ with elements in S converges to an element of S .

9.8.2 Norms and metrics

In these notes, a field K can be either \mathbb{R} or \mathbb{C} .

Definition 489 A norm on a vector space E on a field $K \in \{\mathbb{R}, \mathbb{C}\}$ is a function

$$\|\cdot\| : E \rightarrow \mathbb{R}, \quad x \mapsto \|x\|$$

which satisfies the following properties: $\forall x, y \in E, \forall \lambda \in K$,

1. $\|x\| \geq 0$ (non negativity),
2. $\|x + y\| \leq \|x\| + \|y\|$ (triangle inequality),
3. $\|\lambda x\| = |\lambda| \cdot \|x\|$ (homogeneity),¹¹

¹¹If $\lambda \in \mathbb{R}$, $|\lambda|$ is the absolute value of λ ; if $\lambda = (a, b) \in \mathbb{C}$, then $|\lambda| = \sqrt{a^2 + b^2}$.

4. $\|x\| = 0 \Rightarrow x = 0$ (separation).

Proposition 490 Given a norm $\|\cdot\|$ on E , $\forall x, y \in E$

1. $x = 0 \Rightarrow \|x\| = 0$
2. $\|x - y\| = \|y - x\|$
3. $||\|x\| - \|y\|| \leq \|x - y\|$.

Proof.

1. Since E is a vector space, $\forall x \in E$, $0x = 0$ (from Proposition 135 in Villanacci (2012)). Then

$$\|0\| = \|0x\| \stackrel{(a)}{=} |0| \cdot \|x\| \stackrel{(b)}{=} 0\|x\| = 0$$

where (a) follows from property 3 of norm and (b) from the definition of absolute value.

2. $\|x - y\| = \|-y + x\| \stackrel{12}{=} \|-y - (-x)\| \stackrel{(c)}{=} \|(-1)y + (-1)(-x)\| = \|(-1)(y - x)\| =$
 $\|-1\| \cdot \|y - x\| = \|y - x\|$

where (c) follows from Proposition 135.4 in Villanacci (2012).

3. From the definition of absolute value, we want to show that

$$-\|x - y\| \leq \|x\| - \|y\| \leq \|x - y\|.$$

Indeed,

$$\|x\| = \|x - y + y\| \leq \|x - y\| + \|y\|,$$

i.e., $\|x - y\| \geq \|x\| - \|y\|$, and

$$\|y\| = \|y - x + x\| \leq \|y - x\| + \|x\| = \|x - y\| + \|x\|,$$

i.e., $-\|x - y\| \leq \|x\| - \|y\|$, as desired.

■

Proposition 491 If properties 2. and 3. in Definition 489 hold true, then property 1. in the same definition holds true.

Proof. We want to show that

$$\begin{aligned} \langle \|x + y\| \leq \|x\| + \|y\| \wedge \|\lambda x\| = |\lambda| \cdot \|x\| \rangle \\ \Rightarrow \langle \forall x \in E, \|x\| \geq 0 \rangle \end{aligned}$$

Observe that if $x = 0$, from Proposition 490.2 (which uses only property 3 of Definition 489) we have $\|x\| = 0$. Then

$$\begin{aligned} \|0\| &\leq \|x - x\| \leq \|x\| + \|-x\| \text{ and therefore} \\ \|x\| &\geq -\|x\| = -\|-1\| \cdot \|x\| = -\|x\|. \end{aligned}$$

Now, if $\|x\| < 0$, we would have a negative number strictly larger than a positive number, which is a contradiction. ■

Definition 492 The pair $(E, \|\cdot\|)$, where E is a vector space and $\|\cdot\|$ is a norm, is called a normed vector space.

Remark 493 Normed spaces are, by definition, vector space.

Definition 494 A seminorm is a function satisfying properties 1, 2 and 3 in Definition 489.

¹² $\forall v \in V \quad (-1)v = -v \text{ and } -(-v) = (-1)((-1)v) = ((-1)(-1))v = 1 \cdot v = v.$

Definition 495 Given a non-empty set X , a function $d : X \times X \rightarrow \mathbb{R}$ is called a metric or a distance on X if $\forall x, y, z \in X$,

1. $d(x, y) \geq 0$ (non negativity),
 2. $d(x, y) = 0 \Leftrightarrow x = y$ (coincidence),
 3. $d(x, y) = d(y, x)$ (symmetry),
 4. $d(x, z) \leq d(x, y) + d(y, z)$ (triangle inequality),
- (X, d) is called a metric space.

Definition 496 Given a normed vector space $(E, \|\cdot\|)$, the metric

$$d : E^2 \rightarrow \mathbb{R}, (x, y) \mapsto \|x - y\|$$

is called the metric induced by the norm $\|\cdot\|$.

Proposition 497 Given a normed vector space $(E, \|\cdot\|)$,

$$d : E \times E \rightarrow \mathbb{R}, (x, y) \mapsto \|x - y\|$$

is a metric and (E, d) is a metric space.

Proof.

1. It follows from the fact that $x, y \in E \Rightarrow x - y \in E$ and property 1 of the norm.
2. It follows from property 1 of the norm and Proposition 490.1.
3. It follows from Proposition 490.2.
4. $d(x, z) = \|x - z\| = \|(x - y) + (y - z)\| \leq \|x - y\| + \|y - z\| = d(x, y) + d(y, z)$.

■

Proposition 498 If $\|\cdot\|$ is a norm on a vector space E and

$$d : E \times E \rightarrow \mathbb{R}, (x, y) \mapsto \|x - y\|$$

then $\forall x, y, z \in E, \forall \lambda \in K$

- a. $d(x, 0) = \|x\|$
- b. $d(x + z, y + z) = d(x, y)$ (translation invariance)
- c. $d(\lambda x, \lambda y) = |\lambda|d(x, y)$ (homogeneity).

Proof.

- a. $d(x, 0) = \|x - 0\| = \|x\|$
- b. $d(x + z, y + z) = \|(x + z) - (y + z)\| = \|x - y\| = d(x, y)$
- c. $d(\lambda x, \lambda y) = \|\lambda x - \lambda y\| = \|\lambda(x - y)\| = |\lambda| \cdot \|x - y\| = |\lambda|d(x, y)$.

■

Proposition 499 Let (E, d) be a metric space such that d satisfies translation invariance and homogeneity. Then

$$n : E \rightarrow \mathbb{R}, x \mapsto d(x, 0)$$

is a norm and $\forall x, y \in E, n(y - x) = d(x, y)$.

Proof.

1. $n(x) = d(x, 0) \geq 0$, where the inequality follows from property 1 in Definition 4,

2.

$$\begin{aligned} n(x+y) &= d(x+y, 0) \stackrel{(a)}{=} d(x+y-y, 0-y) = d(x, -y) \stackrel{(b)}{\leq} d(x, 0) + d(0, -y) = \\ &\stackrel{(c)}{=} d(x, 0) + d(-y, 0) \stackrel{(d)}{=} d(0, y) + d(0, x) = n(y) + n(x), \end{aligned}$$

where (a) follows from translation invariance, (b) from triangle inequality in Definition 495, (c) from symmetry in Definition 495 and (d) from homogeneity.

3.

$$n(\lambda x) = d(\lambda x, 0) = |\lambda|d(x, 0) = |\lambda|n(x),$$

4.

$$n(x) = 0 \Rightarrow d(x, 0) = 0 \Rightarrow x = 0.$$

It follows that

$$n(y-x) = d(y-x, 0) = d(y-x+x, 0+x) = d(y, x) = d(x, y).$$

■

Remark 500 The above Proposition suggests that the following statement is **false**:

Given a metric space (E, d) , then $n_d : E \rightarrow \mathbb{R}, : x \mapsto d(x, 0)$ is a norm on E .

The fact that the above statement is false is verified below. Take an arbitrary vector space E with the discrete metric d ,

$$d : E \times E \rightarrow \mathbb{R}, \quad d(x, y) = \begin{cases} 0 & \text{if } x = y \\ 1 & \text{if } x \neq y \end{cases}$$

First of all, let's verify that d does not satisfy (translation invariance and homogeneity), otherwise from Proposition 499, we would contradict the desired result. Indeed homogeneity fails.

Take $x \neq y$ and $\lambda = 2$ then

$$d(x, y) = 1 \quad \text{and} \quad d(\lambda x, \lambda y) = 1$$

$$|\lambda|d(x, y) = 2 \neq 1.$$

Let's now show that in the case of the discrete metric

$$n : E \rightarrow \mathbb{R}, \quad x \mapsto d(x, 0)$$

is not a norm. Take $x \neq 0$ and $\lambda = 2$ then

$$||\lambda x|| = d(\lambda x, 0) = 1$$

$$|\lambda|d(x, 0) = 2.$$

9.9 Exercises

Problem sets: 3,4,5,6.

From Lipschutz (1965), starting from page 54: 1, 18, 19, 20, 23, 28 (observe that Lipschutz uses the word “range” in the place of “image”);

starting from page 120: 1, 3, 6, 7, 25, 29.

Chapter 10

Functions

10.1 Limits of functions

In what follows we take for given metric spaces (X, d) and (X', d') and sets $S \subseteq X$ and $T \subseteq X'$.

Definition 501 Given $x_0 \in D(S)$, i.e., given an accumulation point x_0 for S , and $f : S \rightarrow T$, we write

$$\lim_{x \rightarrow x_0} f(x) = l$$

if

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } x \in (B_{(X,d)}(x_0, \delta) \cap S) \setminus \{x_0\} \Rightarrow f(x) \in B_{(X',d')}(l, \varepsilon)$$

or

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } (x \in S \wedge 0 < d(x, x_0) < \delta) \Rightarrow d'(f(x), l) < \varepsilon$$

Proposition 502 Given $x_0 \in D(S)$ and $f : S \rightarrow T$,

$$\begin{aligned} & \langle \lim_{x \rightarrow x_0} f(x) = l \rangle \\ & \Leftrightarrow \\ & \left\langle \begin{array}{l} \text{for any sequence } (x_n)_{n \in \mathbb{N}} \text{ in } S \text{ such that } \forall n \in \mathbb{N}, x_n \neq x_0 \text{ and } \lim_{n \rightarrow +\infty} x_n = x_0, \\ \lim_{n \rightarrow +\infty} f(x_n) = l. \end{array} \right\rangle \end{aligned}$$

Proof. for the following proof see also Proposition 6.2.4, page 123 in Morris.

[\Rightarrow]

Take

a sequence $(x_n)_{n \in \mathbb{N}}$ in S such that $\forall n \in \mathbb{N}, x_n \neq x_0$ and $\lim_{n \rightarrow +\infty} x_n = x_0$.

We want to show that $\lim_{n \rightarrow +\infty} f(x_n) = l$, i.e.,

$$\forall \varepsilon > 0, \exists n_0 \in \mathbb{N} \text{ such that } \forall n > n_0, d(f(x_n), l) < \varepsilon.$$

Since $\lim_{x \rightarrow x_0} f(x) = l$,

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } x \in S \wedge 0 < d(x, x_0) < \delta \Rightarrow d(f(x), l) < \varepsilon.$$

Since $\lim_{n \rightarrow +\infty} x_n = x_0$

$$\forall \delta > 0, \exists n_0 \in \mathbb{N} \text{ such that } \forall n > n_0, 0 <^{(*)} d(x_n, x_0) < \delta,$$

where $(*)$ follows from the fact that $\forall n \in \mathbb{N}, x_n \neq x_0$.

Therefore, combining the above results, we get

$$\forall \varepsilon > 0, \exists n_0 \in \mathbb{N} \text{ such that } \forall n > n_0, d(f(x_n), l) < \varepsilon$$

as desired.

[\Leftarrow]

Suppose otherwise, then

$$\begin{aligned} \exists \varepsilon > 0 \text{ such that } \forall \delta_n = \frac{1}{n}, \text{ i.e., } \forall n \in \mathbb{N}, \exists x_n \in S \text{ such that} \\ x_n \in S \wedge 0 < d(x_n, x_0) < \frac{1}{n} \text{ and } d(f(x), l) \geq \varepsilon. \end{aligned} \quad (10.1)$$

Consider $(x_n)_{n \in \mathbb{N}}$; then, from the above and from Proposition 416, $x_n \rightarrow x_0$, and from the above (specifically the fact that $0 < d(x_n, x_0)$), we also have that $\forall n \in \mathbb{N}, x_n \neq x_0$. Then by assumption, $\lim_{n \rightarrow +\infty} f(x_n) = l$, i.e., by definition of limit,

$$\forall \varepsilon > 0, \exists N \in \mathbb{N} \text{ such that if } n > N, \text{ then } |f(x_n) - l| < \varepsilon,$$

contradicting (10.1). ■

Proposition 503 (*uniqueness*) Given $x_0 \in D(S)$ and $f : S \rightarrow T$,

$$\left\langle \lim_{x \rightarrow x_0} f(x) = l_1 \text{ and } \lim_{x \rightarrow x_0} f(x) = l_2 \right\rangle \Rightarrow \langle l_1 = l_2 \rangle$$

Proof. It follows from Proposition 502 and Proposition 420. ■

Proposition 504 Given $S \subseteq X, x_0 \in D(S)$ and $f, g : S \rightarrow \mathbb{R}$, and

$$\lim_{x \rightarrow x_0} f(x) = l \text{ and } \lim_{x \rightarrow x_0} g(x) = m$$

1. $\lim_{x \rightarrow x_0} f(x) + g(x) = l + m$;
2. $\lim_{x \rightarrow x_0} f(x) \cdot g(x) = l \cdot m$;
3. if $m \neq 0$ and $\forall x \in S, g(x) \neq 0$, $\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = \frac{l}{m}$.

Proof. It follows from Proposition 502 and Proposition 418. ■

10.2 Continuous Functions

Definition 505 Given a metric space (X, d) and a set $V \subseteq X$, an open neighborhood of V is an open set containing V .

Remark 506 Sometimes, an open neighborhood is simply called a neighborhood.

Definition 507 Take $S \subseteq (X, d), T \subseteq (Y, d'), x_0 \in S$ and $f : S \rightarrow T$. Then, f is $(X, d) - (Y, d')$ continuous at x_0 if

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } x \in (B_{(X, d)}(x_0, \delta) \cap S) \Rightarrow f(x) \in B_{(Y, d')}(f(x_0), \varepsilon),$$

i.e.,

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } x \in S \wedge d(x, x_0) < \delta \Rightarrow d'(f(x), f(x_0)) < \varepsilon,$$

i.e.,

$$\forall \varepsilon > 0, \exists \delta > 0, f(B_{(X, d)}(x_0, \delta) \cap S) \subseteq B_{(Y, d')}(f(x_0), \varepsilon),$$

i.e.,

for any open neighborhood V of $f(x_0)$,
there exists an open neighborhood U of x_0 such that $f(U \cap S) \subseteq V$.

If f is continuous at x_0 for every x_0 in S , f is continuous on S .

Remark 508 If x_0 is an isolated point of S , f is continuous at x_0 . If x_0 is an accumulation point for S , f is continuous at x_0 if and only if $\lim_{x \rightarrow x_0} f(x) = f(x_0)$.

Proposition 509 Suppose that $Z \subseteq X''$, where (X''', d'') is a metric space and

$$f : S \rightarrow T, \quad g : W \supseteq f(S) \rightarrow Z$$

$$h : S \rightarrow Z, \quad h(x) = g(f(x))$$

If f is continuous at $x_0 \in S$ and g is continuous at $f(x_0)$, then h is continuous at x_0 .

Proof. Exercise (see Apostol (1974), page 79) or Ok, page 206. ■

Proposition 510 Take $f, g : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$. If f and g are continuous, then

1. $f + g$ is continuous;
2. $f \cdot g$ is continuous;
3. if $\forall x \in S, g(x) \neq 0$, $\frac{f}{g}$ is continuous.

Proof. If x_0 is an isolated point of S , from Remark 508, we are done. If x_0 is an accumulation point for S , the result follows from Remark 508 and Proposition 504. ■

Proposition 511 Let $f : S \subseteq X \rightarrow \mathbb{R}^m$, and for any $j \in \{1, \dots, m\}$ $f_j : S \rightarrow \mathbb{R}$ be such that $\forall x \in S$,

$$f(x) = (f_j(x))_{j=1}^m$$

Then,

$$\langle f \text{ is continuous} \rangle \Leftrightarrow \langle \forall j \in \{1, \dots, m\}, f_j \text{ is continuous} \rangle$$

Proof. The proof follows the strategy used in Proposition 421. ■

Definition 512 Given for any $i \in \{1, \dots, n\}$, $S_i \subseteq \mathbb{R}$, $f : \times_{i=1}^n S_i \rightarrow \mathbb{R}$ is continuous in each variable separately if $\forall i \in \{1, \dots, n\}$ and $\forall x_i^0 \in S_i$,

$$f_{x_i^0} : \times_{k \neq i} S_k \rightarrow \mathbb{R},$$

$$f_{x_i^0}((x_k)_{k \neq i}) = f(x_1, \dots, x_{i-1}, x_i^0, x_{i+1}, \dots, x_n)$$

is continuous.

Proposition 513 Given for any $i \in \{1, \dots, n\}$,

$$f : \times_{i=1}^n S_i \rightarrow \mathbb{R} \text{ is continuous} \Rightarrow f \text{ is continuous in each variable separately}$$

Proof. Exercise. ■

Remark 514 It is false that

$$f \text{ is continuous in each variable separately} \Rightarrow f \text{ is continuous}$$

To see that consider $f : \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$f(x, y) = \begin{cases} \frac{xy}{x^2+y^2} & \text{if } (x, y) \neq 0 \\ 0 & \text{if } (x, y) = 0 \end{cases}$$

The following Proposition is useful to show continuity of functions using the results about continuity of functions from \mathbb{R} to \mathbb{R} .

Proposition 515 For any $k \in \{1, \dots, n\}$, take $S_k \subseteq X$, and define $S := \times_{k=1}^n S_k \subseteq X^n$. Moreover, take $i \in \{1, \dots, n\}$ and let

$$g : S_i \rightarrow Y, \quad : x_i \mapsto g(x_i)$$

be a continuous function and

$$f : S \rightarrow Y, \quad (x_k)_{k=1}^n \mapsto g(x_i).$$

Then f is continuous.

Example 516 An example of the objects described in the above Proposition is the following one.

$$g : [0, \pi] \rightarrow \mathbb{R}, \quad g(x) = \sin x,$$

$$f : [0, \pi] \times [-\pi, 0] \rightarrow \mathbb{R}, \quad f(x, y) = \sin x.$$

Proof. of Proposition 515. We want to show that

$$\forall x_0 \in S, \forall \varepsilon > 0 \quad \exists \delta > 0 \text{ such that } d(x, x_0) < \delta \wedge x \in S \Rightarrow d(f(x), f(x_0)) < \varepsilon$$

We know that

$$\forall x_{i0} \in S_i, \forall \varepsilon > 0 \quad \exists \delta' > 0 \text{ such that } d(x_i, x_{i0}) < \delta' \wedge x_i \in S \Rightarrow d(g(x_i), g(x_{i0})) < \varepsilon$$

Take $\delta = \delta'$. Then $d(x, x_0) < \delta \wedge x \in S \Rightarrow d(x_i, x_{i0}) < \delta' \wedge x_i \in S$ and $\varepsilon > d(g(x_i), g(x_{i0})) = d(f(x), f(x_0))$, as desired. ■

Exercise 517 Show that the following function is continuous.

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^3, \quad f(x_1, x_2) = \begin{pmatrix} e^{x_1} + \cos(x_1 \cdot x_2) \\ \frac{\sin^2 x_1}{e^{x_2}} \\ x_1 + x_2 \end{pmatrix}$$

From Proposition 511, it suffices to show that each component function is continuous. We are going to show that $f_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$f_1(x_1, x_2) = e^{x_1} + \cos(x_1 \cdot x_2)$$

is continuous, leaving the proof of the continuity of the other component functions to the reader.

1. $f_{11} : \mathbb{R}^2 \rightarrow \mathbb{R}$, $f_{11}(x_1, x_2) = e^{x_1}$ is continuous from Proposition 515 and “Calculus 1”;
2. $h_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$, $h_1(x_1, x_2) = x_1$ is continuous from Proposition 515 and “Calculus 1”,
 $h_2 : \mathbb{R}^2 \rightarrow \mathbb{R}$, $h_2(x_1, x_2) = x_2$ is continuous from Proposition 515 and “Calculus 1”,
 $g : \mathbb{R}^2 \rightarrow \mathbb{R}$, $g(x_1, x_2) = h_1(x_1, x_2) \cdot h_2(x_1, x_2) = x_1 \cdot x_2$ is continuous from Proposition 510.2,
 $\phi : \mathbb{R} \rightarrow \mathbb{R}$, $\phi(x) = \cos x$ is continuous from “Calculus 1”,
 $f_{12} : \mathbb{R}^2 \rightarrow \mathbb{R}$, $f_{12}(x_1, x_2) = (\phi \circ g)(x_1, x_2) = \cos(x_1 \cdot x_2)$ is continuous from Proposition 509 (continuity of composition).
3. $f_1 = f_{11} + f_{12}$ is continuous from Proposition 510.1.

The following Proposition is useful in the proofs of several results.

Proposition 518 Let S, T be arbitrary sets, $f : S \rightarrow T$, $\{A_i\}_{i=1}^n$ a family of subsets of S and $\{B_i\}_{i=1}^n$ a family of subsets of T . Then

1. “inverse image preserves inclusions, unions, intersections and set differences”, i.e.,
 - a. $B_1 \subseteq B_2 \Rightarrow f^{-1}(B_1) \subseteq f^{-1}(B_2)$,
 - b. $f^{-1}(\cup_{i=1}^n B_i) = \cup_{i=1}^n f^{-1}(B_i)$,
 - c. $f^{-1}(\cap_{i=1}^n B_i) = \cap_{i=1}^n f^{-1}(B_i)$,
 - d. $f^{-1}(B_1 \setminus B_2) = f^{-1}(B_1) \setminus f^{-1}(B_2)$,
2. “image preserves inclusions, unions, only”, i.e.,
 - e. $A_1 \subseteq A_2 \Rightarrow f(A_1) \subseteq f(A_2)$,
 - f. $f(\cup_{i=1}^n A_i) = \cup_{i=1}^n f(A_i)$,
 - g. $f(\cap_{i=1}^n A_i) \subseteq \cap_{i=1}^n f(A_i)$, and
 if f is one-to-one, then $f(\cap_{i=1}^n A_i) = \cap_{i=1}^n f(A_i)$,
 - h. $f(A_1 \setminus A_2) \supseteq f(A_1) \setminus f(A_2)$, and
 if f is one-to-one and onto, then $f(A_1 \setminus A_2) = f(A_1) \setminus f(A_2)$,
3. “relationship between image and inverse image”

- i. $A_1 \subseteq f^{-1}(f(A_1))$, and
 if f is one-to-one, then $A_1 = f^{-1}(f(A_1))$,
 1. $B_1 \supseteq f(f^{-1}(B_1))$, and
 if f is onto, then $B_1 = f(f^{-1}(B_1))$.

Proof.

...

g.

- (i). $y \in f(A_1 \cap A_2) \Leftrightarrow \exists x \in A_1 \cap A_2$ such that $f(x) = y$;
 (ii). $y \in f(A_1) \cap f(A_2) \Leftrightarrow y \in f(A_1) \wedge y \in f(A_2) \Leftrightarrow (\exists x_1 \in A_1 \text{ such that } f(x_1) = y) \wedge (\exists x_2 \in A_2 \text{ such that } f(x_2) = y)$
 To show that (i) \Rightarrow (ii) it is enough to take $x_1 = x$ and $x_2 = x$.
 ... ■

Proposition 519 $f : X \rightarrow Y$ is continuous \Leftrightarrow

$$V \subseteq Y \text{ is open} \Rightarrow f^{-1}(V) \subseteq X \text{ is open.}$$

Proof. [\Rightarrow]

Take a point $x_0 \in f^{-1}(V)$. We want to show that

$$\exists r > 0 \text{ such that } B(x_0, r) \subseteq f^{-1}(V)$$

Define $y_0 = f(x_0) \in V$. Since $V \subseteq Y$ is open,

$$\exists \varepsilon > 0 \text{ such that } B(y_0, \varepsilon) \subseteq V \quad (10.2)$$

Since f is continuous,

$$\forall \varepsilon > 0, \exists \delta > 0, f(B(x_0, \delta)) \subseteq B(f(x_0), \varepsilon) = B(y_0, \varepsilon) \quad (10.3)$$

Then, taken $r = \delta$, we have

$$B(x_0, r) = B(x_0, \delta) \stackrel{(1)}{\subseteq} f^{-1}(f(B(x_0, \delta))) \stackrel{(2)}{\subseteq} f^{-1}(B(y_0, \varepsilon)) \stackrel{(3)}{\subseteq} f^{-1}(V)$$

where (1) follows from 3.i in Proposition 518,

(2) follows from 1.a in Proposition 518 and (10.3)

(3) follows from 1.a in Proposition 518 and (10.2).

[\Leftarrow]

Take $x_0 \in X$ and define $y_0 = f(x_0)$; we want to show that f is continuous at x_0 .

Take $\varepsilon > 0$, then $B(y_0, \varepsilon)$ is open and, by assumption,

$$f^{-1}(B(y_0, \varepsilon)) \subseteq X \text{ is open.} \quad (10.4)$$

Moreover, by definition of y_0 ,

$$x_0 \in f^{-1}(B(y_0, \varepsilon)) \quad (10.5)$$

(10.4) and (10.5) imply that

$$\exists \delta > 0 \text{ such that } B(x_0, \delta) \subseteq f^{-1}(B(y_0, \varepsilon)) \quad (10.6)$$

Then

$$f(B(x_0, \delta)) \stackrel{(1)}{\subseteq} f(f^{-1}(B(y_0, \varepsilon))) \stackrel{(2)}{\subseteq} B(y_0, \varepsilon)$$

where

(1) follows from 2.e in Proposition 518 and (10.6),

(2) follows from 2.1 in Proposition 518 ■

Proposition 520 $f : X \rightarrow Y$ is continuous \Leftrightarrow

$$V \subseteq Y \text{ closed} \Rightarrow f^{-1}(V) \subseteq X \text{ closed.}$$

Proof. $[\Rightarrow]$

V closed in $Y \Rightarrow Y \setminus V$ open. Then

$$f^{-1}(Y \setminus V) = f^{-1}(Y) \setminus f^{-1}(V) = X \setminus f^{-1}(V) \quad (10.7)$$

where the first equality follows from 1.d in Proposition 518.

Since f is continuous and $Y \setminus V$ open, then from (10.7) $X \setminus f^{-1}(V) \subseteq X$ is open and therefore $f^{-1}(V)$ is closed.

$[\Leftarrow]$

We want to show that for every open set V in Y , $f^{-1}(V)$ is open.

$$V \text{ open} \Rightarrow Y \setminus V \text{ closed} \Rightarrow f^{-1}(Y \setminus V) \text{ closed} \Leftrightarrow X \setminus f^{-1}(V) \text{ closed} \Leftrightarrow f^{-1}(V) \text{ open.}$$

■

Definition 521 A function $f : X \rightarrow Y$ is open if

$$S \subseteq X \text{ open} \Rightarrow f(S) \text{ open;}$$

it is closed if

$$S \subseteq X \text{ closed} \Rightarrow f(S) \text{ closed.}$$

Exercise 522 Through simple examples show the relationship between open, closed and continuous functions.

We can summarize our discussion on continuous function in the following Proposition.

Proposition 523 Let f be a function between metric spaces (X, d) and (Y, d') . Then the following statements are equivalent:

1. f is continuous;
2. $V \subseteq Y$ is open $\Rightarrow f^{-1}(V) \subseteq X$ is open;
3. $V \subseteq Y$ closed $\Rightarrow f^{-1}(V) \subseteq X$ closed;
4. $\forall x_0 \in X, \forall (x_n)_{n \in \mathbb{N}} \in X^\infty$ such that $\lim_{n \rightarrow +\infty} x_n = x_0, \quad \lim_{n \rightarrow +\infty} f(x_n) = f(x_0)$.

10.3 Continuous functions on compact sets

Proposition 524 Given $f : X \rightarrow Y$, if S is a compact subset of X and f is continuous, then $f(S)$ is a compact subset (of Y).

Proof. Let \mathcal{F} be an open covering of $f(S)$, so that

$$f(S) \subseteq \cup_{A \in \mathcal{F}} A. \quad (10.8)$$

We want to show that \mathcal{F} admits an open subcover which covers $f(S)$. Since f is continuous,

$$\forall A \in \mathcal{F}, \quad f^{-1}(A) \text{ is open in } X$$

Moreover,

$$S \stackrel{(1)}{\subseteq} f^{-1}(f(S)) \stackrel{(2)}{\subseteq} f^{-1}(\cup_{A \in \mathcal{F}} A) \stackrel{(3)}{=} \cup_{A \in \mathcal{F}} f^{-1}(A)$$

where

- (1) follows from 3.i in Proposition 518,
- (2) follows from 1.a in Proposition 518 and (10.8),
- (3) follows from 1.b in Proposition 518.

In other words $\{f^{-1}(A)\}_{A \in \mathcal{F}}$ is an open cover of S . Since S is compact there exists $A_1, \dots, A_n \in \mathcal{F}$ such that

$$S \subseteq \cup_{i=1}^n f^{-1}(A_i).$$

Then

$$f(S) \stackrel{(1)}{\subseteq} f\left(\bigcup_{i=1}^n (f^{-1}(A_i))\right) \stackrel{(2)}{=} \bigcup_{i=1}^n f(f^{-1}(A_i)) \stackrel{(3)}{\subseteq} \bigcup_{i=1}^n A_i$$

where

- (1) follows from 1.a in Proposition 518,
- (2) follows from 2.f in Proposition 518,
- (3) follows from 3.1 in Proposition 518. ■

Proposition 525 (Extreme Value Theorem) *If S a nonempty, compact subset of X and $f : S \rightarrow \mathbb{R}$ is continuous, then f admits global maximum and minimum on S , i.e.,*

$$\exists x_{\min}, x_{\max} \in S \text{ such that } \forall x \in S, \quad f(x_{\min}) \leq f(x) \leq f(x_{\max}).$$

Proof. From the previous Proposition $f(S)$ is closed and bounded. Therefore, since $f(S)$ is bounded, there exists $M = \sup f(S)$. By definition of sup,

$$\forall \varepsilon > 0, \quad B(M, \varepsilon) \cap f(S) \neq \emptyset$$

Then¹, $\forall n \in \mathbb{N}$, take

$$\alpha_n \in B\left(M, \frac{1}{n}\right) \cap f(S).$$

Then, $(\alpha_n)_{n \in \mathbb{N}}$ is such that $\forall n \in \mathbb{N}$, $\alpha_n \in f(S)$ and $0 < d(\alpha_n, M) < \frac{1}{n}$. Therefore, $\alpha_n \rightarrow M$, and since $f(S)$ is closed, $M \in f(S)$. But $M \in f(S)$ means that $\exists x_{\max} \in S$ such that $f(x_{\max}) = M$ and the fact that $M = \sup f(S)$ implies that $\forall x \in S, \quad f(x) \leq f(x_{\max})$. Similar reasoning holds for x_{\min} . ■

We conclude the section showing a result useful in itself and needed to show the inverse function theorem - see Section 15.3.

Proposition 526 *Let $f : X \rightarrow Y$ be a function from a metric space (X, d) to another metric space (Y, d') . Assume that f is one-to-one and onto. If X is compact and f is continuous, then the inverse function f^{-1} is continuous.*

Proof. Exercise. ■

We are going to use the above result to show that a “well behaved” consumer problem does have a solution.

Let the following objects be given.

Price vector $p \in \mathbb{R}_{++}^n$, consumption vector $x \in \mathbb{R}^n$, consumer's wealth $w \in \mathbb{R}_{++}$, continuous utility function $u : \mathbb{R}^n \rightarrow \mathbb{R}$, $x \mapsto u(x)$. The consumer solves the following problem. For given, $p \in \mathbb{R}_{++}^n, w \in \mathbb{R}_{++}$, find x which gives the maximum value to the utility function u under the constraint $x \in C(p, w)$ defined as

$$\{x = (x_i)_{i=1}^n \in \mathbb{R}^n : \forall i \in \{1, \dots, n\}, x_i \geq 0 \text{ and } px \leq w\}.$$

As an application of Propositions 525 and 444, we have to show that for any $p \in \mathbb{R}_{++}^n, w \in \mathbb{R}_{++}$,

- 1. $C(p, w) \neq \emptyset$,
- 2. $C(p, w)$ is bounded, and
- 3. $C(p, w)$ is closed,
- 1. $0 \in C(p, w)$.

2. Clearly if $S \subseteq \mathbb{R}^n$, then S is bounded iff

S is bounded below, i.e., $\exists \underline{x} = (\underline{x}_i)_{i=1}^n \in \mathbb{R}^n$ such that $\forall x = (x_i)_{i=1}^n \in S$, we have that $\forall i \in \{1, \dots, n\}, x_i \geq \underline{x}_i$, and

S is bounded above, i.e., $\exists \bar{x} = (\bar{x}_i)_{i=1}^n \in \mathbb{R}^n$ such that $\forall x = (x_i)_{i=1}^n \in S$, we have that $\forall i \in \{1, \dots, n\}, x_i \leq \bar{x}_i$.

$C(p, w)$ is bounded below by zero, i.e., we can take $\underline{x} = 0$. $C(p, w)$ is bounded above because for every $i \in \{1, \dots, n\}$,

$$x_i \leq \frac{w - \sum_{i' \neq i} p_{i'} x_{i'}}{p_i} \leq \frac{w}{p_i},$$

¹The fact that $M \in f(S)$ can be also proved as follows: from Proposition 485, $M \in \text{Cl } f(S) = f(S)$, where the last equality follows from the fact that $f(S)$ is closed.

where the first inequality comes from the fact that $px \leq w$, and the second inequality from the fact that $p \in \mathbb{R}_{++}^n$ and $x \in \mathbb{R}_+^n$. Then we can take $\bar{x} = (m, m, \dots, m)$, where $m = \max \left\{ \frac{w}{p_i} \right\}_{i=1}^n$.

3. Define

$$\text{for } i \in \{1, \dots, n\}, \quad g_i : \mathbb{R}^n \rightarrow \mathbb{R}, \quad x = (x_i)_{i=1}^n \mapsto x_i,$$

and

$$h : \mathbb{R}^n \rightarrow \mathbb{R}, \quad x = (x_i)_{i=1}^n \mapsto w - px.$$

All the above functions are continuous and clearly,

$$C(p, w) = \{x = (x_i)_{i=1}^n \in \mathbb{R}^n : \forall i \in \{1, \dots, n\}, \quad g_i(x) \geq 0 \text{ and } h(x) \geq 0\}.$$

Moreover,

$$\begin{aligned} C(p, w) &= \{x = (x_i)_{i=1}^n \in \mathbb{R}^n : \forall i \in \{1, \dots, n\}, \quad g_i(x) \in [0, +\infty) \text{ and } h(x) \in [0, +\infty)\} = \\ &= \cap_{i=1}^n g_i^{-1}([0, +\infty)) \cap h^{-1}([0, +\infty)). \end{aligned}$$

$[0, +\infty)$ is a closed set and since for any g_i is continuous and h is continuous, from Proposition 523.3, the following sets are closed

$$\forall i \in \{1, \dots, n\}, \quad g_i^{-1}([0, +\infty)) \text{ and } h^{-1}([0, +\infty)).$$

Then the desired result follows from the fact that intersection of closed set is closed.

10.4 Exercises

From Lipschutz (1965), starting from page 61: 30, 32, 34; starting from page 106: 19, 20.

Chapter 11

Correspondence, maximum theorem and a fixed point theorem

11.1 Continuous Correspondences

Definition 527 Consider¹ two metric spaces (X, d_X) and (Y, d_Y) . A correspondence φ from X to Y is a rule which associates a subset of Y with each element of X , and it is described by the notation

$$\varphi : X \rightarrow\!\!\rightarrow Y, \quad \varphi : x \mapsto\!\!\mapsto \varphi(x).$$

Remark 528 In other words, a correspondence $\varphi : X \rightarrow\!\!\rightarrow Y$ can be identified with a function from X to 2^Y (the set of all subsets of Y). If we identify x with $\{x\}$, a function from X to Y can be thought as a particular correspondence.

Remark 529 Some authors make part of the Definition of correspondence the fact that φ is not empty valued, i.e., that $\forall x \in X, \varphi(x) \neq \emptyset$.

In what follows, unless otherwise stated, (X, d_X) and (Y, d_Y) are assumed to be metric spaces and are denoted by X and Y , respectively.

Definition 530 Given $U \subseteq X$, $\varphi(U) = \cup_{x \in U} \varphi(x) = \{y \in Y : \exists x \in U \text{ such that } y \in \varphi(x)\}$.

Definition 531 The graph of $\varphi : X \rightarrow\!\!\rightarrow Y$ is

$$\text{graph } \varphi := \{(x, y) \in X \times Y : y \in \varphi(x)\}.$$

Definition 532 Consider $\varphi : X \rightarrow\!\!\rightarrow Y$. φ is Upper Hemi-Continuous (UHC) at $x \in X$ if $\varphi(x) \neq \emptyset$ and for every open neighborhood V of $\varphi(x)$, there exists an open neighborhood U of x such that for every $x' \in U$, $\varphi(x') \subseteq V$ (or $\varphi(U) \subseteq V$).

φ is UHC if it is UHC at every $x \in X$.

Definition 533 Consider $\varphi : X \rightarrow\!\!\rightarrow Y$. φ is Lower Hemi-Continuous (LHC) at $x \in X$ if $\varphi(x) \neq \emptyset$ and for any open set V in Y such that $\varphi(x) \cap V \neq \emptyset$, there exists an open neighborhood U of x such that for every $x' \in U$, $\varphi(x') \cap V \neq \emptyset$.

φ is LHC if it is LHC at every $x \in X$.

Example 534 Consider $X = \mathbb{R}_+$ and $Y = [0, 1]$, and

$$\begin{aligned} \varphi_1(x) &= \begin{cases} [0, 1] & \text{if } x = 0 \\ \{0\} & \text{if } x > 0. \end{cases} \\ \varphi_2(x) &= \begin{cases} \{0\} & \text{if } x = 0 \\ [0, 1] & \text{if } x > 0. \end{cases} \end{aligned}$$

φ_1 is UHC and not LHC; φ_2 is LHC and not UHC.

¹This chapter is based mainly on McLean (1985) and Hildebrand (1974).

Some (partial) intuition about the above definitions can be given as follows.

Upper Hemi-Continuity does not allow "explosions". In other words, φ is not UHC at x if there exists a small enough open neighborhood of x such that φ does "explode", i.e., it becomes much bigger in that neighborhood.

Lower Hemi-Continuity does not allow "implosions". In other words, φ is not LHC at x if there exists a small enough open neighborhood of x such that φ does "implode", i.e., it becomes much smaller in that neighborhood.

In other words, "UHC \Rightarrow no explosion" and "LHC \Rightarrow no implosion" (or "explosion \Rightarrow not UHC" and "implosion \Rightarrow not LHC"). On the other hand, opposite implications are false, i.e.,

it is false that "explosion \Leftarrow not UHC" and "implosion \Leftarrow not LHC", or, in an equivalent manner,

it is false that "no explosion \Rightarrow UHC" and "no implosion \Rightarrow LHC".

An example of a correspondence which neither explodes nor implodes and which is not UHC and not LHC is presented below.

$$\varphi : \mathbb{R}_+ \rightarrow \rightarrow \mathbb{R}, \quad \varphi : x \mapsto \begin{cases} [1, 2] & \text{if } x \in [0, 1) \\ [3, 4] & \text{if } x \in [1, +\infty) \end{cases}$$

φ does not implode or explode if you move away from 1 (in a small open neighborhood of 1): on the right of 1, φ does not change; on the left, it changes completely. Clearly, φ is neither UHC nor LHC (in 1).

The following correspondence is both UHC and LHC:

$$\varphi : \mathbb{R}_+ \rightarrow \rightarrow \mathbb{R}, \quad \varphi : x \mapsto [x, x + 1]$$

A, maybe disturbing, example is the following one

$$\varphi : \mathbb{R}_+ \rightarrow \rightarrow \mathbb{R}, \quad \varphi : x \mapsto (x, x + 1).$$

Observe that the graph of the correspondence under consideration "does not implode, does not explode, does not jump". In fact, the above correspondence is LHC, but it is not UHC in any $x \in \mathbb{R}_+$, as verified below. We want to show that

$$\text{not} \left\langle \begin{array}{l} \text{for every neighborhood } V \text{ of } \varphi(x), \\ \text{there exists a neighborhood } U \text{ of } x \text{ such that for every } x' \in U, \varphi(x') \subseteq V \end{array} \right\rangle$$

i.e.,

$$\left\langle \begin{array}{l} \text{there exists a neighborhood } V^* \text{ of } \varphi(x) \text{ such that.} \\ \text{for every neighborhood } U \text{ of } x \text{ there exists } x' \in U \text{ such that } \varphi(x') \not\subseteq V^* \end{array} \right\rangle$$

Just take $V = \varphi(x) = (x, x + 1)$; then for any open neighborhood U of x and, in fact, $\forall x' \in U \setminus \{x\}$, $\varphi(x') \not\subseteq V$.

Example 535 *The correspondence below*

$$\varphi : \mathbb{R}_+ \rightarrow \rightarrow \mathbb{R}, \quad \varphi : x \mapsto \begin{cases} [1, 2] & \text{if } x \in [0, 1] \\ [3, 4] & \text{if } x \in [1, +\infty) \end{cases}$$

is UHC, but not LHC.

Remark 536 *Summarizing the above results, we can maybe say that a correspondence which is both UHC and LHC, in fact a continuous correspondence, is a correspondence which agrees with our intuition of a graph without explosions, implosions or jumps.*

Proposition 537 1. *If $\varphi : X \rightarrow \rightarrow Y$ is either UHC or LHC and it is a function, then it is a continuous function.*

2. *If $\varphi : X \rightarrow \rightarrow Y$ is a continuous function, then it is a UHC and LHC correspondence.*

Proof.

1.

Case 1. φ is UHC.

First proof. Use the fourth characterization of continuous function in Definition 507.

Second proof. Recall that a function $f : X \rightarrow Y$ is continuous iff $[V \text{ open in } Y] \Rightarrow [f^{-1}(V) \text{ open in } X]$. Take V open in Y . Consider $x \in f^{-1}(V)$, i.e., x such that $f(x) \in V$. By assumption f is UHC and therefore \exists an open neighborhood U of x such that $f(U) \subseteq V$. Then, $U \subseteq f^{-1} \circ f(U) \subseteq f^{-1}(V)$. Then, for any $x \in f^{-1}(V)$, we have found an open set U which contains x and is contained in $f^{-1}(V)$, i.e., $f^{-1}(V)$ is open.

Case 2. φ is LHC.

See Remark 541 below.

2.

The results follows from the definitions and again from Remark 541 below.

■

Definition 538 $\varphi : X \rightarrow \rightarrow Y$ is a continuous correspondence if it is both UHC and LHC.

Very often, checking if a correspondence is UHC or LHC is not easy. We present some *related* concepts which are more convenient to use.

Definition 539 $\varphi : X \rightarrow \rightarrow Y$ is “sequentially LHC” at $x \in X$ if

for every sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$ such that $x_n \rightarrow x$, and for every $y \in \varphi(x)$, there exists a sequence $(y_n)_{n \in \mathbb{N}} \in Y^\infty$ such that $\forall n \in \mathbb{N}$, $y_n \in \varphi(x_n)$ and $y_n \rightarrow y$, φ is “sequentially LHC” if it is “sequentially LHC” at every $x \in X$.

Proposition 540 Consider $\varphi : X \rightarrow \rightarrow Y$. φ is LHC at $x \in X \Leftrightarrow \varphi$ is LHC in terms of sequences at $x \in X$.

Proof.

$[\Rightarrow]$

Consider an arbitrary sequence $(x_n)_{n \in \mathbb{N}} \subseteq X^\infty$ such that $x_n \rightarrow x$ and an arbitrary $y \in \varphi(x)$. For every $r \in \mathbb{N}$, consider $B(y, \frac{1}{r})$. Clearly $B(y, \frac{1}{r}) \cap \varphi(x) \neq \emptyset$, since y belongs to both sets. From the fact that φ is LHC, we have that

$$\forall r \in \mathbb{N}, \exists \text{ a neighborhood } U_r \text{ of } x \text{ such that } \forall z \in U_r, \varphi(z) \cap B(y, \frac{1}{r}) \neq \emptyset. \quad (1)$$

Since $x_n \rightarrow x$,

$$\forall r \in \mathbb{N}, \exists n_r \in \mathbb{N} \text{ such that } n \geq n_r \Rightarrow x_n \in U_r \quad (2).$$

Consider $\{n_1, \dots, n_r, \dots\}$. For any $r \in \mathbb{N}$, if $n_r < n_{r+1}$, define $n'_r := n_r$ and $n'_{r+1} := n_{r+1}$, i.e., “just add a ' to the name of those indices”. If $n_r \geq n_{r+1}$, define $n'_r := n_r$ and $n'_{r+1} := n_r + 1$. Then, $\forall r \in \mathbb{N}$, $n'_{r+1} > n'_r$ and condition (2) still hold, i.e.,

$$\forall r, \exists n'_r \text{ such that } n \geq n'_r \Rightarrow x_n \in U_r \text{ and } n'_{r+1} > n'_r. \quad (3)$$

We can now define the desired sequence $(y_n)_{n \in \mathbb{N}}$. For any $n \in [n'_r, n'_{r+1})$, observe that from (3), $x_n \in U_r$, and, then, from (1), $\varphi(x_n) \cap B(y, \frac{1}{r}) \neq \emptyset$. Then,

$$\text{for any } r, \text{ for any } n \in [n'_r, n'_{r+1}) \cap \mathbb{N}, \text{ choose } y_n \in \varphi(x_n) \cap B(y, \frac{1}{r}) \quad (4).$$

We are left with showing that $y_n \rightarrow y$, i.e., $\forall \varepsilon > 0, \exists \overline{m}$ such that $n \geq \overline{m} \Rightarrow y_n \in B(y, \varepsilon)$. Observe that (4) just says that

- for any $n \in [n'_1, n'_2)$, $y_n \in \varphi(x_n) \cap B(y, \frac{1}{1})$,
- for any $n \in [n'_2, n'_3)$, $y_n \in \varphi(x_n) \cap B(y, \frac{1}{2}) \subseteq B(y, 1)$
- ...
- for any $n \in [n'_r, n'_{r+1})$, $y_n \in \varphi(x_n) \cap B(y, \frac{1}{r}) \subseteq B(y, \frac{1}{r-1})$,

and so on. Then, for any $\varepsilon > 0$, choose $r > \frac{1}{\varepsilon}$ (so that $\frac{1}{r} < \varepsilon$) and $\overline{m} = n'_r$, then from the above observations, $n \in [\overline{m}, n'_{r+1}) \Rightarrow y_n \in B(y, \frac{1}{r}) \subseteq B(y, \varepsilon)$, and for $n > n'_{r+1}$, a fortiori, $y_n \in B(y, \frac{1}{r})$.
[\Leftarrow]

Assume otherwise, i.e., \exists an open set V such that

$$\varphi(x) \cap V \neq \emptyset \quad (5)$$

and such that \forall open neighborhood U of x $\exists x_U \in U$ such that $\varphi(x_U) \cap V = \emptyset$.

Consider the following family of open neighborhood of x :

$$\left\{ B\left(x, \frac{1}{n}\right) : n \in \mathbb{N} \right\}.$$

Then $\forall n \in \mathbb{N}$, $\exists x_n \in B(x, \frac{1}{n})$, and therefore $x_n \rightarrow x$, such that

$$\varphi(x_n) \cap V = \emptyset \quad (6).$$

From (5), we can take $y \in \varphi(x) \cap V$. By assumption, we know that there exists a sequence $(y_n)_{n \in \mathbb{N}} \in Y^\infty$ such that $\forall n \in \mathbb{N}$, $y_n \in \varphi(x_n)$ and $y_n \rightarrow y$. Since V is open and $y \in V$, $\exists \overline{n}$ such that $n > \overline{n} \Rightarrow y_n \in V$. Therefore,

$$y \in \varphi(x_n) \cap V \quad (7).$$

But (7) contradicts (6).
■

Thanks to the above Proposition from now on we talk simply of Lower Hemi-Continuous correspondences.

Remark 541 If $\varphi : X \rightarrow Y$ is LHC and it is a function, then it is a continuous function. The result follows from the characterization of Lower Hemi-Continuity in terms of sequences and from the characterization of continuous functions presented in Proposition 523.

Definition 542 $\varphi : X \rightarrow Y$ is closed, or "sequentially UHC", at $x \in X$ if

for every sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$ such that $x_n \rightarrow x$, and for every sequence $(y_n)_{n \in \mathbb{N}} \in Y^\infty$ such that $y_n \in \varphi(x_n)$ and $y_n \rightarrow y$,
it is the case that $y \in \varphi(x)$.
 φ is closed if it is closed at every $x \in X$.

Proposition 543 φ is closed \Leftrightarrow graph φ is a closed set in $X \times Y$.²

Proof. An equivalent way of stating of the above Definition is the following one: for every sequence $(x_n, y_n)_{n \in \mathbb{N}} \in (X \times Y)^\infty$ such that $\forall n \in \mathbb{N}$, $(x_n, y_n) \in \text{graph } \varphi$ and $(x_n, y_n) \rightarrow (x, y)$, it is the case that $(x, y) \in \text{graph } \varphi$. Then, from the characterization of closed sets in terms of sequences, i.e., Proposition 425, the desired result follows. ■

Remark 544 Because of the above result, many author use the expression " φ has closed graph" in the place of " φ is closed".

Remark 545 The definition of closed correspondence does NOT reduce to continuity in the case of functions, as the following example shows.

$$\varphi_3 : \mathbb{R}_+ \rightarrow \mathbb{R}, \quad \varphi_3(x) = \begin{cases} \{0\} & \text{if } x = 0 \\ \{\frac{1}{x}\} & \text{if } x > 0. \end{cases}$$

φ is a closed correspondence, but it is not a continuous function.

Definition 546 A set $C \subseteq \mathbb{R}^n$ is convex if $\forall x_1, x_2 \in C$ and $\forall \lambda \in [0, 1]$, $(1 - \lambda)x_1 + \lambda x_2 \in C$.

Definition 547 $\varphi : X \rightarrow Y$ is closed (non-empty, convex, compact ...) valued if for every $x \in X$, $\varphi(x)$ is a closed (non-empty, convex, compact ...) set.

² $((X \times Y), d^*)$ with $d^*((x, x'), (y, y')) = \max\{d(x, x'), d'(y, y')\}$ is a metric space.

Proposition 548 Consider $\varphi : X \rightarrow\rightarrow Y$. φ closed $\begin{smallmatrix} \Rightarrow \\ \nLeftarrow \end{smallmatrix}$ φ closed valued.

Proof.

[\Rightarrow]

We want to show that every sequence in $\varphi(x)$ which converges, in fact, converges in $\varphi(x)$. Choose $x \in X$ and a sequence $(y_n)_{n \in \mathbb{N}}$ such that $\{y_n : n \in \mathbb{N}\} \subseteq \varphi(x)$ and such that $y_n \rightarrow y$. Then setting for any $n \in \mathbb{N}$, $x_n = x$, we get $x_n \rightarrow x$, $y_n \in \varphi(x_n)$, $y_n \rightarrow y$. Then, since φ is closed, $y \in \varphi(x)$. This shows that $\varphi(x)$ is a closed set.

[\nLeftarrow]

φ_2 in Example 534 is closed valued, but not closed.

■

Remark 549 Consider $\varphi : X \rightarrow\rightarrow Y$. φ is UHC $\begin{smallmatrix} \nRightarrow \\ \nLeftarrow \end{smallmatrix}$ φ is closed.

[\nRightarrow]

$\varphi_4 : \mathbb{R}_+ \rightarrow\rightarrow \mathbb{R}$, $\varphi_4(x) = [0, 1]$ for every $x \in \mathbb{R}$ is UHC and not closed.

[\nLeftarrow]

φ_3 in Remark 545 is closed and not UHC, simply because it is not a continuous “function”.

Proposition 550 Consider $\varphi : X \rightarrow\rightarrow Y$. If φ is UHC (at x) and closed valued (at x), then φ is closed (at x).

Proof.

Take an arbitrary $x \in X$. We want to show that φ is closed at x , i.e., assume that $x_n \rightarrow x$, $y_n \in \varphi(x_n)$, $y_n \rightarrow y$; we want to show that $y \in \varphi(x)$. Since $\varphi(x)$ is a closed set, it suffices to show that $y \in Cl\varphi(x)$, i.e.,³ $\forall \varepsilon > 0$, $B(y, \varepsilon) \cap \varphi(x) \neq \emptyset$.

Consider $\{B(z, \frac{\varepsilon}{2}) : z \in \varphi(x)\}$. Then, $\cup_{z \in \varphi(x)} B(z, \frac{\varepsilon}{2}) := V$ is open and contains $\varphi(x)$. Since φ is UHC at x , there exists an open neighborhood U of x such that

$$\varphi(U) \subseteq V. \quad (1)$$

Since $x_n \rightarrow x \in U$, $\exists \hat{n} \in \mathbb{N}$ such that $\forall n > \hat{n}$, $x_n \in U$, and, from (1), $\varphi(x_n) \subseteq V$. Since $y_n \in \varphi(x_n)$,

$$\forall n > \hat{n}, \quad y_n \in V := \cup_{z \in \varphi(x)} B(z, \frac{\varepsilon}{2}). \quad (2)$$

From (2), $\forall n > \hat{n}$, $\exists z_n^* \in \varphi(x)$ such that $y_n \in B(z_n^*, \frac{\varepsilon}{2})$ and then

$$d(y_n, z_n^*) < \frac{\varepsilon}{2}. \quad (3)$$

Since $y_n \rightarrow y$, $\exists n^*$ such that $\forall n > n^*$,

$$d(y_n, y) < \frac{\varepsilon}{2} \quad (4)$$

From (3) and (4), $\forall n > \max\{\hat{n}, n^*\}$, $z_n^* \in \varphi(x)$ and $d(y, z_n^*) \leq d(y, y_n) + d(y_n, z_n^*) < \varepsilon$, i.e., $z_n^* \in B(y, \varepsilon) \cap \varphi(x) \neq \emptyset$.

■

Proposition 551 Consider $\varphi : X \rightarrow\rightarrow Y$. If φ is closed and there exists a compact set $K \subseteq Y$ such that $\varphi(X) \subseteq K$, then φ is UHC.

Therefore, in simpler terms, if φ is closed (at x) and Y is compact, then φ is UHC (at x).

Proof.

Assume that there exists $x \in X$ such that φ is not UHC at $x \in X$, i.e., there exist an open neighborhood V of $\varphi(x)$ such that for every open neighborhood U_x of x , $\varphi(U_x) \cap V^C \neq \emptyset$. In particular, $\forall n \in \mathbb{N}$, $\varphi(B(x, \frac{1}{n})) \cap V^C \neq \emptyset$. Therefore, we can construct a sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$ such that $x_n \rightarrow x$ and $\varphi(x_n) \cap V^C \neq \emptyset$. Now, take $y_n \in \varphi(x_n) \cap V^C$. Since $y_n \in \varphi(X) \subseteq K$ and

³See Corollary 485.

K is compact, and therefore sequentially compact, up to a subsequence, $y_n \rightarrow y \in K$. Moreover, since $\forall n \in \mathbb{N}, y_n \in V^C$ and V^C is closed,

$$y \in V^C \quad (1).$$

Since φ is closed and $x_n \rightarrow x, y_n \in \varphi(x_n), y_n \rightarrow y$, we have that $y \in \varphi(x)$. Since, by assumption, $\varphi(x) \subseteq V$, we have that

$$y \in V \quad (2).$$

But (2) contradicts (1). ■

None of the Assumptions of the above Proposition can be dispensed of. All the examples below show correspondences which are not UHC.

Example 552 1.

$$\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}, \varphi(x) = \begin{cases} \{\frac{1}{2}\} & \text{if } x \in [0, 2] \\ \{1\} & \text{if } x > 2. \end{cases}$$

$Y = [0, 1]$, but φ is not closed.

2.

$$\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}, \varphi(x) = \begin{cases} \{0\} & \text{if } x = 0 \\ \{\frac{1}{x}\} & \text{if } x > 0. \end{cases}$$

φ is closed, but $\varphi(X) = \mathbb{R}_+$, which is closed, but not bounded.

3.

$$\varphi : [0, 1] \rightarrow [0, 1], \varphi(x) = \begin{cases} \{0\} & \text{if } x \in [0, 1) \\ \{\frac{1}{2}\} & \text{if } x = 1. \end{cases}$$

φ is closed (in Y), but $Y = [0, 1)$ is not compact. Observe that if you consider

$$\varphi : [0, 1] \rightarrow [0, 1], \varphi(x) = \begin{cases} \{0\} & \text{if } x \in [0, 1) \\ \{\frac{1}{2}\} & \text{if } x = 1. \end{cases},$$

then φ is not closed.

Definition 553 Consider $\varphi : X \rightarrow Y, V \subseteq Y$.

The strong inverse image of V via φ is

$${}^s\varphi^{-1}(V) := \{x \in X : \varphi(x) \subseteq V\};$$

.

The weak inverse image of V via φ is

$${}^w\varphi^{-1}(V) := \{x \in X : \varphi(x) \cap V \neq \emptyset\}.$$

Remark 554 1. $\forall V \subseteq Y, {}^s\varphi^{-1}(V) \subseteq {}^w\varphi^{-1}(V)$.

2. If φ is a function, the usual definition of inverse image coincides with both above definitions.

Proposition 555 Consider $\varphi : X \rightarrow Y$.

- 1.1. φ is UHC \Leftrightarrow for every open set V in $Y, {}^s\varphi^{-1}(V)$ is open in X ;
- 1.2. φ is UHC \Leftrightarrow for every closed set V in $Y, {}^w\varphi^{-1}(V)$ is closed in X ;
- 2.1. φ is LHC \Leftrightarrow for every open set V in $Y, {}^w\varphi^{-1}(V)$ is open in X ;
- 2.2. φ is LHC \Leftrightarrow for every closed set V in $Y, {}^s\varphi^{-1}(V)$ is closed in X .⁴

Proof.

[1.1., \Rightarrow] Consider V open in Y . Take $x_0 \in {}^s\varphi^{-1}(V)$; by definition of ${}^s\varphi^{-1}$, $\varphi(x_0) \subseteq V$. By definition of UHC correspondence, \exists an open neighborhood U of x_0 such that $\forall x \in U, \varphi(x) \in V$. Then $x_0 \in U \subseteq {}^s\varphi^{-1}(V)$.

[1.1., \Leftarrow] Take an arbitrary $x_0 \in X$ and an open neighborhood V of $\varphi(x_0)$. Then $x_0 \in {}^s\varphi^{-1}(V)$ and ${}^s\varphi^{-1}(V)$ is open by assumption. Therefore (just identifying U with ${}^s\varphi^{-1}(V)$), we have proved that φ is UHC.

⁴Part 2.2 of the Proposition will be used in the proof of the Maximum Theorem.

To show 1.2, preliminarily, observe that

$$(w\varphi^{-1}(V))^C = {}^s\varphi^{-1}(V^C). \quad (11.1)$$

(To see that, simply observe that $(w\varphi^{-1}(V))^C := \{x \in X : \varphi(x) \cap V = \emptyset\}$ and ${}^s\varphi^{-1}(V^C) := \{x \in X : \varphi(x) \subseteq V^C\}$)

[1.2., \Rightarrow] V closed $\Leftrightarrow V^C$ open $\xrightarrow{Assum., (1.1)} {}^s\varphi^{-1}(V^C) \stackrel{(11.1)}{=} (w\varphi^{-1}(V))^C$ open $\Leftrightarrow w\varphi^{-1}(V)$ closed.

[1.2., \Leftarrow]

From (1.1.), it suffices to show that \forall open set V in Y , ${}^s\varphi^{-1}(V)$ is open in X . Then,

V open $\Leftrightarrow V^C$ closed $\xrightarrow{Assum.^w} \varphi^{-1}(V^C)$ closed $\Leftrightarrow (w\varphi^{-1}(V^C))^C \stackrel{(11.1)}{=} {}^s\varphi^{-1}(V)$ open.

The proofs of parts 2.1. and 2.2. are similar to the above ones.

■

Remark 556 Observe that φ is UHC \nRightarrow for every closed set V in Y , ${}^s\varphi^{-1}(V)$ is closed in X .

[\nRightarrow]

Consider

$$\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}, \quad \varphi(x) = \begin{cases} [0, 2] & \text{if } x \in [0, 1] \\ [0, 1] & \text{if } x > 1. \end{cases}$$

φ is UHC and $[0, 1]$ is closed, but ${}^s\varphi^{-1}([0, 1]) := \{x \in \mathbb{R}_+ : \varphi(x) \subseteq [0, 1]\} = (1, +\infty)$ is not closed.

[\nLeftarrow]

Consider

$$\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+, \quad \varphi(x) = \begin{cases} [0, \frac{1}{2}] \cup \{1\} & \text{if } x = 0 \\ [0, 1] & \text{if } x > 0. \end{cases}$$

For any closed set in $Y := \mathbb{R}_+$, ${}^s\varphi^{-1}(V)$ can be only one of the following set, and each of them is closed: $\{0\}, \mathbb{R}_+, \emptyset$. On the other hand, φ is not UHC in 0.

Definition 557 Let the vector spaces (X, d_X) , (Y, d_Y) and (Z, d_Z) and the correspondences $\varphi : X \rightarrow Y, \psi : Y \rightarrow Z$ be given. The composition of φ with ψ is

$$\psi \circ \varphi : X \rightarrow Z,$$

$$(\psi \circ \varphi)(x) := \cup_{y \in \varphi(x)} \psi(y) = \{z \in Z : \exists x \in X \text{ such that } z \in \psi(\varphi(x))\}$$

.

Proposition 558 Consider $\varphi : X \rightarrow Y, \psi : Y \rightarrow Z$. If φ and ψ are UHC, then $\psi \circ \varphi$ is UHC.

Proof.

$$\text{Step 1. } {}^s(\psi \circ \varphi)^{-1}(V) = {}^s\varphi^{-1}({}^s\psi^{-1}(V)).$$

$$\begin{aligned} {}^s(\psi \circ \varphi)^{-1}(V) &= \{x \in X : \psi(\varphi(x)) \subseteq V\} = \{x \in X : \forall y \in \varphi(x), \psi(y) \subseteq V\} = \\ &= \{x \in X : \forall y \in \varphi(x), y \in {}^s\psi^{-1}(V)\} = \{x \in X : \varphi(x) \subseteq {}^s\psi^{-1}(V)\} = {}^s\varphi^{-1}({}^s\psi^{-1}(V)). \end{aligned}$$

Step 2. Desired result.

Take V open in Z . From Theorem 555, we want to show that ${}^s(\psi \circ \varphi)(V)$ is open in X . From step 1, we have that ${}^s(\varphi \circ \psi)^{-1}(V) = {}^s\varphi^{-1}({}^s\psi^{-1}(V))$. Now, ${}^s\psi^{-1}(V)$ is open because ψ is UHC, and ${}^s\varphi^{-1}({}^s\psi^{-1}(V))$ is open because φ is UHC.

■

Proposition 559 Consider $\varphi : X \rightarrow Y$. If φ is UHC and compact valued, and $A \subseteq X$ is a compact set, then $\varphi(A)$ is compact.

Proof.

Consider an arbitrary open cover $\{C_\alpha\}_{\alpha \in I}$ for $\varphi(A)$. Since $\varphi(A) := \cup_{x \in A} \varphi(x)$ and φ is compact valued, there exists a finite set $N_x \subseteq I$ such that

$$\varphi(x) \subseteq \cup_{\alpha \in N_x} C_\alpha := G_x. \quad (11.2)$$

Since for every $\alpha \in N_x$, C_α is open, then G_x is open. Since φ is UHC, ${}^s\varphi^{-1}(G_x)$ is open. Moreover, $x \in {}^s\varphi^{-1}(G_x)$: this is the case because, by definition, $x \in {}^s\varphi^{-1}(G_x)$ iff $\varphi(x) \subseteq G_x$, which is just (11.2). Therefore, $\{{}^s\varphi^{-1}(G_x)\}_{x \in A}$ is an open cover of A . Since, by assumption, A is compact, there exists a finite set $\{x_i\}_{i=1}^m \subseteq A$ such that $A \subseteq \cup_{i=1}^m ({}^s\varphi^{-1}(G_{x_i}))$. Finally,

$$\varphi(A) \subseteq \varphi(\cup_{i=1}^m ({}^s\varphi^{-1}(G_{x_i}))) \stackrel{(1)}{\subseteq} \cup_{i=1}^m \varphi({}^s\varphi^{-1}(G_{x_i})) \stackrel{(2)}{\subseteq} \cup_{i=1}^m G_{x_i} = \cup_{i=1}^m \cup_{\alpha \in N_{x_i}} C_\alpha,$$

and $\left\{ \{C_\alpha\}_{\alpha \in N_{x_i}} \right\}_{i=1}^m$ is a finite subcover of $\{C_\alpha\}_{\alpha \in I}$. We are left with showing (1) and (2) above.

(1). In general, it is the case that $\varphi(\cup_{i=1}^m S_i) \subseteq \cup_{i=1}^m \varphi(S_i)$.

$y \in \varphi(\cup_{i=1}^m S_i) \Leftrightarrow \exists x \in \cup_{i=1}^m S_i$ such that $y \in \varphi(x) \Rightarrow \exists i$ such that $y \in \varphi(x) \subseteq \varphi(S_i) \Rightarrow y \in \cup_{i=1}^m \varphi(S_i)$.

(2). In general, it is the case that $\varphi({}^s\varphi^{-1}(A)) \subseteq A$.

$y \in \varphi({}^s\varphi^{-1}(A)) \Rightarrow \exists x \in {}^s\varphi^{-1}(A)$ such that $y \in \varphi(x)$. But, by definition of ${}^s\varphi^{-1}(A)$, and since $x \in {}^s\varphi^{-1}(A)$, it follows that $\varphi(x) \subseteq A$ and therefore $y \in A$.

■

Remark 560 Observe that the assumptions in the above Proposition cannot be dispensed of, as verified below.

Consider $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}, \varphi(x) = [0, 1]$. Observe that φ is UHC and bounded valued but not closed valued, and $\varphi([0, 1]) = [0, 1]$ is not compact.

Consider $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}, \varphi(x) = \mathbb{R}_+$. Observe that φ is UHC and closed valued, but not bounded valued, and $\varphi([0, 1]) = \mathbb{R}_+$ is not compact.

Consider $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+, \varphi(x) = \begin{cases} \{x\} & \text{if } x \neq 1 \\ \{0\} & \text{if } x = 1. \end{cases}$ Observe that φ is not UHC and $\varphi([0, 1]) = [0, 1]$ is not compact.

Add Proposition 5, page 25 and Proposition 6, page 26, from Hildebrand (1974) ... maybe as exercises ...

Remark 561 Below, we summarize some facts we showed in the present Section, in a somehow informal manner.

$$\langle (\text{if } \varphi \text{ is a fcn., it is cnt.}) \Leftrightarrow \langle \varphi \text{ is UHC} \rangle \not\Rightarrow \langle \varphi \text{ is sequentially UHC, i.e., closed} \rangle \not\Leftarrow \langle (\text{if } \varphi \text{ is a fcn., it is cnt.}) \rangle$$

$$\langle (\text{if } \varphi \text{ is a fcn, it is continuous}) \Leftrightarrow \langle \varphi \text{ is LHC} \rangle \Leftrightarrow \langle \varphi \text{ is sequentially LHC} \rangle$$

$$\langle \varphi \text{ UHC and closed valued at } x \rangle \Rightarrow \langle \varphi \text{ is closed at } x \rangle$$

$$\langle \varphi \text{ UHC at } x \rangle \Leftarrow \langle \varphi \text{ is closed at } x \text{ and } \text{Im } \varphi \text{ compact} \rangle$$

11.2 The Maximum Theorem

Theorem 562 (Maximum Theorem) Let the metric spaces (Π, d_Π) , (X, d_X) , the correspondence $\beta : \Pi \rightarrow X$ and a function $u : X \times \Pi \rightarrow \mathbb{R}$ be given.⁵ Define

$$\begin{aligned} \xi : \Pi &\rightarrow X, \\ \xi(\pi) &= \{z \in \beta(\pi) : \forall x \in \beta(\pi), u(z, \pi) \geq u(x, \pi)\} = \arg \max_{x \in \beta(\pi)} u(x, \pi), \end{aligned}$$

⁵ Obviously, β stands for “budget correspondence” and u for “utility function”.

Assume that

β is non-empty valued, compact valued and continuous,
 u is continuous.

Then

1. ξ is non-empty valued, compact valued, UHC and closed, and
- 2.

$$v : \Pi \rightarrow \mathbb{R}, \quad v : \pi \mapsto \max_{x \in \beta(\pi)} u(x, \pi).$$

is continuous.

Proof.

ξ is non-empty valued.

It is a consequence of the fact that β is non-empty valued and compact valued and of the Extreme Value Theorem - see Proposition 525.

ξ is compact valued.

We are going to show that for any $\pi \in \Pi$, $\xi(\pi)$ is a sequentially compact set. Consider a sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$ such that $\{x_n : n \in \mathbb{N}\} \subseteq \xi(\pi)$. Since $\xi(\pi) \subseteq \beta(\pi)$ and $\beta(\pi)$ is compact by assumption, without loss of generality, up to a subsequence, $x_n \rightarrow x_0 \in \beta(\pi)$. We are left with showing that $x_0 \in \xi(\pi)$. Take an arbitrary $z \in \beta(\pi)$. Since $\{x_n : n \in \mathbb{N}\} \subseteq \xi(\pi)$, we have that $u(x_n, \pi) \geq u(z, \pi)$. By continuity of u , taking limits with respect to n of both sides, we get $u(x_0, \pi) \geq u(z, \pi)$, i.e., $x_0 \in \xi(\pi)$, as desired.

ξ is UHC.

From Proposition 555, it suffices to show that given an arbitrary closed set V in X , ${}^w\xi^{-1}(V) := \{\pi \in \Pi : \xi(\pi) \cap V \neq \emptyset\}$ is closed in Π . Consider an arbitrary sequence $(\pi_n)_{n \in \mathbb{N}}$ such that $\{\pi_n : n \in \mathbb{N}\} \subseteq {}^w\xi^{-1}(V)$ and such that $\pi_n \rightarrow \pi_0$. We have to show that $\pi_0 \in {}^w\xi^{-1}(V)$.

Take a sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$ such that for every n , $x_n \in \xi(\pi_n) \cap V \neq \emptyset$. Since $\xi(\pi_n) \subseteq \beta(\pi_n)$, it follows that $x_n \in \beta(\pi_n)$. We can now show the following

Claim. There exists a subsequence $(x_{n_k})_{n_k \in \mathbb{N}}$ of $(x_n)_{n \in \mathbb{N}}$ such that $x_{n_k} \rightarrow x_0$ and $x_0 \in \beta(\pi_0)$.

Proof of the Claim.

Since $\{\pi_n : n \in \mathbb{N}\} \cup \{\pi_0\}$ is a compact set (Show it), and since, by assumption, β is UHC and compact valued, from Proposition 559, $\beta(\{\pi_n : n \in \mathbb{N}\} \cup \{\pi_0\})$ is compact. Since $\{x_n\}_n \subseteq \beta(\{\pi_n\} \cup \{\pi_0\})$, there exists a subsequence $(x_{n_k})_{n_k \in \mathbb{N}}$ of $(x_n)_{n \in \mathbb{N}}$ which converges to some x_0 . Since β is compact valued, it is closed valued, too. Then, β is UHC and closed valued and from Proposition 550, β is closed. Since

$$\pi_{n_k} \rightarrow \pi_0, \quad x_{n_k} \in \beta(\pi_{n_k}), \quad x_{n_k} \rightarrow x_0,$$

the fact that β is closed implies that $x_0 \in \beta(\pi_0)$.

End of the Proof of the Claim.

Choose an arbitrary element z_0 such that $z_0 \in \beta(\pi_0)$. Since we assumed that $\pi_n \rightarrow \pi_0$ and since β is LHC, there exists a sequence $(z_n)_{n \in \mathbb{N}} \in X^\infty$ such that $z_n \in \beta(\pi_n)$ and $z_n \rightarrow z_0$.

Summarizing, and taking the subsequences of $(\pi_n)_{n \in \mathbb{N}}$ and $(z_n)_{n \in \mathbb{N}}$ corresponding to $(x_{n_k})_{n_k \in \mathbb{N}}$, we have for any n_k ,

$$\begin{aligned} \pi_{n_k} &\rightarrow \pi_0, \\ x_{n_k} &\rightarrow x_0, \quad x_{n_k} \in \xi(\pi_{n_k}), \quad x_0 \in \beta(\pi_0), \\ z_{n_k} &\rightarrow z_0, \quad z_{n_k} \in \beta(\pi_{n_k}), \quad z_0 \in \beta(\pi_0). \end{aligned}$$

Then for any n_k , we have that $u(x_{n_k}, \pi_{n_k}) \geq u(z_{n_k}, \pi_{n_k})$. Since u is continuous, taking limits, we get that $u(x_0, \pi_0) \geq u(z_0, \pi_0)$. Since the choice of z_0 in $\beta(\pi_0)$ was arbitrary, we have then $x_0 \in \xi(\pi_0)$.

Finally, since $(x_{n_k})_{n_k \in \mathbb{N}} \in V^\infty$, $x_{n_k} \rightarrow x_0$ and V is closed, $x_0 \in V$. Then $x_0 \in \xi(\pi_0) \cap V$ and $\pi_0 \in \{\pi \in \Pi : \xi(\pi) \cap V \neq \emptyset\} := {}^w\xi^{-1}(V)$, which was the desired result.

ξ is closed.

ξ is UHC and compact valued, and therefore closed valued. Then, from Proposition 550, it is closed, too.

v is a continuous function.

The basic idea of the proof is that v is a function and “it is equal to” the composition of UHC correspondences; therefore, it is a continuous function. A precise argument goes as follows.

Let the following correspondences be given:

$$\begin{aligned} (\xi, id) : \Pi \rightarrow\!\!\rightarrow X \times \Pi, \quad \pi &\mapsto \xi(\pi) \times \{\pi\}, \\ \beta : X \times \Pi \rightarrow\!\!\rightarrow \mathbb{R}, \quad (x, \pi) &\mapsto \{u(x, \pi)\}. \end{aligned}$$

Then, from Definition 557,

$$(\beta \circ (\xi, id))(\pi) = \cup_{(x, \pi) \in \xi(\pi) \times \{\pi\}} \{u(x, \pi)\}.$$

By definition of ξ ,

$$\forall \pi \in \Pi, \forall \bar{x} \in \xi(\pi), \quad \cup_{(x, \pi) \in \xi(\pi) \times \{\pi\}} \{u(x, \pi)\} = \{u(\bar{x}, \pi)\},$$

and

$$\forall \pi \in \Pi, \quad (\beta \circ (\xi, id))(\pi) = \{u(\bar{x}, \pi)\} = \{v(\pi)\}. \quad (11.3)$$

Now, (ξ, id) is UHC, and since u is a continuous function, β is UHC as well. From Proposition 558, $\beta \circ (\xi, id)$ is UHC and, from 11.3, v is a continuous function.

■

A sometimes more useful version of the Maximum Theorem is one which does not use the fact that β is UHC.

Theorem 563 (*Maximum Theorem*) Consider the correspondence $\beta : \Pi \rightarrow\!\!\rightarrow X$ and the function $u : X \times \Pi \rightarrow \mathbb{R}$ defined in Theorem 562 and Π, X Euclidean spaces.

Assume that

β is non-empty valued, compact valued, convex valued, closed and LHC.

u is continuous.

Then

1. ξ is a non-empty valued, compact valued, closed and UHC correspondence;
2. v is a continuous function.

Proof.

The desired result follows from next Proposition.

■

Proposition 564 Consider the correspondence $\beta : \Pi \rightarrow\!\!\rightarrow X$, with Π and X Euclidean spaces.

Assume that β is non-empty valued, compact valued, convex valued, closed and LHC. Then β is UHC.

Proof.

See Hildenbrand (1974) Lemma 1 page 33. The proof requires also Theorem 1 in Hildenbrand (1974).

■

The following result allows to substitute the requirement “ β is LHC” with the easier to check requirement “ $Cl\beta$ is LHC”.

Proposition 565 Consider the correspondence $\varphi : \Pi \rightarrow\!\!\rightarrow X$. φ is LHC $\Leftrightarrow Cl\varphi$ is LHC.

Proof.

Preliminary Claim.

V open set, $Cl\phi(\pi) \cap V \neq \emptyset \Rightarrow \varphi(\pi) \cap V \neq \emptyset$.

Proof of the Preliminary Claim.

Take $z \in Cl\phi(\pi) \cap V \neq \emptyset$. Since V is open, $\exists \varepsilon > 0$ such that $B(z, \varepsilon) \subseteq V$. Since $z \in Cl\phi(\pi)$, $\exists \{z_n\} \subseteq \varphi(\pi)$ such that $z_n \rightarrow z$. But then $\exists n_\varepsilon$ such that $n > n_\varepsilon \Rightarrow z_n \in B(z, \varepsilon) \subseteq V$. But $z_n \in V$ and $z_n \in \varphi(\pi)$ implies that $\varphi(\pi) \cap V \neq \emptyset$.

End of the Proof of the Preliminary Claim.

[\Rightarrow]

Take an open set V such that $Cl\phi(\pi) \cap V \neq \emptyset$. We want to show that there exists an open set U^* such that $\pi \in U^*$ and $\forall \xi \in U^*$, $Cl\phi(\xi) \cap V \neq \emptyset$. From the preliminary remark, it must be the case that $\varphi(\pi) \cap V \neq \emptyset$. Then, since φ is LHC, there exists an open set U such that $\pi \in U$ and $\forall \xi \in U$, $\varphi(\xi) \cap V \neq \emptyset$. Since $Cl\phi(\pi) \supseteq \varphi(\pi)$, we also have $Cl\phi(\pi) \cap V \neq \emptyset$. Choosing $U^* = U$, we are done.

[\Leftarrow]

Since $\varphi(\pi) \cap V \neq \emptyset$, then $Cl\phi(\pi) \cap V \neq \emptyset$, and, by assumption, \exists open set U' such that $\pi \in U'$ and $\forall \xi \in U'$, $Cl\phi(\xi) \cap V \neq \emptyset$. Then, from the preliminary remark, it must be the case that $\varphi(\pi) \cap V \neq \emptyset$.

■

Remark 566 *In some economic models, a convenient strategy to show that a correspondence β is LHC is the following one. Introduce a correspondence $\hat{\beta}$; show that $\hat{\beta}$ is LHC; show that $Cl \hat{\beta} = \beta$. Then from the above Proposition 565, the desired result follows - see, for example, point 5 the proof of Proposition 579 below.*

11.3 Fixed point theorems

A thorough analysis of the many versions of fixed point theorems existing in the literature is outside the scope of this notes. Below, we present a useful relatively general version of fixed point theorems both in the case of functions and correspondences.

Theorem 567 (*The Brouwer Fixed Point Theorem*)

For any $n \in \mathbb{N}$, let S be a nonempty, compact, convex subset of \mathbb{R}^n . If $f : S \rightarrow S$ is a continuous function, then $\exists x \in S$ such that $f(x) = x$.

Proof. For a (not self-contained) proof, see Ok (2007), page 279. ■

Just to try to avoid having a Section without a proof, let's show the following extremely simple version of that theorem.

Proposition 568 *If $f : [0, 1] \rightarrow [0, 1]$ is a continuous function, then $\exists x \in [0, 1]$ such that $f(x) = x$.*

Proof. If $f(0) = 0$ or $f(1) = 1$, the result is true. Then suppose otherwise, i.e., $f(0) \neq 0$ and $f(1) \neq 1$, i.e., since the domain of f is $[0, 1]$, suppose that $f(0) > 0$ and $f(1) < 1$. Define

$$g : [0, 1] \rightarrow \mathbb{R}, \quad : x \mapsto x - f(x).$$

Clearly, g is continuous, $g(0) = -f(0) < 0$ and $g(1) = 1 - f(1) > 0$. Then, from the intermediate value for continuous functions, $\exists x \in [0, 1]$ such that $g(x) = x - f(x) = 0$, i.e., $x = f(x)$, as desired. ■

Theorem 569 (*Kakutani's Fixed Point Theorem*)

For any $n \in \mathbb{N}$, let S be a nonempty, compact, convex subset of \mathbb{R}^n . If $\varphi : S \rightarrow S$ is a convex valued, closed correspondence, then $\exists x \in S$ such that $\varphi(x) \ni x$.

Proof. For a proof, see Ok (2007), page 331. ■

11.4 Application of the maximum theorem to the consumer problem

Definition 570 (*Mas Colell (1996), page 17*) *Commodities are goods and services available for purchases in the market.*

We assume the number of commodities is finite and equal to C . Commodities are indexed by superscript $c = 1, \dots, C$.

Definition 571 *A commodity vector is an element of the commodity space \mathbb{R}^C .*

Definition 572 (*almost Mas Colell (1996), page 18*) *A consumption set is a subset of the commodity space \mathbb{R}^C . It is denoted by X . Its elements are the vector of commodities the individual can conceivably consume given the physical or institutional constraints imposed by the environment.*

Example 573 *See Mas colell pages 18, 19.*

Common assumptions on X are that it is convex, bounded below and unbounded. Unless otherwise stated, we make the following stronger

Assumption 1 $X = \mathbb{R}_+^C := \{x \in \mathbb{R}^C : x \geq 0\}$.

Definition 574 $p \in \mathbb{R}^C$ is the vector of commodity prices.

Households' choices are limited also by an economic constraint: they cannot buy goods whose value is bigger than their wealth, i.e., it must be the case that $px \leq w$, where w is household's wealth.

Remark 575 w can take different specifications. For example, we can have $w = pe$, where $e \in \mathbb{R}^C$ is the vector of goods owned by the household, i.e., her endowments.

Assumption 2 All commodities are traded in markets at publicly observable prices, expressed in monetary unit terms.

Assumption 3 All commodities are assumed to be strictly goods (and not "bad"), i.e., $p \in \mathbb{R}_{++}^C$.

Assumption 4 Households behave as if they cannot influence prices.

Definition 576 *The budget set is*

$$\beta(p, w) := \{x \in \mathbb{R}_+^C : px \leq w\}.$$

With some abuse of notation we define the budget correspondence as

$$\beta : \mathbb{R}_{++}^C \times \mathbb{R}_{++} \rightarrow \mathbb{R}^C, \beta(p, w) = \{x \in \mathbb{R}_+^C : px \leq w\}.$$

Definition 577 *The utility function is*

$$u : X \rightarrow \mathbb{R}, \quad x \mapsto u(x)$$

Definition 578 *The Utility Maximization Problem (UMP) is*

$$\max_{x \in \mathbb{R}_+^C} u(x) \quad \text{s.t.} \quad px \leq w, \text{ or } x \in \beta(p, w).$$

$\xi : \mathbb{R}_{++}^C \times \mathbb{R}_{++} \rightarrow \mathbb{R}^C, \xi(p, w) = \arg \max(UMP)$ *is the demand correspondence.*

Theorem 579 ξ is a non-empty valued, compact valued, closed and UHC correspondence and

$$v : \mathbb{R}_{++}^C \times \mathbb{R}_{++} \rightarrow \mathbb{R}, v : (p, w) \mapsto \max(UMP),$$

i.e., the indirect utility function, is a continuous function.

Proof.

As an application of the (second version of) the Maximum Theorem, i.e., Theorem 563, we have to show that β is non-empty valued, compact valued, convex valued, closed and LHC.

1. β is non-empty valued.

$$x = \left(\frac{w}{Cp^c} \right)_{c=1}^C \in \beta(p, w) \text{ (or, simpler, } 0 \in \beta(p, w)).$$

2. β is compact valued.

$\beta(p, w)$ is closed because is the intersection of the inverse image of closed sets via continuous functions.

$\beta(p, w)$ is bounded below by zero.

$\beta(p, w)$ is bounded above because for every c , $x^c \leq \frac{w - \sum_{c' \neq c} p^{c'} x^{c'}}{p^c} \leq \frac{w}{p^c}$, where the first inequality comes from the fact that $px \leq w$, and the second inequality from the fact that $p \in \mathbb{R}_{++}^C$ and $x \in \mathbb{R}_+^C$.

3. β is convex valued.

To see that, simply, observe that $(1 - \lambda)px' + \lambda px'' \leq (1 - \lambda)w + \lambda w = w$.

4. β is closed.

We want to show that for every sequence $\{(p_n, w_n)\}_n \subseteq \mathbb{R}_{++}^C \times \mathbb{R}_{++}$ such that

$$(p_n, w_n) \rightarrow (p, w), \quad x_n \in \beta(p_n, w_n), \quad x_n \rightarrow x,$$

it is the case that $x \in \beta(p, w)$.

Since $x_n \in \beta(p_n, w_n)$, we have that $p_n x_n \leq w_n$ and $x_n \geq 0$. Taking limits of both sides of both inequalities, we get $px \leq w$ and $x \geq 0$, i.e., $x \in \beta(p, w)$.

5. β is LHC.

We proceed as follows: a. $\text{Int } \beta$ is LHC; b. $\text{Cl } \text{Int } \beta = \beta$. Then, from Proposition 565 the result follows.

a. Observe that $\text{Int } \beta(p, w) := \{x \in \mathbb{R}_+^C : x \gg 0 \text{ and } px < w\}$ and that $\text{Int } \beta(p, w) \neq \emptyset$, since $x = \left(\frac{w}{2Cp^c} \right)_{c=1}^C \in \text{Int } \beta(p, w)$. We want to show that the following is true.

For every sequence $(p_n, w_n)_n \in (\mathbb{R}_{++}^C \times \mathbb{R}_{++})^\infty$ such that $(p_n, w_n) \rightarrow (p, w)$ and for any $x \in \text{Int } \beta(p, w)$,

there exists a sequence $\{x_n\}_n \subseteq \mathbb{R}_+^C$ such that $\forall n, x_n \in \text{Int } \beta(p_n, w_n)$ and $x_n \rightarrow x$.

$p_n x_n - w_n \rightarrow px - w < 0$ (where the strict inequality follows from the fact that $x \in \text{Int } \beta(p, w)$).

Then, $\exists N$ such that $n \geq N \Rightarrow p_n x_n - w_n < 0$.

For $n \leq N$, choose an arbitrary $x_n \in \text{Int } \beta(p_n, w_n) \neq \emptyset$. Since $p_n x_n - w_n < 0$, for every $n > N$, there exists $\varepsilon_n > 0$ such that $z \in B(x, \varepsilon_n) \Rightarrow p_n z - w_n < 0$.

For any $n > N$, choose $x_n = x + \frac{1}{\sqrt{C}} \min \left\{ \frac{\varepsilon_n}{2}, \frac{1}{n} \right\} \cdot \mathbf{1}$. Then,

$$d(x, x_n) = \left(C \left(\frac{1}{\sqrt{C}} \min \left\{ \frac{\varepsilon_n}{2}, \frac{1}{n} \right\} \right)^2 \right)^{\frac{1}{2}} = \min \left\{ \frac{\varepsilon_n}{2}, \frac{1}{n} \right\} < \varepsilon_n,$$

i.e., $x_n \in B(x, \varepsilon_n)$ and therefore

$$p_n x_n - w_n < 0 \quad (1).$$

Since $x_n \gg x$, we also have

$$x_n \gg 0 \quad (2).$$

(1) and (2) imply that $x_n \in \text{Int } \beta(p_n, w_n)$. Moreover, since $x_n \gg x$, we have $0 \leq \lim_{n \rightarrow +\infty} (x_n - x) = \lim_{n \rightarrow +\infty} \frac{1}{\sqrt{C}} \cdot \min \left\{ \frac{\varepsilon_n}{2}, \frac{1}{n} \right\} \cdot \mathbf{1} \leq \lim_{n \rightarrow +\infty} \frac{1}{n} \frac{1}{\sqrt{C}} \cdot \mathbf{1} = 0$, i.e., $\lim_{n \rightarrow +\infty} x_n = x$.⁶

b.

It follows from the fact that the budget correspondence is the intersection of the inverse images of half spaces via continuous functions.

2.

It follows from Proposition 580, part (4), and the Maximum Theorem.

■

⁶Or simply

$$0 \leq \lim_{n \rightarrow \infty} d(x, x_n) \leq \lim_{n \rightarrow \infty} \frac{1}{n} = 0.$$

Proposition 580 For every $(p, w) \in \mathbb{R}_{++}^C \times \mathbb{R}_{++}$,

- (1) $\forall \alpha \in \mathbb{R}_{++}, \xi(\alpha p, \alpha w) = \xi(p, w)$;
- (2) if u is LNS, $\forall x \in \mathbb{R}_+^C, x \in \xi(p, w) \Rightarrow px = w$;
- (3) if u is quasi-concave, ξ is convex valued;
- (4) if u is strictly quasi-concave, ξ is single valued, i.e., it is a function.

Proof.

(1)

It simply follows from the fact that $\forall \alpha \in \mathbb{R}_{++}, \beta(\alpha p, \alpha w) = \beta(p, w)$.

(2)

Suppose otherwise, then $\exists x' \in \mathbb{R}_+^C$ such that $x' \in \xi(p, w)$ and $px' < w$. Therefore, $\exists \varepsilon' > 0$ such that $B(x', \varepsilon') \subseteq \beta(p, w)$ (take $\varepsilon' = d(x', H(p, w))$). Then, from the fact that u is LNS, there exists x^* such that $x^* \in B(x', \varepsilon') \subseteq \beta(p, w)$ and $u(x^*) > u(x')$, i.e., $x' \notin \xi(p, w)$, a contradiction.

(3)

Assume there exist x', x'' such that $x', x'' \in \xi(p, w)$. We want to show that $\forall \lambda \in [0, 1], x^\lambda := (1 - \lambda)x' + \lambda x'' \in \xi(p, w)$. Observe that $u(x') = u(x'') := u^*$. From the quasi-concavity of u , we have $u(x^\lambda) \geq u^*$. We are therefore left with showing that $x^\lambda \in \beta(p, w)$, i.e., β is convex valued. To see that, simply, observe that $px^\lambda = (1 - \lambda)px' + \lambda px'' \leq (1 - \lambda)w + \lambda w = w$.

(4) Assume otherwise. Following exactly the same argument as above we have $x', x'' \in \xi(p, w)$, and $px^\lambda \leq w$. Since u is strictly quasi concave, we also have that $u(x^\lambda) > u(x') = u(x'') := u^*$, which contradicts the fact that $x', x'' \in \xi(p, w)$.

■

Proposition 581 If u is a continuous LNS utility function, then the indirect utility function has the following properties.

For every $(p, w) \in \mathbb{R}_{++}^C \times \mathbb{R}_{++}$,

- (1) $\forall \alpha \in \mathbb{R}_{++}, v(\alpha p, \alpha w) = v(p, w)$;
- (2) Strictly increasing in w and for every c , non increasing in p^c ;
- (3) for every $\bar{v} \in \mathbb{R}, \{(p, w) : v(p, w) \leq \bar{v}\}$ is convex.
- (4) continuous.

Proof.

(1) It follows from Proposition 580 (2).

(2)

If w increases, say by Δw , then, from Proposition 580 (2), $px(p, w) < w + \Delta w$. Define $x(p, w) := x'$. Then, $\exists \varepsilon' > 0$ such that $B(x', \varepsilon') \subseteq \beta(p, w + \Delta w)$ (take $\varepsilon' = d(x', H(p, w + \Delta w))$). Then, from the fact that u is LNS, there exists x^* such that $x^* \in B(x', \varepsilon') \subseteq \beta(p, w + \Delta w)$ and $u(x^*) > u(x')$. The result follows observing that $v(p, w + \Delta w) \geq u(x^*)$.

Similar proof applies to the case of a decrease in p . Assume $\Delta p^{c'} < 0$. Define $\Delta := (\Delta^c)_{c=1}^C \in \mathbb{R}^C$ with $\Delta^c = 0$ iff $c \neq c'$ and $\Delta^{c'} = \Delta p^{c'}$. Then,

$$\begin{aligned} px(p, w) = w &\Rightarrow (p + \Delta)x(p, w) = px(p, w) + \Delta p^{c'} x^{c'}(p, w) = \\ &= w + \Delta p^{c'} x^{c'}(p, w) \leq w. \end{aligned}$$

The remaining part of the proof is the same as in the case of an increase of w .

(3) Take $(p', w'), (p'', w'') \in \{(p, w) : v(p, w) \leq \bar{v}\} := S(\bar{v})$. We want to show that $\forall \lambda \in [0, 1], (p^\lambda, w^\lambda) := (1 - \lambda)(p', w') + \lambda(p'', w'') \in S(\bar{v})$, i.e., $x \in \xi(p^\lambda, w^\lambda) \Rightarrow u(x) > \bar{v}$.

$$x \in \xi(p^\lambda, w^\lambda) \Rightarrow p^\lambda x \leq w^\lambda \Leftrightarrow (1 - \lambda)p' + \lambda p'' \leq (1 - \lambda)w' + \lambda w''.$$

Then, either $p'x \leq w'$ or $p''x \leq w''$. If $p'x \leq w'$, then $u(x) \leq v(p', w') \leq \bar{v}$. Similarly, if $p''x \leq w''$.

(4)

It was proved in Theorem 579.

■

Part III

Differential calculus in Euclidean spaces

Chapter 12

Partial derivatives and directional derivatives

12.1 Partial Derivatives

The¹ concept of partial derivative is not that different from the concept of “standard” derivative of a function from \mathbb{R} to \mathbb{R} , in fact we are going to see that partial derivatives of a function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ are just standard derivatives of a naturally associated function from \mathbb{R} to \mathbb{R} .

Recall that for any $k \in \{1, \dots, n\}$, $e_n^k = (0, \dots, 1, \dots, 0)$ is the k -th vector in the canonical basis of \mathbb{R}^n .

Definition 582 Let a set $S \subseteq \mathbb{R}^n$, a point $x_0 = (x_{0k})_{k=1}^n \in \text{Int } S$ and a function $f : S \rightarrow \mathbb{R}$ be given. If the following limit exists and it is finite

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h e_n^k) - f(x_0)}{h} = \lim_{x_k \rightarrow x_{0k}} \frac{f(x_0 + (x_k - x_{0k}) e_n^k) - f(x_0)}{x_k - x_{0k}} \quad (12.1)$$

then it is called the partial derivative of f with respect to the k -th coordinate computed in x_0 and it is denoted by any of the following symbols

$$D_{x_k} f(x_0), \quad D_k f(x_0), \quad \frac{\partial f}{\partial x_k}(x_0), \quad \frac{\partial f(x)}{\partial x_k} \Big|_{x=x_0}.$$

Remark 583 As said above, partial derivatives are not really a new concept. We are just treating f as a function of one variable at the time, keeping the other variables fixed. In other words, for simplicity taking $S = \mathbb{R}^n$ and using the notation of the above definition, we can define

$$g_k : \mathbb{R} \rightarrow \mathbb{R}, \quad g_k(x_k) = f(x_0 + (x_k - x_{0k}) e_n^k)$$

a function of only one variable, and, by definition of g_k ,

$$\begin{aligned} g_k'(x_{0k}) &= \lim_{x_k \rightarrow x_{0k}} \frac{g_k(x_k) - g_k(x_{0k})}{x_k - x_{0k}} = \\ &= \lim_{h \rightarrow 0} \frac{g_k(x_{0k} + h) - g_k(x_{0k})}{h} = \lim_{h \rightarrow 0} \frac{f(x_0 + h e_n^k) - f(x_0)}{h} = D_{x_k} f(x_0). \end{aligned} \quad (12.2)$$

Example 584 Given $f : \mathbb{R}^3 \rightarrow \mathbb{R}$,

$$f(x_1, x_2, x_3) = e^{xy} \cos x + \sin yz$$

we have

$$\begin{pmatrix} D_{x_1} f(x) \\ D_{x_2} f(x) \\ D_{x_3} f(x) \end{pmatrix} = \begin{pmatrix} -(\sin x) e^{xy} + y (\cos x) e^{xy} \\ z \cos yz + x (\cos x) e^{xy} \\ y \cos yz \end{pmatrix}$$

¹In this Part, I follow closely Section 5.14 and chapters 12 and 13 in Apostol (1974).

Remark 585 Loosely speaking, we can give the following geometrical interpretation of partial derivatives. Given $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ admitting partial derivatives, $\frac{\partial f(x_0)}{\partial x_1}$ is the slope of the graph of the function obtained cutting the graph of f with a plane which is
 orthogonal to the $x_1 - x_2$ plane, and
 going through the line parallel to the x_1 axis and passing through the point x_0 , line to which we have given the same orientation as the x_1 axis.
 picture to be added.

Definition 586 Given an open subset S in \mathbb{R}^n and a function $f : S \rightarrow \mathbb{R}$, if $\forall k \in \{1, \dots, n\}$, the limit in (12.1) exists, we call the gradient of f in x_0 the following vector


$$(D_k f(x_0))_{k=1}^n$$

and we denote it by

$$Df(x_0)$$

Remark 587 The existence of the gradient for f in x_0 does not imply continuity of the function in x_0 , as the following example shows.

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f(x_1, x_2) = \begin{cases} x_1 + x_2 & \text{if } \begin{array}{l} \text{either } x_1 = 0 \text{ or } x_2 = 0 \\ \text{i.e., } (x_1, x_2) \in (\{0\} \times \mathbb{R}) \cup (\mathbb{R} \times \{0\}) \end{array} \\ 1 & \text{otherwise} \end{cases}$$



$$D_1 f(0) = \lim_{x_1 \rightarrow 0} \frac{f(x_1, 0) - f(0, 0)}{x_1 - 0} = \lim_{x_1 \rightarrow 0} \frac{x_1}{x_1 - 0} = 1$$

and similarly

$$D_2 f(0) = 1.$$

f is not continuous in 0: we want to show that $\exists \varepsilon > 0$ such that $\forall \delta > 0$ there exists $(x_1, x_2) \in \mathbb{R}^2$ such that $(x_1, x_2) \in B(0, \delta)$ and $|f(x_1, x_2) - f(0, 0)| \geq \varepsilon$. Take $\varepsilon = \frac{1}{2}$ and any $(x_1, x_2) \in B(0, \delta)$ such that $x_1 \neq 0$ and $x_2 \neq 0$. Then $|f(x_1, x_2) - f(0, 0)| = 1 > \varepsilon$.

12.2 Directional Derivatives

A first generalization of the concept of partial derivative of a function is presented in Definition 589 below.

Definition 588 Given

$$f : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad x \mapsto f(x),$$

$\forall i \in \{1, \dots, m\}$, the function

$$f_i : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m, \quad x \mapsto i\text{-th component of } f(x).$$

is called the i -th component function of f .

Therefore,

$$\forall x \in S, \quad f(x) = (f_i(x))_{i=1}^m. \quad (12.3)$$

Definition 589 Given $m, n \in \mathbb{N}$, a set $S \subseteq \mathbb{R}^n$, $x_0 \in \text{Int } S$, $u \in \mathbb{R}^n$, $h \in \mathbb{R}$ such that $x_0 + hu \in S$, $f : S \rightarrow \mathbb{R}^m$, we call the directional derivative of f at x_0 in the direction u , denoted by the symbol

$$f'(x_0; u),$$

the limit

$$\lim_{h \rightarrow 0} \frac{f(x_0 + hu) - f(x_0)}{h} \quad (12.4)$$

if it exists and it is finite.

Remark 590 Assume that the limit in (12.4) exists and it is finite. Then, from (12.3) and using Proposition 421,

$$f'(x_0; u) = \lim_{h \rightarrow 0} \frac{f(x_0 + hu) - f(x_0)}{h} = \left(\lim_{h \rightarrow 0} \frac{f_i(x_0 + hu) - f_i(x_0)}{h} \right)_{i=1}^m = (f'_i(x_0; u))_{i=1}^m.$$

If $u = e_n^j$, the j -th element of the canonical basis in \mathbb{R}^n , we then have

$$f'(x_0; e_n^j) = \left(\lim_{h \rightarrow 0} \frac{f_i(x_0 + he_n^j) - f_i(x_0)}{h} \right)_{i=1}^m = (f'_i(x_0; e_n^j))_{i=1}^m \stackrel{(*)}{=} (D_{x_j} f_i(x_0))_{i=1}^m := D_{x_j} f(x_0) \quad (12.5)$$

where equality $(*)$ follows from (12.2).

We can then construct a matrix whose n columns are the above vectors, a matrix which involves all partial derivative of all component functions of f . That matrix is formally defined below.

Definition 591 Assume that $f = (f_i)_{i=1}^m : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ admits all partial derivatives in x_0 . The Jacobian matrix of f at x_0 is denoted by $Df(x_0)$ and is the following $m \times n$ matrix:

$$\begin{aligned} & \begin{bmatrix} D_{x_1} f_1(x_0) & \dots & D_{x_j} f_1(x_0) & \dots & D_{x_n} f_1(x_0) \\ \vdots & & \vdots & & \vdots \\ D_{x_1} f_i(x_0) & \dots & D_{x_j} f_i(x_0) & \dots & D_{x_n} f_i(x_0) \\ \vdots & & \vdots & & \vdots \\ D_{x_1} f_m(x_0) & \dots & D_{x_j} f_m(x_0) & \dots & D_{x_n} f_m(x_0) \end{bmatrix} = \\ &= \begin{bmatrix} D_{x_1} f(x_0) & \dots & D_{x_j} f(x_0) & \dots & D_{x_n} f(x_0) \end{bmatrix} = \\ &= \begin{bmatrix} f'(x_0; e_n^1) & \dots & f'(x_0; e_n^j) & \dots & f'(x_0; e_n^n) \end{bmatrix}. \end{aligned}$$

Remark 592 How to easily write the Jacobian matrix of a function.

To compute the Jacobian of f is convenient to construct a table as follows.

1. In the first column, write the m vector component functions $f_1, \dots, f_i, \dots, f_m$ of f .
2. In the first row, write the subvectors $x_1, \dots, x_j, \dots, x_n$ of x .
3. For each i and j , write the partial Jacobian matrix $D_{x_j} f_i(x)$ in the entry at the intersection of the i -th row and j -th column.

We then obtain the following table,

	x_1	\dots	x_j	\dots	x_n
f_1	$D_{x_1} f_1(x)$		$D_{x_j} f_1(x)$		$D_{x_n} f_1(x)$
\dots					
f_i	$D_{x_1} f_i(x)$		$D_{x_j} f_i(x)$		$D_{x_n} f_i(x)$
\dots					
f_m	$D_{x_1} f_m(x)$		$D_{x_j} f_m(x)$		$D_{x_n} f_m(x)$

where the Jacobian matrix is the part of the table between square brackets.

Example 593 Given $f : \mathbb{R}^4 \rightarrow \mathbb{R}^5$,

$$f(x, y, z, t) = \begin{pmatrix} \frac{xy}{x^2+1} \\ \frac{x+yz}{e^x} \\ \frac{xyz}{e^t} \\ x+y+z+t \\ x^2+t^2 \end{pmatrix}$$

its Jacobian matrix is

$$\begin{bmatrix} \frac{y}{x^2+1} - 2x^2 \frac{y}{(x^2+1)^2} & \frac{x}{x^2+1} & 0 & 0 \\ \frac{1}{e^x} - \frac{1}{e^x}(x+yz) & \frac{z}{e^x} & \frac{y}{e^x} & 0 \\ ty \frac{z}{e^t} & tx \frac{z}{e^t} & tx \frac{y}{e^t} & xy \frac{z}{e^t} - txy \frac{z}{e^t} \\ 1 & 1 & 1 & 1 \\ 2x & 0 & 0 & 2t \end{bmatrix}_{5 \times 4}$$

Remark 594 From Remark 590,

$$\forall u \in \mathbb{R}^n, f'(x_0; u) \text{ exists} \Rightarrow Df(x_0) \text{ exists} \quad (12.6)$$

On the other hand, the opposite implication does not hold true. Consider the example in Remark 587. There, we have seen that

$$D_x f(0) = D_y f(0) = 1$$

But if $u = (u_1, u_2)$ with $u_1 \neq 0$ and $u_2 \neq 0$, we have

$$\lim_{h \rightarrow 0} \frac{f(0+hu) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{1-0}{h} = \infty$$

Remark 595 Again loosely speaking, we can give the following geometrical interpretation of directional derivatives. Take $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ admitting directional derivatives. $f(x_0; u)$ with $\|u\| = 1$ is the slope the graph of the function obtained cutting the graph of f with a plane which is

orthogonal to the $x_1 - x_2$ plane, and

going through the line going through the points x_0 and $x_0 + u$, line to which we have given the same orientation as u .

picture to be added.

Example 596 Take

$$f : \mathbb{R}^n \rightarrow \mathbb{R}, \quad x \mapsto x \cdot x = \|x\|^2$$

Then, the existence of $f'(x_0; u)$ can be checked computing the following limit.

$$\begin{aligned} \lim_{h \rightarrow 0} \frac{f(x_0+hu) - f(x_0)}{h} &= \lim_{h \rightarrow 0} \frac{(x_0+hu)(x_0+hu) - x_0x_0}{h} = \\ &= \lim_{h \rightarrow 0} \frac{x_0x_0 + hx_0u + hu x_0 + h^2uu - x_0x_0}{h} = \\ &= \lim_{h \rightarrow 0} \frac{2hx_0u + h^2uu}{h} = \lim_{h \rightarrow 0} 2x_0u + hu u = 2x_0u \end{aligned}$$

Exercise 597 Verify that

$$f'(x_0; -u) = -f'(x_0; u).$$

Solution:


$$\begin{aligned} f'(x_0; -u) &= \lim_{h \rightarrow 0} \frac{f(x_0 + h(-u)) - f(x_0)}{h} = \lim_{h \rightarrow 0} \frac{f(x_0 - hu) - f(x_0)}{h} = \\ &= -\lim_{k \rightarrow 0} \frac{f(x_0 + (-h)u) - f(x_0)}{-h} \stackrel{k := -h}{=} -\lim_{k \rightarrow 0} \frac{f(x_0 + ku) - f(x_0)}{k} = -f'(x_0; u). \end{aligned}$$

Remark 598 *It is **not** the case that*

$$\forall u \in \mathbb{R}^n, f'(x_0; u) \text{ exists} \Rightarrow f \text{ is continuous in } x_0 \quad (12.7)$$

as the following example shows. Consider

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f(x, y) = \begin{cases} \frac{xy^2}{x^2 + y^4} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0, \text{ i.e., } (x, y) \in \{0\} \times \mathbb{R} \end{cases}$$



Let's compute $f'(0; u)$. If $u_1 \neq 0$.

$$\lim_{h \rightarrow 0} \frac{f(0 + hu) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{hu_1 \cdot h^2 u_2^2}{(h^2 u_1^2 + h^4 u_2^4) h} = \lim_{h \rightarrow 0} \frac{u_1 \cdot u_2^2}{u_1^2 + h^2 u_2^4} = \frac{u_2^2}{u_1}$$

If $u_1 = 0$, we have

$$\lim_{h \rightarrow 0} \frac{f(0 + hu) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{f(0, hu_2) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{0}{h} = 0$$

On the other hand, if $x = y^2$ and $x, y \neq 0$, i.e., along the graph of the parabola $x = y^2$ except the origin, we have

$$f(x, y) = f(y^2, y) = \frac{y^4}{y^4 + y^4} = \frac{1}{2}$$

while

$$f(0, 0) = 0.$$

Roughly speaking, the existence of partial derivatives in a given point in all directions implies “continuity along straight lines” through that point; it does not imply “continuity along all possible curves through that point”, as in the case of the parabola in the picture above.

Remark 599 We are now left with two problems:

1. Is there a definition of derivative whose existence implies continuity?
2. Is there any “easy” way to compute the directional derivative?

Appendix (to be corrected)

There are other definitions of directional derivatives used in the literature.

Let the following objects be given: $m, n \in \mathbb{N}$, a set $S \subseteq \mathbb{R}^n$, $x_0 \in \text{Int } S$, $u \in \mathbb{R}^n$, $h \in \mathbb{R}$ such that $x_0 + hu \in S$, $f : S \rightarrow \mathbb{R}^m$,

Definition 600 (our definition following Apostol (1974)) We call the directional derivative of f at x_0 in the direction u according to Apostol, denoted by the symbol

$$f'_A(x_0; u),$$

the limit

$$\lim_{h \rightarrow 0} \frac{f(x_0 + hu) - f(x_0)}{h} \quad (12.8)$$

if it exists and it is finite.

Definition 601 (Girsanov (1972)) We call the directional derivative of f at x_0 in the direction u according to Girsanov, denoted by the symbol

$$f'_G(x_0; u),$$

the limit

$$\lim_{h \rightarrow 0^+} \frac{f(x_0 + hu) - f(x_0)}{h} \quad (12.9)$$

if it exists and it is finite.

Definition 602 (Wikipedia) Take $u \in \mathbb{R}^n$ such that $\|u\| = 1$. We call the directional derivative of f at x_0 in the direction u according to Wikipedia, denoted by the symbol

$$f'_W(x_0; u),$$

the limit

$$\lim_{h \rightarrow 0^+} \frac{f(x_0 + hu) - f(x_0)}{h} \quad (12.10)$$

if it exists and it is finite.

Fact 1. For given $x_0 \in S, u \in \mathbb{R}^n$

$$A \Rightarrow G \Rightarrow W,$$

while the opposite implications do not hold true. In particular, to see way $A \not\Rightarrow G$, just take $f : \mathbb{R} \rightarrow \mathbb{R}$,

$$f(x) = \begin{cases} 0 & \text{if } x < 0 \\ 1 & \text{if } x \geq 0 \end{cases},$$

and observe that while the right derivative in 0 is

$$\lim_{h \rightarrow 0^+} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0^+} \frac{1 - 1}{h} = 0,$$

while the left derivative is

$$\lim_{h \rightarrow 0^-} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0^-} \frac{0 - 1}{h} = +\infty.$$

Fact 2. For given $x_0 \in S$,

$$f'_W(x, u) \text{ exists} \Rightarrow f'_G(x, v) \text{ exists for any } v = \alpha u \text{ and } \alpha \in \mathbb{R}_{++}.$$

Proof.

$$\begin{aligned} f'_G(x, v) &= \lim_{h \rightarrow 0^+} \frac{f(x_0 + hv) - f(x_0)}{h} = \alpha \lim_{h \rightarrow 0^+} \frac{f(x_0 + h\alpha u) - f(x_0)}{\alpha h} \stackrel{k=\alpha h > 0}{=} \\ &= \alpha \lim_{h \rightarrow 0^+} \frac{f(x_0 + ku) - f(x_0)}{k} = \alpha f'_W(x, u). \end{aligned}$$

Fact 3. Assume that $u \neq 0$ and $x_0 \in \mathbb{R}^n$. Then the following implications are true:

$$\forall u \in \mathbb{R}^n, f'_A(x, u) \text{ exists} \Leftrightarrow \forall u \in \mathbb{R}^n, f'_G(x, u) \text{ exists} \Leftrightarrow \forall u \in \mathbb{R}^n \text{ such that } \|u\| = 1, f'_W(x, u) \text{ exists.}$$

Proof.

From Fact 1, we are left with showing just two implications.

$G \Rightarrow A$.

We want to show that

$$\forall u \in \mathbb{R}^n, \lim_{h \rightarrow 0^+} \frac{f(x_0 + hu) - f(x_0)}{h} \in \mathbb{R} \Rightarrow \forall v \in \mathbb{R}^n, \lim_{h \rightarrow 0} \frac{f(x_0 + hv) - f(x_0)}{h} \in \mathbb{R}.$$

Therefore, it suffices to show that $l := \lim_{h \rightarrow 0^-} \frac{f(x_0 + hv) - f(x_0)}{h} \in \mathbb{R}$. Take $u = -v$. Then,

$$l = \lim_{h \rightarrow 0^-} \frac{f(x_0 - hu) - f(x_0)}{h} = - \lim_{h \rightarrow 0^-} \frac{f(x_0 - hu) - f(x_0)}{-h} \stackrel{k=-h}{=} - \lim_{k \rightarrow 0^+} \frac{f(x_0 + ku) - f(x_0)}{k} \in \mathbb{R}.$$

$W \Rightarrow G$.

The proof of this implication is basically the proof of Fact 2. We want to show that

$$\forall u \in \mathbb{R}^n \text{ such that } \|u\| = 1, \lim_{h \rightarrow 0^+} \frac{f(x_0 + hu) - f(x_0)}{h} \in \mathbb{R} \Rightarrow \forall v \in \mathbb{R}^n \setminus \{0\}, l := \lim_{h \rightarrow 0^+} \frac{f(x_0 + hv) - f(x_0)}{h} \in \mathbb{R}.$$

In fact,

$$l := \lim_{h \rightarrow 0^+} \frac{f\left(x_0 + h\|v\| \frac{v}{\|v\|}\right) - f(x_0)}{h} \in \mathbb{R},$$

simply because $\left\| \frac{v}{\|v\|} \right\| = 1$.

Remark 603 We can give the following geometrical interpretation of directional derivatives. First of all observe that from Proposition 606,

$$f'(x_0; u) := \lim_{h \rightarrow 0} \frac{f(x_0 + hu) - f(x_0)}{h} = df_{x_0}(u).$$

If $f : \mathbb{R} \rightarrow \mathbb{R}$, we then have

$$f'(x_0; u) = f'(x_0) \cdot u.$$

Therefore, if $u = 1$, we have

$$f'(x_0; u) = f'(x_0),$$

and if $u > 0$, we have

$$\text{sign } f'(x_0; u) = \text{sign } f'(x_0).$$

Take now $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ admitting directional derivatives. Then,

$$f'(x_0; u) = Df(x_0) \cdot u \quad \text{with } \|u\| = 1$$

is the slope the graph of the function obtained cutting the graph of f with a plane which is orthogonal to the $x_1 - x_2$ plane, and along the line going through the points x_0 and $x_0 + u$, in the direction from x_0 to $x_0 + u$.

Chapter 13

Differentiability

13.1 Total Derivative and Differentiability

If $f : \mathbb{R} \rightarrow \mathbb{R}$, we say that f is differentiable in x_0 , if the following limit exists and it is finite

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

and we write

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h}$$

or, in equivalent manner,

$$\lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0) - f'(x_0) \cdot h}{h} = 0$$

and

$$f(x_0 + h) - (f(x_0) + f'(x_0) \cdot h) = r(h)$$

where

$$\lim_{h \rightarrow 0} \frac{r(h)}{h} = 0,$$

or

$$f(x_0 + h) = f(x_0) + f'(x_0) \cdot h + r(h),$$

or using what said in Section 7.5, and more specifically using definition 7.8,

$$f(x_0 + h) = f(x_0) + l_{f'(x_0)}(h) + r(h)$$

where $l_{f'(x_0)} \in \mathcal{L}(\mathbb{R}, \mathbb{R})$

and $\lim_{h \rightarrow 0} \frac{r(h)}{h} = 0$

Definition 604 Given a set $S \subseteq \mathbb{R}^n$, $x_0 \in \text{Int } S$, $f : S \rightarrow \mathbb{R}^m$, we say that f is differentiable at x_0 if there exists

a linear function $df_{x_0} : \mathbb{R}^n \rightarrow \mathbb{R}^m$

such that for any $u \in \mathbb{R}^n, u \neq 0$, such that $x_0 + u \in S$,

$$\lim_{u \rightarrow 0} \frac{f(x_0 + u) - f(x_0) - df_{x_0}(u)}{\|u\|} = 0 \quad (13.1)$$

In that case, the linear function df_{x_0} is called the total derivative or the differential or simply the derivative of f at x_0 .

Remark 605 Obviously, given the condition of the previous Definition, we can say that f is differentiable at x_0 if there exists a linear function $df_{x_0} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that $\forall u \in \mathbb{R}^n$ such that $x_0 + u \in S$

$$f(x_0 + u) = f(x_0) + df_{x_0}(u) + r(u), \quad \text{with} \quad \lim_{u \rightarrow 0} \frac{r(u)}{\|u\|} = 0 \quad (13.2)$$

or

$$f(x_0 + u) = f(x_0) + df_{x_0}(u) + \|u\| \cdot E_{x_0}(u), \quad \text{with} \quad \lim_{u \rightarrow 0} E_{x_0}(u) = 0 \quad (13.3)$$

The above equations are called the first-order Taylor formula (of f at x_0 in the direction u). Condition (13.3) is the most convenient one to use in many instances.

Proposition 606 Assume that $f : S \rightarrow \mathbb{R}^m$ is differentiable at x_0 , then

$$\forall u \in \mathbb{R}^n, \quad f'(x_0; u) = df_{x_0}(u).$$

Proof.

$$\begin{aligned} f'(x_0; u) &:= \lim_{h \rightarrow 0} \frac{f(x_0 + hu) - f(x_0)}{h} \stackrel{(1)}{=} \\ &= \lim_{h \rightarrow 0} \frac{f(x_0) + df_{x_0}(hu) + \|hu\| \cdot E_{x_0}(hu) - f(x_0)}{h} \stackrel{(2)}{=} \lim_{h \rightarrow 0} \frac{h df_{x_0}(u) + |h| \|u\| \cdot E_{x_0}(hu)}{h} \stackrel{(3)}{=} \\ &= \lim_{h \rightarrow 0} df_{x_0}(u) + \lim_{h \rightarrow 0} \text{sign}(h) \cdot \|u\| \cdot E_{x_0}(hu) \stackrel{(4)}{=} df_{x_0}(u) + \|u\| \lim_{hu \rightarrow 0} \text{sign}(h) \cdot E_{x_0}(hu) \stackrel{(5)}{=} df_{x_0}(u), \end{aligned}$$

where

(1) follows from (13.3) with hu in the place of u ,

(2) from the fact that df_{x_0} is linear and therefore (Exercise) continuous, and from a property of a norm,

(3) from the fact that $\frac{|h|}{h} = \text{sign}(h)$,¹

(4) from the fact that $h \rightarrow 0$ implies that $hu \rightarrow 0$,

(5) from the assumption that f is differentiable in x_0 . ■

Remark 607 The above Proposition implies that if the differential exists, then it is unique - from the fact that the limit is unique, if it exists.

Proposition 608 If $f : S \rightarrow \mathbb{R}^m$ is differentiable at x_0 , then f is continuous at x_0 .

Proof. We have to prove that

$$\lim_{u \rightarrow 0} f(x_0 + u) - f(x_0) = 0$$

i.e., from (13.3), it suffices to show that

$$\lim_{u \rightarrow 0} df_{x_0}(u) + \|u\| \cdot E_{x_0}(u) = df_{x_0}(0) + \lim_{u \rightarrow 0} \|u\| \cdot E_{x_0}(u) = 0$$

where the first equality follows from the fact that df_{x_0} is linear and therefore continuous, and the second equality from the fact again that df_{x_0} is linear, and therefore $df_x(0) = 0$, and from (13.2). ■

Remark 609 The above Proposition is the answer to Question 1 in Remark 599. We still do not have a answer to Question 2 and another question naturally arises at this point:

3. Is there an “easy” way of checking differentiability?

¹ sign is the function defined as follows:

$$\text{sign} : \mathbb{R} \rightarrow \{-1, 0, 1\}, \quad x \mapsto \begin{cases} -1 & \text{if } x < 0 \\ 0 & \text{if } x = 0 \\ +1 & \text{if } x > 0. \end{cases}$$

13.2 Total Derivatives in terms of Partial Derivatives.

In Remark 611 below, we answer question 2 in Remark 599: Is there any “easy” way to compute the directional derivative?

Proposition 610 Assume that $f = (f_j)_{j=1}^m : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ is differentiable in x_0 . The matrix associated with df_{x_0} with respect to the canonical bases of \mathbb{R}^n and \mathbb{R}^m is the Jacobian matrix $Df(x_0)$, i.e., using the notation of Section 7.5,

$$[df_{x_0}] = Df(x_0),$$

i.e.,

$$\forall x \in \mathbb{R}^n, \quad df_{x_0}(x) = Df(x_0) \cdot x. \quad (13.4)$$

Proof. From (7.6) in Section 7.5

$$[df_{x_0}] = \begin{bmatrix} df_{x_0}(e_1^n) & \dots & df_{x_0}(e_i^n) & \dots & df_{x_0}(e_n^n) \end{bmatrix}_{m \times n}$$

From Proposition 606,

$$\forall i \in \{1, \dots, n\}, \quad df_{x_0}(e^i) = f'(x_0; e^i),$$

and from (12.5)

$$f'(x_0; e^i) = (D_{x_i} f_j(x_0))_{j=1}^m.$$

Then

$$[df_{x_0}] = \begin{bmatrix} (D_{x_1} f_j(x_0))_{j=1}^m & \dots & (D_{x_i} f_j(x_0))_{j=1}^m & \dots & (D_{x_n} f_j(x_0))_{j=1}^m \end{bmatrix}_{m \times n},$$

as desired. ■

Remark 611 From Proposition 606, part 1, and the above Proposition 610, we have that if f is differentiable in x_0 , then $\forall u \in \mathbb{R}^m$

$$\forall u \in \mathbb{R}^m, \quad f'(x_0; u) = Df(x_0) \cdot u.$$

Remark 612 From (13.4), we get

$$\|df_{x_0}(x)\| = \|[Df(x_0)]_{mn} x\| \stackrel{(1)}{\leq} \sum_{j=1}^m |Df_j(x_0) \cdot x| \stackrel{(2)}{\leq} \sum_{j=1}^m \|Df_j(x_0)\| \cdot \|x\|$$

where (1) follows from Remark 56, (2) from Cauchy-Schwarz inequality in (53). Therefore, defined $\alpha := \sum_{j=1}^m \|Df_j(x_0)\|$, we have that

$$\|df_{x_0}(x)\| \leq \alpha \cdot \|x\|$$

and

$$\lim_{x \rightarrow 0} \|df_{x_0}(x)\| = 0$$

Remark 613 We have seen that

$$\begin{array}{lll} f \text{ differentiable in } x_0 & \Rightarrow & f \text{ admits directional derivative in } x_0 \quad \Rightarrow \quad Df(x_0) \text{ exists} \\ \Downarrow & & \text{(not } \Downarrow) \\ f \text{ continuous in } x_0 & \Rightarrow & f \text{ continuous in } x_0 \quad \Rightarrow \quad f \text{ continuous in } x_0 \end{array}$$

Therefore

$$f \text{ differentiable in } x_0 \quad \Leftarrow \quad Df(x_0) \text{ exists}$$

and

$$f \text{ differentiable in } x_0 \quad \Leftarrow \quad f \text{ admits directional derivative in } x_0$$

We still do not have an answer to question 3 in Remark 609: Is there an easy way of checking differentiability? We will provide an answer in Proposition 641.

Chapter 14

Some Theorems

We first introduce some needed definitions.

Definition 614 Given an open $S \subseteq \mathbb{R}^n$,

$$f : S \rightarrow \mathbb{R}^m, \quad x := (x_j)_{j=1}^n \mapsto f(x) = (f_i(x))_{i=1}^m$$

$$I \subseteq \{1, \dots, m\} \quad \text{and} \quad J \subseteq \{1, \dots, n\},$$

the partial Jacobian of $(f_i)_{i \in I}$ with respect to $(x_j)_{j \in J}$ in $x_0 \in S$ is the following $(\#I) \times (\#J)$ submatrix of $Df(x_0)$

$$\left[\frac{\partial f_i(x_0)}{\partial x_j} \right]_{i \in I, j \in J},$$

and it is denoted by

$$D_{(x_j)_{j \in J}}(f_j)_{i \in I}(x_0)$$

Example 615 Take:

S an open subset of \mathbb{R}^{n_1} , with generic element $x' = (x_j)_{j=1}^{n_1}$,

T an open subset of \mathbb{R}^{n_2} , with generic element $x'' = (x_k)_{k=1}^{n_2}$ and

$$f : S \times T \rightarrow \mathbb{R}^m, \quad (x', x'') \mapsto f(x', x'')$$

Then, defined $n = n_1 + n_2$, we have

$$D_{x'} f(x_0) = \begin{bmatrix} D_{x_1} f_1(x_0) & \dots & D_{x_{n_1}} f_1(x_0) \\ \dots & \dots & \dots \\ D_{x_1} f_i(x_0) & \dots & D_{x_{n_1}} f_i(x_0) \\ \dots & \dots & \dots \\ D_{x_1} f_m(x_0) & \dots & D_{x_{n_1}} f_m(x_0) \end{bmatrix}_{m \times n_1}$$

and, similarly,

$$D_{x''} f(x_0) := \begin{bmatrix} D_{x_{n_1+1}} f_1(x_0) & \dots & D_{x_n} f_1(x_0) \\ \dots & \dots & \dots \\ D_{x_{n_1+1}} f_i(x_0) & \dots & D_{x_n} f_i(x_0) \\ \dots & \dots & \dots \\ D_{x_{n_1+1}} f_m(x_0) & \dots & D_{x_n} f_m(x_0) \end{bmatrix}_{m \times n_2}$$

and therefore

$$Df(x_0) := \begin{bmatrix} D_{x'} f(x_0) & D_{x''} f(x_0) \end{bmatrix}_{m \times n}$$

Definition 616 Given an open set $S \subseteq \mathbb{R}^n$ and $f : S \rightarrow \mathbb{R}$, assume that $\forall x \in S$, $Df(x) := \left(\frac{\partial f(x)}{\partial x_j} \right)_{j=1}^n$ exists. Then, $\forall j \in \{1, \dots, n\}$, we define the j -th partial derivative function as

$$\frac{\partial f}{\partial x_j} : S \rightarrow \mathbb{R}, \quad x \mapsto \frac{\partial f(x)}{\partial x_j}$$

Assuming that the above function has partial derivative with respect to x_k for $k \in \{1, \dots, n\}$, we define it as the mixed second order partial derivative of f with respect to x_j and x_k and we write

$$\frac{\partial^2 f(x)}{\partial x_j \partial x_k} := \frac{\partial \frac{\partial f(x)}{\partial x_j}}{\partial x_k}$$

Definition 617 Given $f : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$, the Hessian matrix of f at x_0 is the $n \times n$ matrix

$$D^2 f(x_0) := \left[\frac{\partial^2 f}{\partial x_j \partial x_k}(x_0) \right]_{j,k=1,\dots,n}$$

Remark 618 $D^2 f(x_0)$ is the Jacobian matrix of the gradient function of f .

Example 619 Compute the Hessian function of $f : \mathbb{R}_{++}^3 \rightarrow \mathbb{R}$,

$$f(x, y, z) = (e^x \cos y + z^2 + x^2 \log y + \log x + \log z + 2t \log t)$$

We first compute the gradient:

$$\begin{pmatrix} 2x \ln y + (\cos y) e^x + \frac{1}{x} \\ -(\sin y) e^x + \frac{x^2}{y} \\ 2z + \frac{1}{z} \\ 2 \ln t + 2 \end{pmatrix}$$

and then the Hessian matrix

$$\begin{bmatrix} 2 \ln y + (\cos y) e^x - \frac{1}{x^2} & -(\sin y) e^x + \frac{2x}{y} & 0 & 0 \\ -(\sin y) e^x + \frac{2x}{y} & -(\cos y) e^x - \frac{x}{y^2} & 0 & 0 \\ 0 & 0 & -\frac{1}{z^2} + 2 & 0 \\ 0 & 0 & 0 & \frac{2}{t} \end{bmatrix}$$

14.1 The chain rule

Proposition 620 (Chain Rule) Given $S \subseteq \mathbb{R}^n$, $T \subseteq \mathbb{R}^m$,

$$f : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m,$$

such that $\text{Im } f \subseteq T$, and

$$g : T \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^p,$$

assume that f and g are differentiable in x_0 and $y_0 = f(x_0)$, respectively. Then

$$h : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^p, \quad h(x) = (g \circ f)(x)$$

is differentiable in x_0 and

$$dh_{x_0} = dg_{f(x_0)} \circ df_{x_0}.$$

Proof. We want to show that there exists a linear function $dh_{x_0} : \mathbb{R}^n \rightarrow \mathbb{R}^p$ such that

$$h(x_0 + u) = h(x_0) + dh_{x_0}(u) + \|u\| \cdot E_{x_0}^*(u), \quad \text{with} \quad \lim_{u \rightarrow 0} E_{x_0}^*(u) = 0,$$

and $dh_{x_0} = dg_{f(x_0)} \circ df_{x_0}$.

Taking u sufficiently small (in order to have $x_0 + u \in S$), we have

$$h(x_0 + u) - h(x_0) = g[f(x_0 + u)] - g[f(x_0)] = g[f(x_0 + u)] - g(y_0)$$

and defined

$$v = f(x_0 + u) - y_0$$

we get

$$h(x_0 + u) - h(x_0) = g(y_0 + v) - g(y_0).$$

Since f is differentiable in x_0 , we get

$$v = df_{x_0}(u) + \|u\| \cdot E_{x_0}(u), \quad \text{with} \quad \lim_{u \rightarrow 0} E_{x_0}(u) = 0. \quad (14.1)$$

Since g is differentiable in $y_0 = f(x_0)$, we get

$$g(y_0 + v) - g(y_0) = dg_{y_0}(v) + \|v\| \cdot E_{y_0}(v), \quad \text{with} \quad \lim_{v \rightarrow 0} E_{y_0}(v) = 0. \quad (14.2)$$

Inserting (14.1) in (14.2), we get

$$g(y_0 + v) - g(y_0) = dg_{y_0}(df_{x_0}(u) + \|u\| \cdot E_{x_0}(u)) + \|v\| \cdot E_{y_0}(v) = dg_{y_0}(df_{x_0}(u)) + \|u\| \cdot dg_{y_0}(E_{x_0}(u)) + \|v\| \cdot E_{y_0}(v)$$

Defined

$$E_{x_0}(u) := \begin{cases} 0 & \text{if } u = 0 \\ df_{y_0}(E_{x_0}(u)) + \frac{\|v\|}{\|u\|} \cdot E_{y_0}(v) & \text{if } u \neq 0 \end{cases},$$

we are left with showing that

$$\lim_{u \rightarrow 0} E_{x_0}(u) = 0.$$

Observe that

$$\lim_{u \rightarrow 0} df_{y_0}(E_{x_0}(u)) = 0$$

since linear functions are continuous and from (14.1). Moreover, since $\lim_{u \rightarrow 0} v = \lim_{u \rightarrow 0} (f(x_0 + u) - y_0) = 0$, from (14.2), we get

$$\lim_{u \rightarrow 0} E_{y_0}(v) = 0.$$

Finally, we have to show that $\lim_{u \rightarrow 0} \frac{\|v\|}{\|u\|}$ is bounded. Now, from the definition of u and from (612), defined $\alpha := \sum_{j=1}^m \|Df_j(x_0)\|$,

$$\|v\| = \|df_{x_0}(u) + \|u\| \cdot E_{x_0}(u)\| \leq \|df_{x_0}(u)\| + \|u\| \|E_{x_0}(u)\| \leq (\alpha + \|E_{x_0}(u)\|) \cdot \|u\|$$

and

$$\lim_{u \rightarrow 0} \frac{\|v\|}{\|u\|} \leq \lim_{u \rightarrow 0} (\alpha + \|E_{x_0}(u)\|) = \alpha,$$

as desired. ■

Remark 621 From Proposition 610 and Proposition 292, or simply (7.10), we also have

$$Dh(x_0)_{p \times n} = Dg(f(x_0))_{p \times m} \cdot Df(x_0)_{m \times n}.$$

Observe that $Dg(f(x_0))$ is obtained computing $Dg(y)$ and then substituting $f(x_0)$ in the place of y . We therefore also write $Dg(f(x_0)) = Dg(y)|_{y=f(x_0)}$

Exercise 622 Compute dh_{x_0} , if $n = 1, p = 1$.

Definition 623 Given $f : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^{m_1}$, $g : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^{m_2}$,

$$(f, g) : \mathbb{R}^n \rightarrow \mathbb{R}^{m_1+m_2}, \quad (f, g)(x) = (f(x), g(x))$$

Remark 624 Clearly,

$$D(f, g)(x_0) = \begin{bmatrix} Df(x_0) \\ Dg(x_0) \end{bmatrix}$$

Example 625 Given

$$\begin{aligned} f : \mathbb{R} &\rightarrow \mathbb{R}^2, & x &\mapsto (\sin x, \cos x) \\ g : \mathbb{R}^2 &\rightarrow \mathbb{R}^2, & (y_1, y_2) &\mapsto (y_1 + y_2, y_1 \cdot y_2) \\ h = g \circ f : \mathbb{R} &\rightarrow \mathbb{R}^2, & x &\mapsto (\sin x + \cos x, \sin x \cdot \cos x) \end{aligned}$$

verify the conclusion of the Chain Rule Proposition.

$$\begin{aligned} Df(x) &= \begin{bmatrix} \cos x \\ -\sin x \end{bmatrix} \\ Dg(y) &= \begin{bmatrix} 1 & 1 \\ y_2 & y_1 \end{bmatrix} \\ Dg(f(x)) &= \begin{bmatrix} 1 & 1 \\ \cos x & \sin x \end{bmatrix} \\ Dg(f(x)) \cdot Df(x) &= \begin{bmatrix} 1 & 1 \\ \cos x & \sin x \end{bmatrix} \begin{bmatrix} \cos x \\ -\sin x \end{bmatrix} = \begin{bmatrix} \cos x - \sin x \\ \cos^2 x - \sin^2 x \end{bmatrix} = Dh(x) \end{aligned}$$

Example 626 Take

$$\begin{aligned} g : \mathbb{R}^k &\rightarrow \mathbb{R}^n, & t &\mapsto g(t) \\ f : \mathbb{R}^n \times \mathbb{R}^k &\rightarrow \mathbb{R}^m, & (x, t) &\mapsto f(x, t) \\ h : \mathbb{R}^k &\rightarrow \mathbb{R}^m, & t &\mapsto f(g(t), t) \end{aligned}$$

Then

$$\tilde{g} := (g, id_{\mathbb{R}^k}) : \mathbb{R}^k \rightarrow \mathbb{R}^n \times \mathbb{R}^k, \quad t \mapsto (g(t), t)$$

and

$$h = f \circ \tilde{g} = f \circ (g, id_{\mathbb{R}^k},)$$

Therefore, assuming that f, g, h are differentiable,

$$\begin{aligned} [Dh(t_0)]_{m \times k} &= [Df(g(t_0), t_0)]_{m \times (n+k)} \cdot \begin{bmatrix} Dg(t_0) \\ I \end{bmatrix}_{(n+k) \times k} = \\ &= [[D_x f(g(t_0), t_0)]_{m \times n} \mid [D_t f(g(t_0), t_0)]_{m \times k}] \cdot \begin{bmatrix} [Dg(t_0)]_{n \times k} \\ I_{k \times k} \end{bmatrix} = \\ &= [D_x f(g(t_0), t_0)]_{m \times n} \cdot [Dg(t_0)]_{n \times k} + [D_t f(g(t_0), t_0)]_{m \times k} \end{aligned}$$

In the case $k = n = m = 1$, the above expression

$$\frac{df(x = g(t), t)}{dt} = \frac{\partial f(g(t), t)}{\partial x} \frac{dg(t)}{dt} + \frac{\partial f(g(t), t)}{\partial t}$$

or

$$\frac{df(g(t), t)}{dt} = \frac{\partial f(x, t)}{\partial x} \Big|_{x=g(t)} \cdot \frac{dg(t)}{dt} + \frac{\partial f(x, t)}{\partial t} \Big|_{x=g(t)}$$

14.2 Mean value theorem

Proposition 627 (Mean Value Theorem) Let S be an open subset of \mathbb{R}^n and $f : S \rightarrow \mathbb{R}^m$ a differentiable function. Let $x, y \in S$ be such that the line segment joining them is contained in S , i.e.,

$$L(x, y) := \{z \in \mathbb{R}^n : \exists \lambda \in [0, 1] \text{ such that } z = (1 - \lambda)x + \lambda y\} \subseteq S.$$

Then

$$\forall a \in \mathbb{R}^m, \quad \exists z \in L(x, y) \quad \text{such that} \quad a \cdot [f(y) - f(x)] = a \cdot [Df(z) \cdot (y - x)]$$

Remark 628 Under the assumptions of the above theorem, the following conclusion is **false**:

$$\exists z \in L(x, y) \text{ such that } f(y) - f(x) = Df(z) \cdot (y - x).$$

But if $f : S \rightarrow \mathbb{R}^{m=1}$, then setting $a \in \mathbb{R}^{m=1}$ equal to 1, we get that the above statement is indeed true.

Proof. of Proposition 627

Define $u = y - x$. Since S is open and $L(x, y) \subseteq S$, $\exists \delta > 0$ such that $\forall t \in (-\delta, 1 + \delta)$ such that $x + tu = (1 - t)x + ty \in S$. Taken $a \in \mathbb{R}^m$, define

$$F : (-\delta, 1 + \delta) \rightarrow \mathbb{R}, \quad t \mapsto a \cdot f(x + tu) = \sum_{j=1}^m a_j \cdot f_j(x + tu)$$

Then

$$F'(t) = \sum_{j=1}^m a_j \cdot [Df_j(x + tu)]_{1 \times n} \cdot u_{n \times 1} = a_{1 \times m} \cdot [Df(x + tu)]_{m \times n} \cdot u_{n \times 1}$$

and F is continuous on $[0, 1]$ and differentiable on $(0, 1)$; then, we can apply “Calculus 1” Mean Value Theorem and conclude that

$$\exists \theta \in (0, 1) \text{ such that } F(1) - F(0) = F'(\theta),$$

and by definition of F and u ,

$$\exists \theta \in (0, 1) \text{ such that } f(y) - f(x) = a \cdot Df(x + \theta u) \cdot (y - x)$$

which choosing $z = x + \theta u$ gives the desired result. ■

Remark 629 Using the results we have seen on directional derivatives, the conclusion of the above theorem can be rewritten as follows.

$$\exists z \in L(x, y) \text{ such that } f(y) - f(x) = f'(z, y - x)$$

As in the case of real functions of real variables, the Mean Value Theorem allows to give a simple relationship between sign of the derivative and monotonicity.

Definition 630 A set $C \subseteq \mathbb{R}^n$ is convex if $\forall x_1, x_2 \in C$ and $\forall \lambda \in [0, 1]$, $(1 - \lambda)x_1 + \lambda x_2 \in C$.

Proposition 631 Let S be an open and convex subset of \mathbb{R}^n and $f : S \rightarrow \mathbb{R}^m$ a differentiable function. If $\forall x \in S$, $df_x = 0$, then f is constant on S .

Proof. Take arbitrary $x, y \in S$. Then since S is convex and f is differential, from the Mean Value Theorem, we have that

$$\forall a \in \mathbb{R}^m, \quad \exists z \in L(x, y) \text{ such that } a \cdot [f(y) - f(x)] = a \cdot [Df(z) \cdot (y - x)] = 0.$$

Taken $a = f(y) - f(x)$, we get that

$$\|f(y) - f(x)\| = 0$$

and therefore

$$f(x) = f(y),$$

as desired. ■

Definition 632 Given $x := (x_i)_{i=1}^n, y := (y_i)_{i=1}^n \in \mathbb{R}^n$,

$$x \geq y \quad \text{means} \quad \forall i \in \{1, \dots, n\}, \quad x_i \geq y_i;$$

$$x > y \quad \text{means} \quad x \geq y \quad \wedge \quad x \neq y;$$

$$x \gg y \quad \text{means} \quad \forall i \in \{1, \dots, n\}, \quad x_i > y_i.$$

Definition 633 $f : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ is increasing if $\forall x, y \in S$, $x > y \Rightarrow f(x) \geq f(y)$.
 f is strictly increasing if $\forall x, y \in S$, $x > y \Rightarrow f(x) > f(y)$.

Proposition 634 Take an open, convex subset S of \mathbb{R}^n , and $f : S \rightarrow \mathbb{R}$ differentiable.

1. If $\forall x \in S$, $Df(x) \geq 0$, then f is increasing;
2. If $\forall x \in S$, $Df(x) \gg 0$, then f is strictly increasing.

Proof. 1. Take $y \geq x$. Since S is convex, $L(x, y) \subseteq S$. Then from the Mean Value Theorem,

$$\exists z \in L(x, y) \text{ such that } f(y) - f(x) = Df(z) \cdot (y - x)$$

Since $y - x \geq 0$ and $Df(z) \geq 0$, the result follows.

2. Take $x > y$. Since S is convex, $L(x, y) \subseteq S$. Then from the Mean Value Theorem,

$$\exists z \in L(x, y) \text{ such that } f(y) - f(x) = Df(z) \cdot (y - x)$$

Since $y - x > 0$ and $Df(z) > 0$, the result follows. ■

Exercise 635 Is the following statement correct: “If $\forall x \in S$, $Df(x) > 0$, then f is strictly increasing”?

Corollary 636 Take an open, convex subset S of \mathbb{R}^n , and $f \in C^1(S, \mathbb{R})$. If $\exists x_0 \in S$ and $u \in \mathbb{R}^n \setminus \{0\}$ such that $f'(x_0, u) > 0$, then $\exists \bar{t} \in \mathbb{R}_{++}$ such that $\forall t \in [0, \bar{t}]$,

$$f(x_0 + tu) \geq f(x_0).$$

Proof. Since f is C^1 and $f'(x_0, u) = Df(x_0) \cdot u > 0$, $\exists r > 0$ such that

$$\forall x \in B(x_0, r), \quad f'(x, u) > 0.$$

Then $\forall t \in (-r, r)$, $\left\|x_0 + \frac{1}{\|u\|}tu - x_0\right\| = t < r$, and therefore

$$f'\left(x_0 + \frac{t}{\|u\|}u, u\right) > 0$$

Then, from the Mean Value Theorem, $\forall t \in [0, \frac{r}{2}]$,

$$f(x_0 + tu) - f(x_0) = f'(x_0 + tu, u) \geq 0.$$

■

Definition 637 Given a function $f : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$, $x_0 \in S$ is a point of local maximum for f if

$$\exists \delta > 0 \text{ such that } \forall x \in B(x_0, \delta), \quad f(x_0) \geq f(x);$$

x_0 is a point of global maximum for f if

$$\forall x \in S, \quad f(x_0) \geq f(x).$$

$x_0 \in S$ is a point of strict local maximum for f if

$$\exists \delta > 0 \text{ such that } \forall x \in B(x_0, \delta), \quad f(x_0) > f(x);$$

x_0 is a point of strict global maximum for f if

$$\forall x \in S, \quad f(x_0) > f(x).$$

Local, global, strict minima are defined in obvious manner

Proposition 638 If $S \subseteq \mathbb{R}^n$, $f : S \rightarrow \mathbb{R}$ admits all partial derivatives in $x_0 \in \text{Int } S$ and x_0 is a point of local maximum or minimum, then $Df(x_0) = 0$.

Proof. Since x_0 is a point of local maximum, $\exists \delta > 0$ such that $\forall x \in B(x_0, \delta)$, $f(x_0) \geq f(x)$. As in Remark 583, for any $k \in \{1, \dots, n\}$, define

$$g_k : \mathbb{R} \rightarrow \mathbb{R}, \quad g_k(x_k) = f(x_0 + (x_k - x_{0k})e_n^k).$$

Then g_k has a local maximum point at x_{0k} . Then from Calculus 1,

$$g'_k(x_{0k}) = 0$$

Since, again from Remark 583, we have

$$D_k f(x_0) = g'_k(x_0).$$

the result follows. ■

14.3 A sufficient condition for differentiability

Definition 639 $f = (f_i)_{i=1}^m : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ is continuously differentiable on $A \subseteq S$, or f is C^1 on S , or $f \in \mathcal{C}(A, \mathbb{R}^m)$ if $\forall i \in \{1, \dots, m\}, j \in \{1, \dots, n\}$,

$$D_{x_j} f_i : A \rightarrow \mathbb{R}, \quad x \mapsto D_{x_j} f_i(x) \quad \text{is continuous.}$$

Definition 640 $f : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$ is twice continuously differentiable on $A \subseteq S$, or f is C^2 on S , or $f \in \mathcal{C}^2(A, \mathbb{R}^m)$ if $\forall j, k \in \{1, \dots, n\}$,

$$\frac{\partial^2 f}{\partial x_j \partial x_k} : A \rightarrow \mathbb{R}, \quad x \mapsto \frac{\partial^2 f}{\partial x_j \partial x_k}(x) \quad \text{is continuous.}$$

Proposition 641 If f is C^1 in an open neighborhood of x_0 , then it is differentiable in x_0 .

Proof. See Apostol (1974), page 357 or Section "Existence of derivative", page 232, in Bartle (1964), where the significant case $f : \mathbb{R}^n \rightarrow \mathbb{R}^{m=1}$ is presented using Cauchy-Schwarz inequality. See also, Theorem 1, page 197, in Taylor and Mann (1984), for the case $f : \mathbb{R}^2 \rightarrow \mathbb{R}$. ■

The above result is the answer to Question 3 in Remark 609. To show that $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is differentiable, it is enough to verify that all its partial derivatives, i.e., the entries of the Jacobian matrix, are continuous functions.

14.4 A sufficient condition for equality of mixed partial derivatives

Proposition 642 If $f : S \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^1$ is C^2 in an open neighborhood of x_0 , then $\forall i$

$$\frac{\partial \frac{\partial f}{\partial x_i}}{\partial x_k}(x_0) = \frac{\partial \frac{\partial f}{\partial x_k}}{\partial x_i}(x_0)$$

or

$$\frac{\partial^2 f}{\partial x_i \partial x_k}(x_0) = \frac{\partial^2 f}{\partial x_k \partial x_i}(x_0)$$

Proof. See Apostol (1974), Section 12.13, page 358. ■

14.5 Taylor's theorem for real valued functions

To get Taylor's theorem for functions $f : \mathbb{R}^m \rightarrow \mathbb{R}$, we introduce some notation in line with the definition of directional derivative:

$$f'(x, u) = \sum_{i=1}^n D_{x_i} f(x) \cdot u_i.$$

Definition 643 Assume S is an open subset of \mathbb{R}^m and the function $f : S \rightarrow \mathbb{R}$ admits partial derivatives at least up to order m , and $x \in S, u \in \mathbb{R}^m$. Then

$$f''(x, u) := \sum_{i=1}^n \sum_{j=1}^n D_{i,j} f(x) \cdot u_i \cdot u_j,$$

$$f'''(x, u) := \sum_{i=1}^n \sum_{j=1}^n \sum_{k=1}^n D_{i,j,k} f(x) \cdot u_i \cdot u_j \cdot u_k$$

and similar definition applies to $f^{(m)}(x, u)$.

Proposition 644 (Taylor's formula) Assume S is an open subset of \mathbb{R}^m and the function $f : S \rightarrow \mathbb{R}$ admits partial derivatives at least up to order m , and $x \in S$, $u \in \mathbb{R}^m$. Assume also that all its partial derivative of order $< m$ are differentiable. If y and x are such that $L(y, x) \subseteq S$, then there exists $z \in L(y, x)$ such that

$$f(y) = f(x) + \sum_{k=1}^{m-1} \frac{1}{k!} f^{(k)}(x, y-x) + \frac{1}{m!} f^{(m)}(z, y-x).$$

Proof. . Since S is open and $L(x, y) \subseteq S$, $\exists \delta > 0$ such that $\forall t \in (-\delta, 1 + \delta)$ such that $x + t(y-x) \in S$. Define $g : (-\delta, 1 + \delta) \rightarrow \mathbb{R}$

$$g(t) = f(x + t(y-x)).$$

From standard "Calculus 1" Taylor's theorem, we have that $\exists \theta \in (0, 1)$ such that

$$f(y) - f(x) = g(1) - g(0) = \sum_{k=1}^{m-1} \frac{1}{k!} g^{(k)}(0) + \frac{1}{m!} g^{(m)}(\theta).$$

Then

$$g'(t) = Df(x + t(y-x)) \cdot (y-x) = \sum_{i=1}^n D_{x_i} f(x + t(y-x)) \cdot (y_i - x_i) = f'(x + t(y-x), y-x),$$

$$g''(t) = \sum_{i=1}^n \sum_{j=1}^n D_{x_i, x_j} f(x + t(y-x)) \cdot (y_i - x_i) \cdot (y_j - x_j) = f''(x + t(y-x), y-x)$$

and similarly

$$g^{(m)}(t) = f^{(m)}(x + t(y-x), y-x)$$

Then the desired result follow substituting 0 in the place of t where needed and choosing $z = x + \theta(y-x)$. ■

Chapter 15

Implicit function theorem

15.1 Some intuition

Below, we present an informal discussion of the Implicit Function Theorem. Assume that

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad (x, t) \mapsto f(x, t)$$

is at least C^1 . The basic goal is to study the nonlinear equation

$$f(x, t) = 0,$$

where x can be interpreted as an endogenous variable and t as a parameter (or an exogenous variable). Assume that

$$\exists (x^0, t^0) \in \mathbb{R}^2 \text{ such that } f(x^0, t^0) = 0$$

and for some $\varepsilon > 0$

$$\exists \text{ a } C^1 \text{ function } g : (t^0 - \varepsilon, t^0 + \varepsilon) \rightarrow \mathbb{R}, t \mapsto g(t)$$

such that

$$f(g(t), t) = 0 \tag{15.1}$$

We can then say that g describes *the solution* to the equation

$$f(x, t) = 0,$$

in the unknown variable x and parameter t , in an open neighborhood of t^0 . Therefore, using the Chain Rule - and in fact, Remark 626 - applied to both sides of (15.1), we get

$$\frac{\partial f(x, t)}{\partial x} \Big|_{x=g(t)} \cdot \frac{dg(t)}{dt} + \frac{\partial f(x, t)}{\partial t} \Big|_{x=g(t)} = 0$$

and

$$\text{assuming that } \frac{\partial f(x, t)}{\partial x} \Big|_{x=g(t)} \neq 0$$

we have


$$\frac{dg(t)}{dt} = - \frac{\frac{\partial f(x, t)}{\partial t} \Big|_{x=g(t)}}{\frac{\partial f(x, t)}{\partial x} \Big|_{x=g(t)}} \tag{15.2}$$

The above expression is the derivative of the function implicitly defined by (15.1) close to the value t^0 . In other words, it is the slope of the level curve $f(x, t) = 0$ at the point $(t, g(t))$.

For example, taken

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad (x, t) \mapsto x^2 + t^2 - 1$$

$f(x, t) = 0$ describes the circle with center in the origin and radius equal to 1. Putting t on the horizontal axis and x on the vertical axis, we have the following picture.



Clearly

$$f((0, 1)) = 0$$

As long as $t \in (-1, 1)$, $g(t) = \sqrt{1 - t^2}$ is such that

$$f(g(t), t) = 0 \quad (15.3)$$

Observe that

$$\frac{d(\sqrt{1 - t^2})}{dt} = -\frac{t}{\sqrt{1 - t^2}}$$

and

$$-\frac{\frac{\partial f(x, t)}{\partial t} |_{x=g(t)}}{\frac{\partial f(x, t)}{\partial x} |_{x=g(t)}} = -\frac{2t}{2x |_{x=g(t)}} = -\frac{t}{\sqrt{1 - t^2}}$$

For example for $t = \frac{1}{\sqrt{2}}$, $g'(t) = -\frac{\frac{1}{\sqrt{2}}}{\sqrt{1 - \frac{1}{2}}} = -1$.

Let's try to present a more detailed geometrical interpretation¹. Consider the set $\{(x, t) \in \mathbb{R}^2 : f(x, t) = 0\}$ presented in the following picture.

Insert picture a., page 80.

In this case, does equation

$$f(x, t) = 0 \quad (15.4)$$

define x as a function of t ? Certainly, the curve presented in the picture is not the graph of a function with x as dependent variable and t as an independent variable for all values of t in \mathbb{R} . In fact,

1. if $t \in (-\infty, t_1]$, there is only one value of x which satisfies equation (15.4);
2. if $t \in (t_1, t_2)$, there are two values of x for which $f(x, t) = 0$;
3. if $t \in (t_2, +\infty)$, there are no values satisfying the equation.

If we consider t belonging to an interval contained in (t_1, t_2) , we have to restrict the admissible range of variation of x in order to conclude that equation (15.4) defines x as a function of t in that interval. For example, we see that if the rectangle R is as indicated in the picture, the given equation defines x as a function of t , for well chosen domain and codomain - naturally associated with R . The graph of that function is indicated in the figure below.

Insert picture b., page 80.

¹This discussion is taken from Sydsaeter (1981), page 80-81.

The size of R is limited by the fact that we need to define a *function* and therefore one and only one value has to be associated with t . Similar rectangles and associated solutions to the equation can be constructed for all other points on the curve, *except one*: (t_2, x_2) . Irrespectively of how small we choose the rectangle around that point, there will be values of t close to t_2 , say t' , such that there are two values of x , say x' and x'' , with the property that both (t', x') and (t', x'') satisfy the equation and lie inside the rectangle. Therefore, equation (15.4) does not define x as a function of t in an open neighborhood of the point (t_2, x_2) . In fact, there the slope of the tangent to the curve is infinite. If you try to use expression (15.2) to compute the slope of the curve defined by $x^2 + t^2 = 1$ in the point $(1, 0)$, you get an expression with *zero* in the denominator.

On the basis of the above discussion, we see that it is crucial to require the condition

$$\frac{\partial f(x, t)}{\partial x} \Big|_{x=g(t)} \neq 0$$

to insure the possibility of locally writing x as a solution (to (15.4)) function of t .

We can informally, summarize what we said as follow.

If f is C^1 , $f(x_0, t_0) = 0$ and $\frac{\partial f(x, t)}{\partial x} \Big|_{(x, t)=(x_0, t_0)} \neq 0$, then $f(x, t) = 0$ define x as a C^1 function g of t in an open neighborhood of t^0 , and $g'(t) = -\frac{\frac{\partial f(x, t)}{\partial t}}{\frac{\partial f(x, t)}{\partial x} \Big|_{x=g(t)}}$.

Next sections provide a formal statement and proof of the Implicit Function Theorem. Some work is needed.

15.2 Functions with full rank square Jacobian

Proposition 645 Taken $a \in \mathbb{R}^n$, $r \in \mathbb{R}_{++}$, assume that

1. $f := (f_i)_{i=1}^n : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is continuous on $\text{Cl}(B(a, r))$;
2. $\forall x \in B(a, r)$, $[Df(x)]_{n \times n}$ exists and $\det Df(x) \neq 0$;
3. $\forall x \in \mathcal{F}(B(a, r))$, $f(x) \neq f(a)$.

Then, $\exists \delta \in \mathbb{R}_{++}$ such that

$$f(B(a, r)) \supseteq B(f(a), \delta).$$

Proof. Define $B := B(a, r)$ and

$$g : \mathcal{F}(B) \rightarrow \mathbb{R}, \quad x \mapsto \|f(x) - f(a)\|.$$

From Assumption 3, $\forall x \in \mathcal{F}(B)$, $g(x) > 0$. Moreover, since g is continuous and $\mathcal{F}(B)$ is compact, g attains a global minimum value $m > 0$ on $\mathcal{F}(B)$. Take $\delta = \frac{m}{2}$; to prove the desired result, it is enough to show that $T := B(f(a), \delta) \subseteq f(B)$, i.e., $\forall y \in T$, $y \in f(B)$. Define

$$h : \text{Cl}(B) \rightarrow \mathbb{R}, \quad x \mapsto \|f(x) - y\|.$$

since h is continuous and $\text{Cl} B$ is compact, h attains a global minimum in a point $c \in \overline{B}$. We now want to show that $c \in B$. Observe that, since $y \in T = B(f(a), \delta = \frac{m}{2})$,

$$h(a) = \|f(a) - y\| < \frac{m}{2} \tag{15.5}$$

Therefore, since c is a global minimum point for h , it must be the case that $h(c) < \frac{m}{2}$. Now take $x \in \mathcal{F}(B)$; then

$$h(x) = \|f(x) - y\| = \|f(x) - f(a) - (y - f(a))\| \stackrel{(1)}{\geq} \|f(x) - f(a)\| - \|y - f(a)\| \stackrel{(2)}{\geq} g(x) - \frac{m}{2} \stackrel{(3)}{\geq} \frac{m}{2},$$

where (1) follows from Remark 54, (2) from (15.5) and (3) from the fact that g has minimum value equal to m . Therefore, $\forall x \in \mathcal{F}(B)$, $h(x) > h(a)$ and h does not attain its minimum on $\mathcal{F}(B)$. Then h and h^2 get their minimum at $c \in B^2$. Since

$$H(x) := h^2(x) = \|f(x) - y\|^2 = \sum_{i=1}^n (f_i(x) - y_i)^2$$

² $\forall x \in \overline{B}$, $h(x) \geq 0$ and $h(x) \geq h(c)$. Therefore, $h^2(x) \geq h^2(c)$.

from Proposition 638, $DH(c) = 0$, i.e.,

$$\forall k \in \{1, \dots, n\}, \quad 2 \sum_{i=1}^n D_{x_k} f_i(c) \cdot (f_i(c) - y_i) = 0$$

i.e.,

$$[Df(c)]_{n \times n} (f(c) - y)_{n \times 1} = 0.$$

Then, from assumption 2,

$$f(c) = y,$$

i.e., since $c \in B$, $y \in f(B)$, ad desired. ■

Proposition 646 (*1st sufficient condition for openness of a function*)

Let an open set $A \subseteq \mathbb{R}^n$ and a function $f : A \rightarrow \mathbb{R}^n$ be given. If

1. f is continuous,
2. f is one-to-one,
3. $\forall x \in A$, $Df(x)$ exists and $\det Df(x) \neq 0$,

then f is open.

Proof. Taken $b \in f(A)$, there exists $a \in A$ such that $f(a) = b$. Since A is open, there exists $r \in \mathbb{R}_{++}$ such that $B(a, r) \subseteq A$. Moreover, since f is one-to-one and since $a \notin \mathcal{F}(B)$, $\forall x \in \mathcal{F}(B)$, $f(x) \neq f(a)$. Then³, for sufficiently small r , $\text{Cl}(B(a, r)) \subseteq A$, and the assumptions of Proposition 645 are satisfied and there exists $\delta \in \mathbb{R}_{++}$ such that

$$f(A) \supseteq f(\text{Cl}(B(a, r))) \supseteq B(f(a), \delta),$$

as desired. ■

Definition 647 Given $f : S \rightarrow T$, and $A \subseteq S$, the function $f|_A$ is defined as follows

$$f|_A : A \rightarrow f(A), \quad f|_A(x) = f(x).$$

Proposition 648 Let an open set $A \subseteq \mathbb{R}^n$ and a function $f : A \rightarrow \mathbb{R}^n$ be given. If

1. f is C^1 ,
 2. $\exists a \in A$ such that $\det Df(a) \neq 0$,
- then $\exists r \in \mathbb{R}_{++}$ such that f is one-to-one on $B(a, r)$, and, therefore, $f|_{B(a, r)}$ is invertible.

Proof. Consider $(\mathbb{R}^n)^n$ with generic element $z := (z^i)_{i=1}^n$, where $\forall i \in \{1, \dots, n\}$, $z^i \in \mathbb{R}^n$, and define

$$h : \mathbb{R}^{n^2} \rightarrow \mathbb{R}, \quad : (z^i)_{i=1}^n \mapsto \det \begin{bmatrix} Df_1(z^1) \\ \vdots \\ Df_i(z^i) \\ \vdots \\ Df_n(z^n) \end{bmatrix}.$$

Observe that h is continuous because f is C^1 and the determinant function is continuous in its entries. Moreover, from Assumption 2,

$$h(a, \dots, a, \dots, a) = \det Df(a) \neq 0.$$

Therefore, $\exists r \in \mathbb{R}_{++}$ such that

$$\forall (z^i)_{i=1}^n \in B((a, \dots, a, \dots, a), r'), \quad h((z^i)_{i=1}^n) \neq 0.$$

Observe that $B^* := B((a, \dots, a, \dots, a), r) \cap \beta \neq \emptyset$, where $\beta := \{(z^i)_{i=1}^n \in \mathbb{R}^{n^2} : \forall i \in \{1, \dots, n\}, z^i = z^1\}$. Define

$$\text{proj} : (\mathbb{R}^n)^n \rightarrow \mathbb{R}^n, \text{proj} : (z^i)_{i=1}^n \mapsto z^1,$$

³Simply observe that $\forall r \in \mathbb{R}_{++}$, $\overline{B}(x, \frac{r}{2}) \subseteq B(x, r)$.

and observe that $\text{proj}(B^*) = B(a, r) \subseteq \mathbb{R}^n$ and $\forall i \in \{1, \dots, n\}, \forall z_i \in B(a, r), (z^1, \dots, z^i, \dots, z^n) \in B^*$ and therefore $h(z^1, \dots, z^i, \dots, z^n) \neq 0$, or, summarizing,

$$\exists r \in \mathbb{R}_{++} \text{ such that } \forall i \in \{1, \dots, n\}, \forall z_i \in B(a, r), \quad h(z^1, \dots, z^i, \dots, z^n) \neq 0$$

We now want to show that f is one-to-one on $B(a, r)$. Suppose otherwise, i.e., given $x, y \in B(a, r)$, $f(x) = f(y)$, but $x \neq y$. We can now apply the Mean Value Theorem (see Remark 628) to f^i for any $i \in \{1, \dots, n\}$ on the segment $L(x, y) \subseteq B(a, r)$. Therefore $\forall i \in \{1, \dots, n\}, \exists z^i \in L(x, y)$ such that

$$\forall i \in \{1, \dots, n\}, \exists z^i \in L(x, y) \text{ such that } 0 = f_i(x) - f_i(y) = Df(z^i)(y - x)$$

i.e.,


$$\begin{bmatrix} Df_1(z^1) \\ \dots \\ Df_i(z^i) \\ \dots \\ Df_n(z^n) \end{bmatrix} (y - x) = 0$$

Observe that $\forall i, z^i \in B(a, r)$ and therefore $(z^i)_{i=1}^n \in B((a, \dots, a, \dots, a), r'')$, and therefore

$$\det \begin{bmatrix} Df_1(z^1) \\ \dots \\ Df_i(z^i) \\ \dots \\ Df_n(z^n) \end{bmatrix} = h\left((z^i)_{i=1}^n\right) \neq 0$$

and therefore $y = x$, a contradiction. ■

Remark 649 *The above result is not a global result, i.e., it is false that if f is C^1 and its Jacobian has full rank everywhere in the domain, then f is one to one. Just take the function \tan .*



The next result gives a global property.

Proposition 650 (2nd sufficient condition for openness of a function) *Let an open set $A \subseteq \mathbb{R}^n$ and a function $f : A \rightarrow \mathbb{R}^n$ be given. If*

1. f is C^1 ,
 2. $\forall x \in A, \det Df(x) \neq 0$,
- then f is an open function.*

Proof. Take an open set $S \subseteq A$. From Proposition 648, $\forall x \in S$ there exists $r \in \mathbb{R}_{++}$ such that f is one-to-one on $B(x, r)$. Then, from Proposition 646, $f(B(x, r))$ is open in \mathbb{R}^n . We can then write $S = \cup_{x \in S} B(x, r)$ and

$$f(S) = f(\cup_{x \in S} B(x, r)) = \cup_{x \in S} f(B(x, r))$$

where the second equality follows from Proposition 518.2.f1, and then $f(S)$ is an open set. ■

15.3 The inverse function theorem

Proposition 646 shows that a C^1 function with full rank square Jacobian in a point a has a local inverse in an open neighborhood of a . The inverse function theorem give local differentiability properties of that local inverse function.

Lemma 651 *If g is the inverse function of $f : X \rightarrow Y$ and $A \subseteq X$, then $g|_{f(A)}$ is the inverse of $f|_A$, and*

if g is the inverse function of $f : X \rightarrow Y$ and $B \subseteq Y$, then $g|_B$ is the inverse of $f|_{g(B)}$.

Proof. Exercise. ■

Proposition 652 *Let an open set $S \subseteq \mathbb{R}^n$ and a function $f : S \rightarrow \mathbb{R}^n$ be given. If*

1. f is C^1 , and
 2. $\exists a \in S$, $\det Df(a) \neq 0$,
- then there exist two open sets $X \subseteq S$ and $Y \subseteq f(S)$ and a unique function g such that*
1. $a \in X$ and $f(a) \in Y$,
 2. $Y = f(X)$,
 3. f is one-to-one on X ,
 4. g is the inverse of $f|_X$,
 5. g is C^1 .

Proof. Since f is C^1 , $\exists r_1 \in \mathbb{R}_{++}$ such that $\forall x \in B(a, r_1)$, $\det Df(x) \neq 0$. Then, from Proposition 648, f is one-to-one on $B(a, r_1)$. Then take $r_2 \in (0, r_1)$, and define $B := B(a, r_2)$. Observe that $\text{Cl}(B)(a, r_2) \subseteq B(a, r_1)$. Using the fact that f is one-to-one on $B(a, r_1)$ and therefore on $B(a, r_2)$, we get that Assumption 3 in Proposition 645 is satisfied - while the other two are trivially satisfied. Then, $\exists \delta \in \mathbb{R}_{++}$ such that

$$f(B(a, r_2)) \supseteq B(f(a), \delta) := Y.$$

Define also

$$X := f^{-1}(Y) \cap B, \tag{15.6}$$

an open set because Y and B are open sets and f is continuous. Since f is one-to-one and continuous on the compact set $\text{Cl}(B)$, from Proposition 526, there exists a unique continuous inverse function $\hat{g} : f(\text{Cl}(B)) \rightarrow \text{Cl}(B)$ of $f|_{\text{Cl}(B)}$. From definition of Y ,

$$Y \subseteq f(B) \subseteq f(\text{Cl}(B)). \tag{15.7}$$

From definition of X ,

$$f(X) = Y \cap f(B) = Y.$$

Then, from Lemma 651,

$$g = \hat{g}_Y \text{ is the inverse of } f|_X.$$

The above shows conclusions 1-4 of the Proposition. (About conclusion 1, observe that $a \in f^{-1}(B(f(a), \delta)) \cap B(a, r_2) = X$ and $f(a) \in f(X) = Y$.)

We are then left with proving condition 5.

Following what said in the proof of Proposition 648, we can define

$$h : \mathbb{R}^{n^2} \rightarrow \mathbb{R}, \quad : (z^i)_{i=1}^n \mapsto \det \begin{bmatrix} Df_1(z^1) \\ \dots \\ Df_i(z^i) \\ \dots \\ Df_n(z^n) \end{bmatrix}.$$

and get that, from Assumption 2,

$$h(a, \dots, a, \dots, a) = \det Df(a) \neq 0,$$

and, see the proof of Proposition 648 for details,

$$\exists r \in \mathbb{R}_{++} \text{ such that } \forall i \in \{1, \dots, n\}, \forall z_i \in B(a, r), \quad h(z^1, \dots, z^i, \dots, z^n) \neq 0, \quad (15.8)$$

and trivially also

$$\exists r \in \mathbb{R}_{++} \text{ such that } \forall z \in B(a, r), \quad h(z, \dots, z, \dots, z) = \det Df(z) \neq 0. \quad (15.9)$$

Assuming, without loss of generality that we took $r_1 < r$, we have that

$$\text{Cl}(B) := \text{Cl}(B)(a, r_2) \subseteq B(a, r_1) \subseteq B(a, r).$$

Then $\forall z^1, \dots, z^n \in \text{Cl}(B)$, $h(z^1, \dots, z^i, \dots, z^n) \neq 0$. Writing $g = (g^j)_{j=1}^n$, we want to prove that $\forall i \in \{1, \dots, n\}$, g^i is C^1 . We go through the following two steps: 1. $\forall y \in Y$, $\forall i, k \in \{1, \dots, n\}$, $D_{y_k} g^i(y)$ exists, and 2. it is continuous.

Step 1.

We want to show that the following limit exists and it is finite:

$$\lim_{h \rightarrow 0} \frac{g^i(y + he_n^k) - g^i(y)}{h}.$$

Define

$$\begin{aligned} x &= (x_i)_{i=1}^n = g(y) \subseteq X \subseteq \text{Cl}(B) \\ x' &= (x'_i)_{i=1}^n = g(y + he_n^k) \subseteq X \subseteq \text{Cl}(B) \end{aligned} \quad (15.10)$$

Then

$$f(x') - f(x) = (y + he_n^k) - y = he_n^k.$$

We can now apply the Mean Value Theorem to f^i for $i \in \{1, \dots, n\}$: $\exists z^i \in L(x, x') \subseteq \text{Cl}(B)$, where the inclusion follows from the fact that $x, x' \in \text{Cl}(B)$ a convex set, such that

$$\forall i \in \{1, \dots, n\}, \quad \frac{f^i(x') - f^i(x)}{h} = \frac{Df^i(z^i)(x' - x)}{h}$$

and therefore

$$\begin{bmatrix} Df_1(z^1) \\ \dots \\ Df_i(z^i) \\ \dots \\ Df_n(z^n) \end{bmatrix} \frac{1}{h} (x' - x) = e_n^k.$$

Define

$$A = \begin{bmatrix} Df_1(z^1) \\ \dots \\ Df_i(z^i) \\ \dots \\ Df_n(z^n) \end{bmatrix}$$

Then, from (15.8), the above system admits a unique solution, i.e., using (15.10),

$$\frac{g(y + he_n^k) - g(y)}{h} = \frac{1}{h} (x' - x)$$

and, using Cramer theorem (i.e., Theorem 328),

$$\frac{g(y + he_n^k) - g(y)}{h} = \frac{\varphi(z^1, \dots, z^n)}{h(z^1, \dots, z^n)}$$

where φ takes values which are determinants of a matrix involving entries of A . We are left with showing that

$$\lim_{h \rightarrow 0} \frac{\varphi(z^1, \dots, z^n)}{h(z^1, \dots, z^n)}$$

exists and it is finite, i.e., the limit of the numerator exists and its finite and the limit of the denominator exists is finite and nonzero.

Then, if $h \rightarrow 0$, $y + he_n^k \rightarrow y$, and, being g continuous, $x' \rightarrow x$ and, since $z^i \in L(x, x')$, $z^i \rightarrow x$ for any i . Then, $h(z^1, \dots, z^n) \rightarrow h(x, \dots, x) \neq 0$, because, from 15.10, $x \in \text{Cl}(B)$ and from (15.9). Moreover, $\varphi(z^1, \dots, z^n) \rightarrow \varphi(x, \dots, x)$.

Step 2.

Since

$$\lim_{h \rightarrow 0} \frac{g^i(y + he_n^k) - g^i(y)}{h} = \frac{\varphi(x, \dots, x)}{h(x, \dots, x)}$$

and φ and h are continuous functions, the desired result follows. ■

15.4 The implicit function theorem

Theorem 653 Given S, T open subsets of \mathbb{R}^n and \mathbb{R}^k respectively and a function

$$f : S \times T \rightarrow \mathbb{R}^n, \quad (x, t) \mapsto f(x, t),$$

assume that

1. f is C^1 ,
there exists $(x_0, t_0) \in S \times T$ such that
2. $f(x_0, t_0) = 0$,
3. $D_x f(x_0, t_0)_{n \times n}$ is invertible.

Then there exist $N(x_0) \subseteq S$ open neighborhood of x_0 , $N(t_0) \subseteq T$ open neighborhood of t_0 and a unique function

$$g : N(t_0) \rightarrow N(x_0)$$

such that

1. g is C^1 ,
2. $\{(x, t) \in N(x_0) \times N(t_0) : f(x, t) = 0\} = \{(x, t) \in N(x_0) \times N(t_0) : x = g(t)\} := \text{graph } g$.⁴

Proof. See Apostol (1974). proof to be added. ■

Remark 654 Conclusion 2. above can be rewritten as

$$\forall t \in N(t_0), \quad f(g(t), t) = 0 \tag{15.11}$$

Computing the Jacobian of both sides of (15.11), using Remark 626, we get

$$\forall t \in N(t_0), \quad 0 = [D_x f(g(t), t)]_{n \times n} \cdot [Dg(t)]_{n \times k} + [D_t f(g(t), t)]_{n \times k} \tag{15.12}$$

and using Assumption 3 of the Implicit Function Theorem, we get

$$\forall t \in N(t_0), \quad [Dg(t)]_{n \times k} = -[D_x f(g(t), t)]_{n \times n}^{-1} \cdot [D_t f(g(t), t)]_{n \times k}$$

Observe that (15.12) can be rewritten as the following k systems of equations: $\forall i \in \{1, \dots, k\}$,

$$[D_x f(g(t), t)]_{n \times n} \cdot [D_{t_i} g(t)]_{n \times 1} = -[D_{t_i} f(g(t), t)]_{n \times 1}$$

⁴Then $g(t_0) = x_0$.

Exercise 655 ⁵ Discuss the application of the Implicit Function Theorem to $f : \mathbb{R}^5 \rightarrow \mathbb{R}^2$

$$f(x_1, x_2, t_1, t_2, t_3) \mapsto \begin{pmatrix} 2e^{x_1} + x_2 t_1 - 4t_2 + 3 \\ x_2 \cos x_1 - 6x_1 + 2t_1 - t_3 \end{pmatrix}$$

at $(x^0, t^0) = (0, 1, 3, 2, 7)$.

Let's check that each assumption of the Theorem is verified.

1. $f(x_0, t_0) = 0$. Obvious.

2. f is C^1 .

We have to compute the Jacobian of the function and check that each entry is a continuous function.

	x_1	x_2	t_1	t_2	t_3
$2e^{x_1} + x_2 t_1 - 4t_2 + 3$	$2e^{x_1}$	t_1	x_2	-4	0
$x_2 \cos x_1 - 6x_1 + 2t_1 - t_3$	$-x_2 \sin x_1 - 6$	$\cos x_1$	2	0	-1

3. $[D_x f(x_0, t_0)]_{n \times n}$ is invertible.

$$[D_x f(x_0, t_0)] = \begin{bmatrix} 2e^{x_1} & t_1 \\ -x_2 \sin x_1 - 6 & \cos x_1 \end{bmatrix}_{|(0,1,3,2,7)} = \begin{bmatrix} 2 & 3 \\ -6 & 1 \end{bmatrix}$$

whose determinant is 20.


Therefore, we can apply the Implicit Function Theorem and compute the Jacobian of $g : N(t_0) \subseteq \mathbb{R}^2 \rightarrow N(x_0) \subseteq \mathbb{R}^3$:

$$\begin{aligned} Dg(t) &= - \begin{bmatrix} 2e^{x_1} & t_1 \\ -x_2 \sin x_1 - 6 & \cos x_1 \end{bmatrix}^{-1} \begin{bmatrix} x_2 & -4 & 0 \\ 2 & 0 & -1 \end{bmatrix} = \\ &= \frac{1}{6t_1 + 2(\cos x_1)e^{x_1} + t_1 x_2 \sin x_1} \begin{bmatrix} 2t_1 - x_2 \cos x_1 & 4 \cos x_1 & -t_1 \\ -6x_2 - 4e^{x_1} - x_2^2 \sin x_1 & 4x_2 \sin x_1 + 24 & 2e^{x_1} \end{bmatrix} \end{aligned}$$

Exercise 656 Given the utility function $u : \mathbb{R}_{++}^2 \rightarrow \mathbb{R}_{++}$, $(x, y) \mapsto u(x, y)$ satisfying the following properties

i. u is C^2 , ii. $\forall (x, y) \in \mathbb{R}_{++}^2$, $Du(x, y) > 0$, iii. $\forall (x, y) \in \mathbb{R}_{++}^2$, $D_{xx}u(x, y) < 0$, $D_{yy}u(x, y) < 0$, $D_{xy}u(x, y) > 0$,

compute the Marginal Rate of Substitution in (x_0, y_0) and say if the graph of each indifference curve is concave.



⁵The example is taken from Rudin (1976), pages 227-228.

15.5 Some geometrical remarks on the gradient

In what follows we make some geometrical, not rigorous remarks on the meaning of the gradient, using the implicit function theorem. Consider an open subset X of \mathbb{R}^2 , a C^1 function

$$f : X \rightarrow \mathbb{R}, \quad (x, y) \mapsto f(x, y)$$

where $a \in \mathbb{R}$. Assume that set

$$L(a) := \{(x, y) \in X : f(x, y) = a\}$$

is such that $\forall (x, y) \in X$, $\frac{\partial f(x, y)}{\partial y} \neq 0$ and $\frac{\partial f(x, y)}{\partial x} \neq 0$, then

1. $L(a)$ is the graph of a C^1 function from a subset of \mathbb{R} to \mathbb{R} ;
 2. $(x^*, y^*) \in L(a) \Rightarrow$ the line going through the origin and the point $Df(x^*, y^*)$ is orthogonal to the line going through the origin and parallel to the tangent line to $L(a)$ at (x^*, y^*) ; or the line tangent to the curve $L(a)$ in (x^*, y^*) is orthogonal to the line to which the gradient belongs to.
 3. $(x^*, y^*) \in L(a) \Rightarrow$ the directional derivative of f at (x^*, y^*) in the the direction u such that $\|u\| = 1$ is the largest one if $u = \frac{Df(x^*, y^*)}{\|Df(x^*, y^*)\|}$.
1. It follows from the Implicit Function Theorem.
 2. The slope of the line going through the origin and the vector $Df(x^*, y^*)$ is

$$\frac{\frac{\partial f(x^*, y^*)}{\partial y}}{\frac{\partial f(x^*, y^*)}{\partial x}} \quad (15.13)$$

Again from the Implicit Function Theorem, the slope of the tangent line to $L(a)$ in (x^*, y^*) is

$$-\frac{\frac{\partial f(x^*, y^*)}{\partial x}}{\frac{\partial f(x^*, y^*)}{\partial y}} \quad (15.14)$$

The product between the expressions in (15.13) and (15.14) is equal to -1 .

3. the directional derivative of f at (x^*, y^*) in the the direction u is

$$f'((x^*, y^*); u) = Df(x^*, y^*) \cdot u = \|Df(x^*, y^*)\| \cdot \|u\| \cdot \cos \theta$$

where θ is an angle in between the two vectors. Then the above quantity is the greatest possible iff $\cos \theta = 1$, i.e., u is colinear with $Df(x^*, y^*)$, i.e., $u = \frac{Df(x^*, y^*)}{\|Df(x^*, y^*)\|}$.

15.6 Extremum problems with equality constraints.

Given the open set $X \subseteq \mathbb{R}^n$, consider the C^1 functions

$$f : X \rightarrow \mathbb{R}, \quad f : x \mapsto f(x),$$

$$g : X \rightarrow \mathbb{R}^m, \quad g : x \mapsto g(x) := (g_j(x))_{j=1}^m$$

with $m \leq n$. Consider also the following “maximization problem”:

$$(P) \quad \max_{x \in X} f(x) \quad \text{subject to} \quad g(x) = 0 \quad (15.15)$$

The set

$$C := \{x \in X : g(x) = 0\}$$

is called the constraint set associated with problem (15.15).

Definition 657 The solution set to problem (15.15) is the set

$$\{x^* \in C : \forall x \in C, f(x^*) \geq f(x)\},$$

and it is denoted by $\arg \max$ (15.15).

The function

$$\mathcal{L} : X \times \mathbb{R}^m \rightarrow \mathbb{R}, \quad \mathcal{L} : (x, \lambda) \mapsto f(x) + \lambda^T g(x)$$

is called Lagrange function associated with problem (15.15).

Theorem 658 Given the open set $X \subseteq \mathbb{R}^n$ and the C^1 functions

$$f : X \rightarrow \mathbb{R}, \quad f : x \mapsto f(x), \quad g : X \rightarrow \mathbb{R}^m, \quad g : x \mapsto g(x) := (g_j(x))_{j=1}^m,$$

assume that

1. f and g are C^1 functions,
 2. x_0 is a solution to problem (15.15),⁶ and
 3. $\text{rank}[Dg(x_0)]_{m \times n} = m$.
- Then, there exists $\lambda_0 \in \mathbb{R}^m$, such that, $D\mathcal{L}(x_0, \lambda_0) = 0$, i.e.,

$$\begin{cases} Df(x_0) + \lambda_0 Dg(x_0) = 0 \\ g(x_0) = 0 \end{cases} \quad (15.16)$$

Proof. Define $x' := (x_i)_{i=1}^m \in \mathbb{R}^m$ and $t = (x_{m+k})_{k=1}^{n-m} \in \mathbb{R}^{n-m}$ and therefore $x = (x', t)$. From Assumption 3, without loss of generality,

$$\det[D_{x'}g(x_0)]_{m \times m} \neq 0. \quad (15.17)$$

We want to show that there exists $\lambda_0 \in \mathbb{R}^m$ which is a solution to the system

$$[Df(x_0)]_{1 \times n} + \lambda_{1 \times m} [Dg(x_0)]_{m \times n} = 0. \quad (15.18)$$

We can rewrite (15.18) as follows

$$\begin{bmatrix} D_{x'}f(x_0)_{1 \times m} & | & D_tf(x_0)_{1 \times (n-m)} \end{bmatrix} + \lambda_{1 \times m} \begin{bmatrix} D_{x'}g(x_0)_{m \times m} & | & Dtg(x_0)_{m \times (n-m)} \end{bmatrix} = 0$$

or

$$\begin{cases} \begin{bmatrix} D_{x'}f(x_0)_{1 \times m} \end{bmatrix} + \lambda_{1 \times m} \begin{bmatrix} D_{x'}g(x_0)_{m \times m} \end{bmatrix} = 0 & (1) \\ \begin{bmatrix} D_tf(x_0)_{1 \times (n-m)} \end{bmatrix} + \lambda_{1 \times m} \begin{bmatrix} Dtg(x_0)_{m \times (n-m)} \end{bmatrix} = 0 & (2) \end{cases} \quad (15.19)$$

From (15.17), there exists a unique solution λ_0 to subsystem (1) in (15.19). If $n = m$, we are done. Assume now that $n > m$. We have now to verify that λ_0 is a solution to subsystem (2) in (15.19), as well. To get the desired result, we are going to use the Implicit Function Theorem. Summarizing, we have that

$$1. \ g \text{ is } C^1, \quad 2. \ g(x'_0, t_0) = 0, \quad 3. \ \det[D_{x'}g(x'_0, t_0)]_{m \times m} \neq 0,$$

i.e., all the assumption of the Implicit Function Theorem are verified. Then we can conclude that there exist $N(x_0) \subseteq \mathbb{R}^m$ open neighborhood of x'_0 , $N(t_0) \subseteq \mathbb{R}^{n-m}$ open neighborhood of t_0 and a unique function $\varphi : N(t_0) \rightarrow N(x_0)$ such that

$$1. \ \varphi \text{ is } C^1, \quad 2. \ \varphi(t_0) = x'_0, \quad 3. \ \forall t \in N(t_0), \quad g(\varphi(t), t) = 0. \quad (15.20)$$

Define now

$$F : N(t_0) \subseteq \mathbb{R}^{n-m} \rightarrow \mathbb{R}, \quad : t \mapsto f(\varphi(t), t),$$

and

$$G : N(t_0) \subseteq \mathbb{R}^{n-m} \rightarrow \mathbb{R}^m, \quad : t \mapsto g(\varphi(t), t).$$

⁶The result does apply in the case in which x_0 is a local maximum for Problem (15.15). Obviously the result apply to the case of (local) minima, as well.

Then, from (15.20) and from Remark 654, we have that $\forall t \in N(t_0)$,

$$0 = [DG(t)]_{m \times (n-m)} = [D_{x'}g(\varphi(t), t)]_{m \times m} \cdot [D\varphi(t)]_{m \times (n-m)} + [D_tg(\varphi(t), t)]_{m \times (n-m)}. \quad (15.21)$$

Since⁷, from (15.20), $\forall t \in N(t_0)$, $g(\varphi(t), t) = 0$ and since

$$x_0 := (x'_0, t_0) \stackrel{(15.20)}{=} (\varphi(t_0), t_0) \quad (15.22)$$

is a solution to problem (15.15), we have that $f(x_0) = F(t_0) \geq F(t)$, i.e., briefly,

$$\forall t \in N(t_0), \quad F(t_0) \geq F(t).$$

Then, from Proposition 638, $DF(t_0) = 0$. Then, from the definition of F and the Chain Rule, we have

$$[D_{x'}f(\varphi(t_0), t_0)]_{1 \times m} \cdot [D\varphi(t_0)]_{m \times (n-m)} + [D_tf(\varphi(t_0), t_0)]_{1 \times (n-m)} = 0. \quad (15.23)$$

Premultiplying (15.21) by λ , we get

$$\lambda_{1 \times m} \cdot [D_{x'}g(\varphi(t), t)]_{m \times m} \cdot [D\varphi(t)]_{m \times (n-m)} + \lambda_{1 \times m} \cdot [D_tg(\varphi(t), t)]_{m \times (n-m)} = 0. \quad (15.24)$$

Adding up (15.23) and (15.24), computed at $t = t_0$, we get

$$([D_{x'}f(\varphi(t_0), t_0)] + \lambda \cdot [D_{x'}g(\varphi(t_0), t_0)]) \cdot [D\varphi(t_0)] + [D_tf(\varphi(t_0), t_0)] + \lambda \cdot [D_tg(\varphi(t_0), t_0)] = 0,$$

and from (15.22),

$$([D_{x'}f(x_0)] + \lambda \cdot [D_{x'}g(x_0)]) \cdot [D\varphi(t_0)] + [D_tf(x_0)] + \lambda \cdot [D_tg(x_0)] = 0. \quad (15.25)$$

Then, from the definition of λ_0 as the unique solution to (1) in (15.19), we have that $[D_{x'}f(x_0)] + \lambda_0 \cdot [D_{x'}g(x_0)] = 0$, and then from (15.25) computed at $\lambda = \lambda_0$, we have

$$[D_tf(x_0)] + \lambda_0 \cdot [D_tg(x_0)] = 0,$$

i.e., (2) in (15.19), the desired result. ■

15.7 Exercises on part III

Problem sets: all of the problems on part III.

See Tito Pietra's file (available on line): Exercises $1 \rightarrow 14$ (excluding exercises 3, 5, 15).

⁷The only place where the proof has to be slightly changed to get the result for local maxima is here.

Part IV

Nonlinear programming

Chapter 16

Convex sets

16.1 Definition

Definition 659 A set $C \subseteq \mathbb{R}^n$ is convex if $\forall x_1, x_2 \in C$ and $\forall \lambda \in [0, 1]$, $(1 - \lambda)x_1 + \lambda x_2 \in C$.

Definition 660 A set $C \subseteq \mathbb{R}^n$ is strictly convex if $\forall x_1, x_2 \in C$ such that $x_1 \neq x_2$, and $\forall \lambda \in (0, 1)$, $(1 - \lambda)x_1 + \lambda x_2 \in \text{Int } C$.

Remark 661 If C is strictly convex, then C is convex, but not vice-versa.

Proposition 662 The intersection of an arbitrary family of convex sets is convex.

Proof. We want to show that given a family $\{C_i\}_{i \in I}$ of convex sets, if $x, y \in C := \bigcap_{i \in I} C_i$ then $(1 - \lambda)x + \lambda y \in C$. $x, y \in C$ implies that $x, y \in C_i$, $\forall i \in I$. Since C_i is convex, $\forall i \in I$, $\forall \lambda \in [0, 1]$, $(1 - \lambda)x + \lambda y \in C_i$, and $\forall \lambda \in [0, 1]$ $(1 - \lambda)x + \lambda y \in C$. ■

Exercise 663 $\forall n \in \mathbb{N}, \forall i \in \{1, \dots, n\}$, I_i is an interval in \mathbb{R} , then

$$\times_{i=1}^n I_i$$

is a convex set.

16.2 Separation of convex sets

Definition 664 Let A and B be subsets of \mathbb{R}^n , and let $H := \{x \in \mathbb{R}^n : c_0 + c \cdot x = 0\}$ be a hyperplane in \mathbb{R}^n .¹ Then, H is said to

1. separate A and B if

$$A \subseteq H_- := \{x \in \mathbb{R}^n : c_0 + c \cdot x \leq 0\} \quad \text{and} \quad B \subseteq H_+ := \{x \in \mathbb{R}^n : c_0 + c \cdot x \geq 0\},$$

i.e.,

$$\forall a \in A, \quad \forall b \in B, \quad c_0 + c \cdot a \leq 0 \leq c_0 + c \cdot b,$$

i.e.,

$$\forall a \in A, \quad \forall b \in B, \quad c \cdot a \leq -c_0 \leq c \cdot b;$$

2. separate A and B properly if it separates them and $A \cup B \not\subseteq H$;

3. separate A and B strictly if

$$A \subseteq \{x \in \mathbb{R}^n : c_0 + c \cdot x < 0\} \quad \text{and} \quad B \subseteq \{x \in \mathbb{R}^n : c_0 + c \cdot x > 0\}.$$

¹When we introduce a hyperplane by equation $c_0 + cx = 0$, consistently with the definition of hyperplane, we assume $c \neq 0$.

Example 665 The convex sets $\{(x, y) \in \mathbb{R}^2 : x \leq 0\}$ and $\{(x, y) \in \mathbb{R}^2 : x > 0, y > \frac{1}{x}\}$ in \mathbb{R}^2 cannot be strictly separated, but they are properly separated by the y -axis.

Clearly, it is not always possible to separate two convex sets by a hyperplane. For instance, there is no line in \mathbb{R}^2 separating the set $\{0\}$ and the closed unit disc $\{(x, y) \in \mathbb{R}^2 : x^2 + y^2 \leq 1\}$.

Remark 666 Let A, B be two sets in \mathbb{R}^n such that at least one of them is nonempty. If they can be strictly separated, then they can also be properly separated.

Remark 667 Let H be a hyperplane in \mathbb{R}^n , and let A and B be two subsets of H . Then, H separates A and B , but does not separate them properly.

Proposition 668 Let a hyperplane $H := \{x \in \mathbb{R}^n : c_0 + c \cdot x = 0\}$ be given. Then, 2.

$$\langle H \text{ separates properly } A \text{ and } B \rangle \Leftrightarrow \langle H \text{ separates properly } \text{Cl}(A) \text{ and } \text{Cl}(B) \rangle.$$

Proof. [\Leftarrow]

Obvious.

[\Rightarrow]

We first present two proofs of the fact H separates $\text{Cl}(A)$ and $\text{Cl}(B)$, and then we show that the separation is proper.

1st proof.

$$\langle A \subseteq H_- \rangle \Rightarrow \langle \text{Cl}(A) \subseteq \text{Cl}(H_-) = H_- \rangle,$$

where we used the fact that H_- is closed.

Similarly, $B \subseteq H_+ \Rightarrow \text{Cl}(H_+) \subseteq H_+$

2nd proof.

Take $(a^*, b^*) \in \text{Cl}(A) \times \text{Cl}(B)$. Then there exists sequences $(a_n)_{n \in \mathbb{N}} \in A^\infty$ and $(b_n)_{n \in \mathbb{N}} \in B^\infty$ such that $a_n \rightarrow a^*$ and $b_n \rightarrow b^*$. By assumption,

$$\forall n \in \mathbb{N}, \quad c_0 + c \cdot a_n \leq 0 \leq c_0 + c \cdot b_n.$$

Taking limits for $n \rightarrow +\infty$, we get

$$c_0 + c \cdot a^* \leq 0 \leq c_0 + c \cdot b^*,$$

as desired.

We now show that the separation is proper;

$$A \cup B \not\subseteq H \Rightarrow A \cup B \cap H^C \neq \emptyset \Rightarrow \text{Cl}(A) \cup \text{Cl}(B) \cap H^C \neq \emptyset.$$

■

The following three Propositions are presented without proof. Detailed, self-contained proofs of those results are contained, for example, in Villanacci, A., (in progress), Basic Convex Analysis, mimeo, Università degli Studi di Firenze.

Proposition 669 Let A be a closed nonempty convex set in \mathbb{R}^n such that $0 \notin A$. Then, there exists a hyperplane in \mathbb{R}^n that strictly separates A and $\{0\}$.

Corollary 670 Let B be a closed nonempty convex set in \mathbb{R}^n such that $w \notin B$. Then, there exists a hyperplane in \mathbb{R}^n that strictly separates B and $\{w\}$.

Proof. Exercise. ■

Proposition 671 Let A be a nonempty convex set in \mathbb{R}^n such that $0 \notin A$. Then, there exists a hyperplane H in \mathbb{R}^n such that $A \subseteq H_-$ and $\{0\} \subseteq H_+$.

Proposition 672 Let A be a nonempty convex set in \mathbb{R}^n such that $0 \notin \text{Int}(A)$. Then, there exists a hyperplane H in \mathbb{R}^n such that $A \subseteq H_-$ and $\{0\} \subseteq H_+$.

Proposition 673 Let A and B be nonempty convex sets in \mathbb{R}^n . If one of the following conditions holds, then there exists a hyperplane H such that $A \subseteq H_-$ and $B \subseteq H_+$:

1. $0 \notin A - B$;
2. $0 \notin \text{Int}(A - B)$;
3. $\text{Int}B \neq \emptyset$ and $0 \notin A - \text{Int}(B)$.

Proof. 1.

Since A and B are convex, we show that $A - B$ is convex. Let $x, y \in A - B$ where $x = a_1 - b_1$, $y = a_2 - b_2$, and $a_1, a_2 \in A, b_1, b_2 \in B$. Let $\lambda \in [0, 1]$. Then,

$$(1 - \lambda)x + \lambda y = (1 - \lambda)(a_1 - b_1) + \lambda(a_2 - b_2) = [(1 - \lambda)a_1 + \lambda a_2] - [(1 - \lambda)b_1 + \lambda b_2] \in A - B$$

by convexity of A and B .

Hence, from our assumption and Lemma ??, there exists a hyperplane $H = \{x \in \mathbb{R}^n : c \cdot x = c_0\}$ that separates $\{0\}$ and $A - B$. Without loss of generality, $A - B \subseteq H_+$ and $c_0 = 0$. Then, $\forall a \in A, b \in B$,

$$c \cdot (a - b) \geq 0 \Leftrightarrow c \cdot a \geq c \cdot b.$$

Hence, there must exist $H' = \{x' \in \mathbb{R}^n : c \cdot x' = c_1\}$ such that

$$c \cdot a \geq c_1 \geq c \cdot b,$$

i.e., such that $B \subseteq H'_-$ and $A \subseteq H'_+$.

2.

From our assumption and 672, there exists a hyperplane that separates $\{0\}$ and $A - B$, whence, following the Proof of point 1 above, we are done.

3.

From our assumption and point 1 above, there exists a hyperplane H that separates A and $\text{Int}(B)$. From Remark 668, H separates $\text{Cl}(A)$ and $\text{Cl}(\text{Int}(B)) = \text{Cl}(B)$, where last equality follows from the Assumption that $\text{Int}(B) \neq \emptyset$ and Proposition 662.7. Then since the closure of a set contains the set, the desired result follows. ■

Proposition 674 *Let A and B be subsets of \mathbb{R}^n . Then*

$$0 \notin A - B \Leftrightarrow A \cap B = \emptyset.$$

Proof.

$$0 \notin A - B \Leftrightarrow$$

$$\neg(0 \in A - B) \Leftrightarrow$$

$$\neg(\exists a \in A, \exists b \in B \text{ such that } a - b = 0) \Leftrightarrow$$

$$\forall a \in A, \forall b \in B, \quad a \neq b \Leftrightarrow$$

$$A \cap B = \emptyset.$$

■

Proposition 675 *Let A and B be nonempty convex sets in \mathbb{R}^n . If one of the following conditions holds true, then there exists a hyperplane H such that $A \subseteq H_-$ and $B \subseteq H_+$.*

1. $A \cap B = \emptyset$;
2. $\text{Int}B \neq \emptyset$ and $A \cap \text{Int}(B) = \emptyset$.

16.3 Farkas' Lemma

Proposition 676 *\mathbb{R}^2*

1. $v_1, \dots, v_m, w \in \mathbb{R}^n$, and
 2. $v_1 x \leq 0, \dots, v_m x \leq 0 \Rightarrow w x \leq 0$,
- then

$$\exists \lambda := (\lambda_i)_{i=1}^m \in \mathbb{R}_+^m \quad \text{such that} \quad w = \sum_{i=1}^m \lambda_i v_i.$$

²In this Section, I follow very closely Section 8.1.2 in Montrucchio (1998).

Proof. Define

$$C = \left\{ y \in \mathbb{R}^n : \sum_{i=1}^m \gamma_i v_i \text{ with } (\gamma_i)_{i=1}^m \in \mathbb{R}_+^m \right\}.$$

Observe that since $0 \in \mathbb{R}_+^m$, then

$$0 \in C, \quad (16.1)$$

and

$$\forall \lambda \in \mathbb{R}_+ \text{ and } \forall y \in C, \text{ we have } \lambda y \in C. \quad (16.2)$$

We want to show that $w \in C$.

Claim 1. C is a convex, nonempty and closed set.

Proof of Claim 1.

The fact that C is a convex and nonempty is obvious. Let's check closedness. Take $(x_k)_{k \in \mathbb{N}} \in C^\infty$ such that $x_k \rightarrow x$. We want to show that $x \in C$. Indeed, $x_k \in C$ means that there exists $(\gamma_{ik})_{i=1}^m$ such that

$$x_k = \sum_{i=1}^m \gamma_{ik} v_i \rightarrow x. \quad (16.3)$$

Then

$$\forall \varepsilon > 0 \quad \exists N_\varepsilon \in \mathbb{N} \text{ such that } \forall k > N_\varepsilon, \quad \left\| \sum_{i=1}^m \gamma_{ik} v_i - x \right\| < \varepsilon.$$

Now, in general, we have that $\|x\| = \|x - y + y\| \leq \|x - y\| + \|y\|$, i.e.,

$$\|x\| - \|y\| \leq \|x - y\|.$$

Therefore,

$$\left\| \sum_{i=1}^m \gamma_{ik} v_i \right\| - \|x\| \leq \left\| \sum_{i=1}^m \gamma_{ik} v_i - x \right\| < \varepsilon, \quad (16.4)$$

and

$$0 \leq \left\| \sum_{i=1}^m \gamma_{ik} v_i \right\| < \|x\| + \varepsilon. \quad (16.5)$$

Then, as verified below, $((\gamma_{ik})_{i=1}^m)_{k \in \mathbb{N}}$ is a bounded sequence. Suppose otherwise; then, since by assumption, for any $i \in \{1, \dots, m\}$ and any $k \in \mathbb{N}$, $\gamma_{ik} \geq 0$, we have that there exists $i^* \in \{1, \dots, m\}$ such that $\gamma_{ik^*} \rightarrow +\infty$. Then,

$$\lim_{k \rightarrow +\infty} \left\| \sum_{i=1}^m \gamma_{ik} v_i \right\| = \lim_{k \rightarrow +\infty} \sum_{i=1}^m \gamma_{ik} \|v_i\| = +\infty,$$

violating (16.5).

Then, up to a subsequence,

$$\gamma_k := (\gamma_{ik})_{i=1}^m \rightarrow \tilde{\gamma} \in \mathbb{R}_+^m. \quad (16.6)$$

Then,

$$\sum_{i=1}^m \gamma_{ik} v_i \rightarrow \sum_{i=1}^m \tilde{\gamma}_i v_i \stackrel{(16.3)}{=} x,$$

and, then, from (16.6), $x \in C$, as desired.

End of the Proof of Claim 1.

Suppose now our Claim is false, i.e., $w \notin C$. We can then apply Corollary 669 to conclude that there exist $a \in \mathbb{R}^n \setminus \{0\}$ and $\gamma \in \mathbb{R}$ such that

$$\forall x \in C, \quad aw > \gamma > ax. \quad (16.7)$$

From (16.1), $0 \in C$ and then from (16.7), we have

$$\forall x \in C, \quad aw > \gamma > 0. \quad (16.8)$$

Claim 2.

$$\forall x \in C, \quad ax \leq 0.$$

Proof of Claim 2.

Suppose otherwise, i.e., $\exists x^* \in C$ such that

$$ax^* > 0.$$

From (16.2), and the fact that $x^* \in C$, we have that for any $\lambda \geq 0$, $\lambda x^* \in C$. Then, from (16.7), we get

$$\forall \lambda \geq 0, \quad \gamma > a\lambda x^* = \overset{(\geq 0)}{\lambda} \overset{(>0)}{(ax^*)} > 0,$$

which is clearly false: if $\lambda = 2\frac{\gamma}{ax^*} > 0$, we get $\gamma > 2\frac{\gamma}{ax^*}ax^*$, i.e., $1 > 2$.

End of the Proof of Claim 2.

Summarizing, we have shown that

$$\exists a \in \mathbb{R}^n \setminus \{0\} \text{ such that } \forall x \in C, \quad ax \leq 0 \text{ and } aw > 0.$$

Since $v_1, v_2, \dots, v_m \in C$, we have that

$$av_1 \leq 0, \dots, av_m \leq 0 \text{ and } aw > 0,$$

contradicting the assumption. ■

Proposition 677 (*A version of Farkas Lemma*) Given $A \in \mathbb{M}(m, n)$, $b \in \mathbb{R}^n$,

- either 1. $\exists x \in \mathbb{R}^n$ such that $Ax \leq 0$ and $bx > 0$,
or 2. $\exists y \in \mathbb{R}^m$ such that $yA = b$ and $y \geq 0$,
but not both.

Proof. We want to show that either

1.

$$\begin{cases} Ax \leq 0 \\ bx > 0 \end{cases} \quad (16.9)$$

has a solution, or

2.

$$\begin{cases} yA = b \\ y \geq 0 \end{cases} \quad (16.10)$$

has a solution, but not both.

Claim. It suffices to show

a. $(\neg 1) \Rightarrow 2$,

b. $2 \Rightarrow (\neg 1)$.

Proof of the Claim.

Indeed, a. $\Leftrightarrow (\neg 2) \Rightarrow 1$ and b. $\Leftrightarrow 1 \Rightarrow (\neg 2)$. Therefore, showing a. and b. implies showing

$$(\neg 1) \Leftrightarrow 2, \quad \text{and} \quad (\neg 2) \Leftrightarrow 1. \quad (16.11)$$

Observe also that we have that $1 \vee 2$. Suppose otherwise, i.e., $(\neg 1) \wedge (\neg 2)$. But $(\neg 1)$ and (16.11) imply 2, a contradiction.

End of the proof of the Claim.

We are now left with showing a. and b.

a. Suppose that (16.9) has a no solution. Then $Ax \leq 0$ implies that $bx \leq 0$. Then from Proposition 676, identifying v_i with $R^i(A)$, the i -th row of A , and w with b , we have that

$$\exists \lambda := (\lambda_i)_{i=1}^m \in \mathbb{R}_+^m \quad \text{such that} \quad b = \lambda A,$$

i.e., λ is a solution to (16.10).

b. By assumption, there exists $y \in \mathbb{R}_+^m$ such that $yA = b$. Then, taken $x \in \mathbb{R}^n$, we also have $yAx = bx$. Now if $bx > 0$, since $y \geq 0$, we have that $\exists i \in \{1, \dots, m\}$ such that $R^i(A) \cdot x > 0$, and therefore (16.9) has no solution. ■

Chapter 17

Concave functions

Consider¹ a set $X \subseteq \mathbb{R}^n$, a set $\Pi \subseteq \mathbb{R}^k$ and the functions $f : X \times \Pi \rightarrow \mathbb{R}$, $g : X \times \Pi \rightarrow \mathbb{R}^m$, $h : X \times \Pi \rightarrow \mathbb{R}^l$. The goal of this Chapter is to study the problem:

for given f, g, h and for given $\pi \in \Pi$,

$$\max_{x \in X} f(x, \pi) \quad \text{s.t.} \quad g(x, \pi) \geq 0 \quad \text{and} \quad h(x, \pi) = 0,$$

under suitable assumptions. The role of concavity (and differentiability) of the functions f, g and h is crucial.

In what follows, unless needed, we omit the depends on π .

17.1 Different Kinds of Concave Functions

Maintained Assumptions in this Chapter. Unless otherwise stated,

X is an open and convex subset of \mathbb{R}^n .

f is a function such that

$$f : X \rightarrow \mathbb{R}, \quad : x \mapsto f(x).$$

For each type of concavity we study, we present

1. the definition in the case in which f is C^0 (i.e., continuous),
2. an attempt of a “partial characterization” of that definition in the case in which f is C^1 and C^2 ; by partial characterization, we mean a statement which is either sufficient or necessary for the concept presented in the case of continuous f ;
3. the relationship between the different partial characterizations;
4. the relationship between the type of concavity and critical points and local or global extrema of f .

Finally, we study the relationship between different kinds of concavities.

The following pictures are taken from David Cass’s Microeconomic Course I followed at the University of Pennsylvania (in 1985) and summarize points 1., 2. and 3. above.

¹This part is based on Cass (1991).

Concave and Quasi-Concave Functions

Assume $X \subset \mathbb{R}^n$ is convex and open, $f: X \rightarrow \mathbb{R}$.

- f is concave if

f continuous

f differentiable

f twice differentiable


$$\{(y, x) \in \mathbb{R}^{n+1} : x \in X \text{ \& \& } y \leq f(x)\}$$

is convex



$$x', x'' \in X \text{ \& } 0 \leq \lambda \leq 1 \Rightarrow$$

$$f((1-\lambda)x' + \lambda x'') \geq (1-\lambda)f(x') + \lambda f(x'')$$




both "flats" and "kinks" are possible

$$x', x'' \in X \Rightarrow$$



$$f(x'') \leq f(x') + Df(x')(x'' - x')$$



"flats" but not "kinks" are possible

$$x \in X \text{ \& } \Delta x \in \mathbb{R}^n \Rightarrow$$



$$\Delta x^T D^2 f(x) \Delta x \leq 0$$

• f is strictly concave if it is concave and

f continuous

f differentiable

f twice differentiable

$$x \in X, \Delta x \in \mathbb{R}^n \text{ \& } \Delta x \neq 0 \Rightarrow$$


$$\Delta x^T D^2 f(x) \Delta x < 0$$

$$x', x'' \in X, x' \neq x'' \Rightarrow$$

$$f((1-\lambda)x' + \lambda x'') > (1-\lambda)f(x') + \lambda f(x'')$$


$$\Leftrightarrow$$

$$\Leftrightarrow$$



$$(1-\lambda)x' + \lambda x''$$

"kinks" but not "flats" are possible



neither "flats" nor "kinks" are possible

Note: When X is open (as well convex), (i) if f is concave, then it is continuous, while (ii) it is meaningful to assume, for example, that f is differentiable (i.e., that f has a first-order differential at each point in X).

f is quasi-concave if


f continuous

$\{x \in X: f(x) \geq \alpha\}$ is convex for every $\alpha \in \mathbb{R}$



$$x', x'' \in X \text{ \& } 0 \leq \lambda \leq 1 \Rightarrow$$

$$f((1-\lambda)x' + \lambda x'') \geq \min\{f(x'), f(x'')\}$$



f differentiable


f twice differentiable

$$x', x'' \in X \text{ \& } f(x'') - f(x') \geq 0 \Rightarrow$$

$$Df(x')(x'' - x') \geq 0$$



$$\Delta x^T D^2 f(x) \Delta x \leq 0$$



f is (at worst) "single-peaked", and may have "flats"

f is **strictly quasi-concave** if it is quasi-concave and

f continuous

f differentiable

f twice differentiable

$$x', x'' \in X, x' \neq x'' \text{ and } 0 < \lambda < 1 \Rightarrow$$



$$x \in X, \Delta x \in \mathbb{R}^n, \Delta x \neq 0 \text{ \&}$$

$$f(x'') - f(x') \geq 0 \Rightarrow$$

$$Df(x) \Delta x = 0 \Rightarrow$$

$$\{(1-\lambda)x' + \lambda x''\} > \min\{f(x'), f(x'')\}$$

$$\Delta x^T D^2 f(x) \Delta x < 0$$



f has no "flats"

Note: There are alternative possible definitions of strict quasi-concavity when f is differentiable. This is the definition most frequently adopted in economic applications.

17.1.1 Concave Functions.

Definition 678 Consider a C^0 function f . f is concave iff $\forall x', x'' \in X, \forall \lambda \in [0, 1]$,

$$f((1 - \lambda)x' + \lambda x'') \geq (1 - \lambda)f(x') + \lambda f(x'').$$

Proposition 679 Consider a C^0 function f .

f is concave

\Leftrightarrow

$M = \{(x, y) \in X \times \mathbb{R} : y \leq f(x)\}$ is convex.

Proof.

$[\Rightarrow]$

Take $(x', y'), (x'', y'') \in M$. We want to show that

$$\forall \lambda \in [0, 1], ((1 - \lambda)x' + \lambda x'', (1 - \lambda)y' + \lambda y'') \in M.$$

But, from the definition of M , we get that

$$(1 - \lambda)y' + \lambda y'' \leq (1 - \lambda)f(x') + \lambda f(x'') \leq f((1 - \lambda)x' + \lambda x'').$$

$[\Leftarrow]$

From the definition of M , $\forall x', x'' \in X, (x', f(x')) \in M$ and $(x'', f(x'')) \in M$.

Since M is convex,

$$((1 - \lambda)x' + \lambda x'', (1 - \lambda)f(x') + \lambda f(x'')) \in M$$

and from the definition of M ,

$$(1 - \lambda)f(x') + \lambda f(x'') \leq f((1 - \lambda)x' + \lambda x'')$$

as desired.

■

Proposition 680 (Some properties of concave functions).

1. If $f, g : X \rightarrow \mathbb{R}$ are concave functions and $a, b \in \mathbb{R}_+$, then the function $af + bg : X \rightarrow \mathbb{R}$, $af + bg : x \mapsto af(x) + bg(x)$ is a concave function.

2. If $f : X \rightarrow \mathbb{R}$ is a concave function and $F : A \rightarrow \mathbb{R}$, with $A \supseteq \text{Im } f$, is nondecreasing and concave, then $F \circ f$ is a concave function.

Proof.

1. This result follows by a direct application of the definition.
2. Let $x', x'' \in X$ and $\lambda \in [0, 1]$. Then

$$(F \circ f)((1 - \lambda)x' + \lambda x'') \stackrel{(1)}{\geq} F((1 - \lambda)f(x') + \lambda f(x'')) \stackrel{(2)}{\geq} (1 - \lambda) \cdot (F \circ f)(x') + \lambda \cdot (F \circ f)(x''),$$

where (1) comes from the fact that f is concave and F is non decreasing, and

(2) comes from the fact that F is concave.

■

Remark 681 (from Sydsæter (1981)). With the notation of part 2 of the above Proposition, the assumption that F is concave cannot be dropped, as the following example shows. Take $f, F : \mathbb{R}_{++} \rightarrow \mathbb{R}_{++}$, $f(x) = \sqrt{x}$ and $F(y) = y^3$. Then f is concave and F is strictly increasing, but $F \circ f(x) = x^{\frac{3}{2}}$ and its second derivative is $\frac{3}{4}x^{-\frac{1}{2}} > 0$. Then, from Calculus I, we know that $F \circ f$ is strictly convex and therefore it is not concave.

Of course, the monotonicity assumption cannot be dispensed either. Consider $f(x) = -x^2$ and $F(y) = -y$. Then, $(F \circ f)(x) = x^2$, which is not concave.

Proposition 682 Consider a differentiable function f .

f is concave

\Leftrightarrow

$$\forall x', x'' \in X, f(x'') - f(x') \leq Df(x')(x'' - x').$$

Proof.[\Rightarrow]

From the definition of concavity, we have that for $\lambda \in (0, 1)$,

$$(1 - \lambda) f(x') + \lambda f(x'') \leq f(x' + \lambda(x'' - x')) \quad \Rightarrow$$

$$\lambda(f(x'') - f(x')) \leq f(x' + \lambda(x'' - x')) - f(x') \quad \Rightarrow$$

$$f(x'') - f(x') \leq \frac{f(x' + \lambda(x'' - x')) - f(x')}{\lambda}.$$

Taking limits of both sides of the last inequality for $\lambda \rightarrow 0$, we get the desired result.

[\Leftarrow]

Consider $x', x'' \in X$ and $\lambda \in (0, 1)$. For $\lambda \in \{0, 1\}$, the desired result is clearly true. Since X is convex, $x^\lambda := (1 - \lambda)x' + \lambda x'' \in X$. By assumption,

$$f(x'') - f(x^\lambda) \leq Df(x^\lambda)(x'' - x^\lambda) \quad \text{and}$$

$$f(x') - f(x^\lambda) \leq Df(x^\lambda)(x' - x^\lambda)$$

Multiplying the first expression by λ , the second one by $(1 - \lambda)$ and summing up, we get

$$\lambda(f(x'') - f(x^\lambda)) + (1 - \lambda)(f(x') - f(x^\lambda)) \leq Df(x^\lambda)(\lambda(x'' - x^\lambda) + (1 - \lambda)(x' - x^\lambda))$$

Since

$$\lambda(x'' - x^\lambda) + (1 - \lambda)(x' - x^\lambda) = x^\lambda - x^\lambda = 0,$$

we get

$$\lambda f(x'') + (1 - \lambda) f(x') \leq f(x^\lambda),$$

i.e., the desired result. \blacksquare

Definition 683 Given a symmetric matrix $A_{n \times n}$, A is negative semidefinite if $\forall x \in \mathbb{R}^n$, $xAx \leq 0$. A is negative definite if $\forall x \in \mathbb{R}^n \setminus \{0\}$, $xAx < 0$.

Proposition 684 Consider a C^2 function f .

f is concave

\Leftrightarrow

$\forall x \in X$, $D^2 f(x)$ is negative semidefinite.

Proof.[\Rightarrow]

We want to show that $\forall u \in \mathbb{R}^n$, $\forall x_0 \in X$, it is the case that $u^T D^2 f(x_0) u \leq 0$. Since X is open, $\forall x_0 \in X \exists a \in \mathbb{R}_{++}$ such that $|h| < a \Rightarrow (x_0 + hu) \in X$. Taken $I := (-a, a) \subseteq \mathbb{R}$, define

$$g : I \rightarrow \mathbb{R}, \quad g : h \mapsto f(x_0 + hu) - f(x_0) - Df(x_0)hu.$$

Observe that

$$g'(h) = D_x f(x_0 + hu) \cdot u + Df(x_0) \cdot u$$

and

$$g''(h) = u \cdot D^2 f(x_0 + hu) \cdot u$$

Since f is a concave function, from Proposition 682, we have that $\forall h \in I$, $g(h) \leq 0$. Since $g(0) = 0$, $h = 0$ is a maximum point. Then, $g'(0) = 0$ and

$$g''(0) \leq 0 \quad (1).$$

Moreover, $\forall h \in I$, $g'(h) = Df(x_0 + hu)u - Df(x_0)u$ and $g''(h) = u^T D^2 f(x_0 + hu)u$. Then,

$$g''(0) = u \cdot D^2 f(x_0) \cdot u \quad (2).$$

(1) and (2) give the desired result.

[\Leftarrow]

Consider $x, x^0 \in X$. From Taylor's Theorem (see Proposition 644), we get

$$f(x) = f(x^0) + Df(x^0)(x - x^0) + \frac{1}{2} (x - x^0)^T D^2 f(\bar{x})(x - x^0)$$

where $\bar{x} = x^0 + \theta(x - x^0)$, for some $\theta \in (0, 1)$. Since, by assumption, $(x - x^0)^T D^2 f(\bar{x})(x - x^0) \leq 0$, we have that

$$f(x) - f(x^0) \leq Df(x^0)(x - x^0),$$

the desired result.

■

Some Properties.

Proposition 685 Consider a concave function f . If x_0 is a local maximum point, then it is a global maximum point.

Proof.

By definition of local maximum point, we know that $\exists \delta > 0$ such that $\forall x \in B(x_0, \delta)$, $f(x_0) \geq f(x)$. Take $y \in X$; we want to show that $f(x_0) \geq f(y)$.

Since X is convex,

$$\forall \lambda \in [0, 1], (1 - \lambda)x^0 + \lambda y \in X.$$

Take $\lambda^0 > 0$ and sufficiently small to have $(1 - \lambda^0)x^0 + \lambda^0 y \in B(x^0, \delta)$. To find such λ^0 , just solve the inequality $\|(1 - \lambda^0)x^0 + \lambda^0 y - x^0\| = \|\lambda^0(y - x^0)\| = |\lambda^0| \|y - x^0\| < \delta$, where, without loss of generality, $y \neq x^0$.

Then,

$$f(x^0) \geq f((1 - \lambda^0)x^0 + \lambda^0 y) \stackrel{f \text{ concave}}{\geq} (1 - \lambda^0)f(x^0) + \lambda^0 f(y),$$

or $\lambda^0 f(x^0) \geq \lambda^0 f(y)$. Dividing both sides of the inequality by $\lambda^0 > 0$, we get $f(x^0) \geq f(y)$.

■

Proposition 686 Consider a differentiable and concave function f . If $Df(x^0) = 0$, then x^0 is a global maximum point.

Proof.

From Proposition 682, if $Df(x^0) = 0$, we get that $\forall x \in X$, $f(x^0) \geq f(x)$, the desired result.

■

17.1.2 Strictly Concave Functions.

Definition 687 Consider a C^0 function f . f is strictly concave iff $\forall x', x'' \in X$ such that $x' \neq x''$, $\forall \lambda \in (0, 1)$,

$$f((1 - \lambda)x' + \lambda x'') > (1 - \lambda)f(x') + \lambda f(x'').$$

Proposition 688 Consider a C^1 function f .

f is strictly concave

$\Leftrightarrow \forall x', x'' \in X$ such that $x' \neq x''$,

$$f(x'') - f(x') < Df(x')(x'' - x').$$

Proof.

[\Rightarrow]

Since strict concavity implies concavity, it is the case that

$$\forall x', x'' \in X, f(x'') - f(x') \leq Df(x')(x'' - x'). \quad (17.1)$$

By contradiction, suppose f is not strictly concave. Then, from 17.1, we have that

$$\exists x', x'' \in X, x' \neq x'' \text{ such that } f(x'') = f(x') + Df(x')(x'' - x'). \quad (17.2)$$

From the definition of strict concavity and 17.2, for $\lambda \in (0, 1)$,

$$f((1 - \lambda)x' + \lambda x'') > (1 - \lambda)f(x') + \lambda f(x'') + \lambda Df(x')(x'' - x')$$

or

$$f((1 - \lambda)x' + \lambda x'') > f(x') + \lambda Df(x')(x'' - x'). \quad (17.3)$$

Applying 17.1 to the points $x(\lambda) := (1 - \lambda)x' + \lambda x''$ and x' , we get that for $\lambda \in (0, 1)$,

$$f((1 - \lambda)x' + \lambda x'') \leq f(x') + Df(x')((1 - \lambda)x' + \lambda x'' - x')$$

or

$$f((1 - \lambda)x' + \lambda x'') \leq f(x') + \lambda Df(x')(x'' - x'). \quad (17.4)$$

And 17.4 contradicts 17.3.

[\Leftarrow] The proof is very similar to that one in Proposition 679.

■

Proposition 689 Consider a C^2 function f . If

$$\forall x \in X, \quad D^2 f(x) \text{ is negative definite,}$$

then f is strictly concave.

Proof.

The proof is similar to that of Proposition 684.

■

Remark 690 In the above Proposition, the opposite implication does not hold. The standard counterexample is $f : \mathbb{R} \rightarrow \mathbb{R}, f : x \mapsto -x^4$.

Some Properties.

Proposition 691 Consider a strictly concave, C^0 function f . If x^0 is a local maximum point, then it is a strict global maximum point, i.e., the unique global maximum point.

Proof.

First, we show that a. it is a global maximum point, and then b. the desired result.

a. It follows from the fact that strict concavity is stronger than concavity and from Proposition 685.

b. Suppose otherwise, i.e., $\exists x', x^0 \in X$ such that $x' \neq x^0$ and both of them are global maximum points. Then, $\forall \lambda \in (0, 1)$, $(1 - \lambda)x' + \lambda x^0 \in X$, since X is convex, and

$$f((1 - \lambda)x' + \lambda x^0) > (1 - \lambda)f(x') + \lambda f(x^0) = f(x') = f(x^0),$$

a contradiction.

■

Proposition 692 Consider a strictly concave, differentiable function f . If $Df(x^0) = 0$, then x^0 is a strict global maximum point.

Proof.

Take an arbitrary $x \in X$ such that $x \neq x^0$. Then from Proposition 688, we have that $f(x) < f(x^0) + Df(x^0)(x - x^0) = f(x^0)$, the desired result.

■

17.1.3 Quasi-Concave Functions.

Definitions.

Definition 693 Consider a C^0 function f . f is quasi-concave iff $\forall x', x'' \in X, \forall \lambda \in [0, 1]$,

$$f((1 - \lambda)x' + \lambda x'') \geq \min \{f(x'), f(x'')\}.$$

Proposition 694 If $f : X \rightarrow \mathbb{R}$ is a quasi-concave function and $F : \mathbb{R} \rightarrow \mathbb{R}$ is non decreasing, then $F \circ f$ is a quasi-concave function.

Proof.

Without loss of generality, assume

$$f(x'') \geq f(x') \quad (1).$$

Then, since f is quasi-concave, we have

$$f((1 - \lambda)x' + \lambda x'') \geq f(x') \quad (2).$$

Then,

$$F(f((1 - \lambda)x' + \lambda x'')) \stackrel{(a)}{\geq} F(f(x')) \stackrel{(b)}{=} \min \{F(f(x')), F(f(x''))\},$$

where (a) comes from (2) and the fact that F is nondecreasing, and (b) comes from (1) and the fact that F is nondecreasing.

■

Proposition 695 Consider a C^0 function f . f is quasi-concave $\Leftrightarrow \forall \alpha \in \mathbb{R}, B(\alpha) := \{x \in X : f(x) \geq \alpha\}$ is convex.

Proof.

[\Rightarrow] [Strategy: write what you want to show].

We want to show that $\forall \alpha \in \mathbb{R}$ and $\forall \lambda \in [0, 1]$, we have that

$$\langle x', x'' \in B(\alpha) \rangle \Rightarrow \langle (1 - \lambda)x' + \lambda x'' \in B(\alpha) \rangle,$$

i.e.,

$$\langle f(x') \geq \alpha \text{ and } f(x'') \geq \alpha \rangle \Rightarrow \langle f((1 - \lambda)x' + \lambda x'') \geq \alpha \rangle.$$

But by Assumption,

$$f((1 - \lambda)x' + \lambda x'') \geq \min \{f(x'), f(x'')\} \stackrel{\text{def } x', x''}{\geq} \alpha.$$

[\Leftarrow]

Consider arbitrary $x', x'' \in X$. Define $\alpha := \min \{f(x'), f(x'')\}$. Then $x', x'' \in B(\alpha)$. By assumption, $\forall \lambda \in [0, 1]$, $(1 - \lambda)x' + \lambda x'' \in B(\alpha)$, i.e.,

$$f((1 - \lambda)x' + \lambda x'') \geq \alpha := \min \{f(x'), f(x'')\}.$$

■

Proposition 696 Consider a differentiable function f . f is quasi-concave $\Leftrightarrow \forall x', x'' \in X$,

$$f(x'') - f(x') \geq 0 \Rightarrow Df(x')(x'' - x') \geq 0.$$

Proof.

[\Rightarrow] [Strategy: Use the definition of directional derivative.]

Take x', x'' such that $f(x'') \geq f(x')$. By assumption,

$$f((1 - \lambda)x' + \lambda x'') \geq \min \{f(x'), f(x'')\} = f(x')$$

and

$$f((1-\lambda)x' + \lambda x'') - f(x') \geq 0.$$

Dividing both sides of the above inequality by $\lambda > 0$, and taking limits for $\lambda \rightarrow 0^+$, we get

$$\lim_{\lambda \rightarrow 0^+} \frac{f(x' + \lambda(x'' - x')) - f(x')}{\lambda} = Df(x')(x'' - x') \geq 0.$$

[\Leftarrow]

Without loss of generality, take

$$f(x') = \min\{f(x'), f(x'')\} \quad (1).$$

Define

$$\varphi : [0, 1] \rightarrow \mathbb{R}, \quad \varphi : \lambda \mapsto f((1-\lambda)x' + \lambda x'').$$

We want to show that

$$\forall \lambda \in [0, 1], \quad \varphi(\lambda) \geq \varphi(0).$$

Suppose otherwise, i.e., $\exists \lambda^* \in [0, 1]$ such that $\varphi(\lambda^*) < \varphi(0)$. Observe that in fact it cannot be $\lambda^* \in \{0, 1\}$: if $\lambda^* = 0$, we would have $\varphi(0) < \varphi(0)$, and if $\lambda^* = 1$, we would have $\varphi(1) < \varphi(0)$, i.e., $f(x'') < f(x')$, contradicting (1). Then, we have that

$$\exists \lambda^* \in (0, 1) \text{ such that } \varphi(\lambda^*) < \varphi(0) \quad (2).$$

Observe that from (1), we also have that

$$\varphi(1) \geq \varphi(0) \quad (3).$$

Therefore, see Lemma 697, $\exists \lambda^{**} > \lambda^*$ such that

$$\varphi'(\lambda^{**}) > 0 \quad (4), \text{ and}$$

$$\varphi(\lambda^{**}) < \varphi(0) \quad (5).$$

From (4), and using the definition of φ' , and the Chain Rule,² we get

$$0 < \varphi'(\lambda^{**}) = [Df((1-\lambda^{**})x' + \lambda^{**}x'')](x'' - x') \quad (6).$$

Define $x^{**} := (1-\lambda^{**})x' + \lambda^{**}x''$. From (5), and the assumption, we get that

$$f(x^{**}) < f(x').$$

Therefore, by assumption,

$$0 \leq Df(x^{**})(x' - x^{**}) = Df(x^{**})(-\lambda^{**})(x'' - x'),$$

i.e.,

$$[Df(x^{**})]^T(x'' - x') \leq 0 \quad (7).$$

But (7) contradicts (6).

■

Lemma 697 Consider a function $g : [a, b] \rightarrow \mathbb{R}$ with the following properties:

1. g is differentiable on (a, b) ;
 2. there exists $c \in (a, b)$ such that $g(b) \geq g(a) > g(c)$.
- Then, $\exists t \in (c, b)$ such that $g'(t) > 0$ and $g(t) < g(a)$.

²Defined $v : [0, 1] \rightarrow X \subseteq \mathbb{R}^n, \lambda \mapsto (1-\lambda)x' + \lambda x''$, we have that $\varphi = f \circ v$. Therefore, $\varphi'(\lambda^*) = Df(v(\lambda^*)) \cdot Dv(\lambda^*)$.

Proof.

Without loss of generality and to simplify notation, assume $g(a) = 0$. Define $A := \{x \in [c, b] : g(x) = 0\}$.

Observe that $A = [c, b] \cap g^{-1}(0)$ is closed; and it is non empty, because g is continuous and by assumption $g(c) < 0$ and $g(b) \geq 0$.

Therefore, A is compact, and we can define $\xi := \min A$.

Claim. $x \in [c, \xi) \Rightarrow g(x) < 0$.

Suppose not, i.e., $\exists y \in (c, \xi)$ such that $g(y) \geq 0$. If $g(y) = 0$, ξ could not be $\min A$. If $g(y) > 0$, since $g(c) < 0$ and g is continuous, there exists $x' \in (c, y) \subseteq (c, \xi)$, again contradicting the definition of ξ . End of the proof of the Claim.

Finally, applying Lagrange Theorem to g on $[c, \xi]$, we have that $\exists t \in (c, \xi)$ such that $g'(t) = \frac{g(\xi) - g(c)}{\xi - c}$. Since $g(\xi) = 0$ and $g(c) < 0$, we have that $g'(t) < 0$. From the above Claim, the desired result then follows.

■

Proposition 698 Consider a C^2 function f . If f is quasi-concave then

$$\forall x \in X, \forall \Delta \in \mathbb{R}^n \text{ such that } Df(x) \cdot \Delta = 0, \quad \Delta^T D^2 f(x) \Delta \leq 0.$$

Proof.

for another proof- see Laura's file

Suppose otherwise, i.e., $\exists x \in X$, and $\exists \Delta \in \mathbb{R}^n$ such that $Df(x) \cdot \Delta = 0$ and $\Delta^T D^2 f(x) \Delta > 0$.

Since the function $h : X \rightarrow \mathbb{R}$, $h : x \mapsto \Delta^T D^2 f(x) \Delta$ is continuous and X is open, $\forall \lambda \in [0, 1]$, $\exists \varepsilon > 0$ such that if $\|x - x^0\| < \varepsilon$, then

$$\Delta \cdot D^2 f(\lambda x + (1 - \lambda)x^0) \cdot \Delta > 0 \quad (1).$$

Define $\bar{x} := x^0 + \mu \frac{\Delta}{\|\Delta\|}$, with $0 < \mu < \varepsilon$. Then,

$$\|\bar{x} - x^0\| = \left\| \mu \frac{\Delta}{\|\Delta\|} \right\| = \mu < \varepsilon$$

and \bar{x} satisfies (1). Observe that

$$\Delta = \frac{\|\Delta\|}{\mu} (\bar{x} - x^0).$$

Then, we can rewrite (1) as

$$(\bar{x} - x^0)^T D^2 f(\lambda \bar{x} + (1 - \lambda)x^0) (\bar{x} - x^0) > 0$$

From Taylor Theorem, $\exists \lambda \in (0, 1)$ such that

$$f(\bar{x}) = f(x^0) + (\bar{x} - x^0)^T Df(x^0) + \frac{1}{2} (\bar{x} - x^0)^T D^2 f(\lambda \bar{x} + (1 - \lambda)x^0) (\bar{x} - x^0).$$

Since $Df(x^0) \Delta = 0$ and from (1), we have

$$f(\bar{x}) > f(x^0) \quad (2).$$

Letting $\tilde{x} = x^0 + \mu(-\Delta/\|\Delta\|)$, using the same procedure as above, we can conclude that

$$f(\tilde{x}) > f(x^0) \quad (3).$$

But, since $x^0 = \frac{1}{2}(\bar{x} + \tilde{x})$, (2) and (3) contradict the Definition of quasi-concavity.

■

Remark 699 In the above Proposition, the opposite implication does not hold. Consider $f : \mathbb{R} \rightarrow \mathbb{R}$, $f : x \mapsto x^4$. From Proposition 695, that function is clearly not quasi-concave. Take $\alpha > 0$. Then $B(\alpha) = \{x \in \mathbb{R} : x^4 \geq \alpha\} = (-\infty, -\sqrt[4]{\alpha}) \cup (\sqrt[4]{\alpha}, +\infty)$ which is not convex.


On the other hand observe the following. $f'(x) = 4x^3$ and $4x^3 \Delta = 0$ if either $x = 0$ or $\Delta = 0$. In both cases $\Delta^T D^2 f(x) \Delta = 0$. (This example is taken from Avriel M. and others (1988), page 91).

Some Properties.

Remark 700 Consider a quasi concave function f . It is NOT the case that

if x_0 is a local maximum point, then it is a global maximum point. To see that, consider the following function.

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f : x \mapsto \begin{cases} -x^2 + 1 & \text{if } x < 1 \\ 0 & \text{if } x \geq 1 \end{cases}$$



Proposition 701 Consider a C^0 quasi-concave function f . If x_0 is a strict local maximum point, then it is a strict global maximum point.

Proof.

By assumption, $\exists \delta > 0$ such that if $x \in B(x_0, \delta) \cap X$ and $x_0 \neq x$, then $f(x_0) > f(x)$.

Suppose the conclusion of the Proposition is false; then $\exists x' \in X$ such that $f(x') \geq f(x_0)$.

Since f is quasi-concave,

$$\forall \lambda \in [0, 1], \quad f((1 - \lambda)x_0 + \lambda x') \geq f(x_0). \quad (1)$$

For sufficiently small λ , $(1 - \lambda)x_0 + \lambda x' \in B(x_0, \delta)$ and (1) above holds, contradicting the fact that x_0 is the *strict* local maximum point.

■

Proposition 702 Consider $f : (a, b) \rightarrow \mathbb{R}$. f monotone $\Rightarrow f$ quasi-concave.

Proof.

Without loss of generality, take $x'' \geq x'$.

Case 1. f is increasing. Then $f(x'') \geq f(x')$. If $\lambda \in [0, 1]$, then $(1 - \lambda)x' + \lambda x'' = x' + \lambda(x'' - x') \geq x'$ and therefore $f((1 - \lambda)x' + \lambda x'') \geq f(x')$.


Case 2. f is decreasing. Then $f(x'') \leq f(x')$. If $\lambda \in [0, 1]$, then $(1 - \lambda)x' + \lambda x'' = (1 - \lambda)x' - (1 - \lambda)x'' + x'' = x'' - (1 - \lambda)(x'' - x') \leq x''$ and therefore $f((1 - \lambda)x' + \lambda x'') \geq f(x'')$.

■

Remark 703 The following statement is false: If f_1 and f_2 are quasi-concave and $a, b \in \mathbb{R}_+$, then $af_1 + bf_2$ is quasi-concave.

It is enough to consider $f_1, f_2 : \mathbb{R} \rightarrow \mathbb{R}$, $f_1(x) = x^3 + x$, and $f_2(x) = -4x$. Since $f'_1 > 0$, then f_1 and, of course, f_2 are monotone and then, from Proposition 702, they are quasi-concave. On the other hand, $g(x) = f_1(x) + f_2(x) = x^3 - x$ has a strict local maximum in $x = -1$ which is not a strict global maximum, and therefore, from Proposition 701, g is not quasi-concave.

$$x^3 - 3x$$



Remark 704 Consider a differentiable quasi-concave function f . It is NOT the case that if $Df(x^0) = 0$, then x^0 is a global maximum point.

Just consider $f : \mathbb{R} \rightarrow \mathbb{R}, f : x \mapsto x^3$ and $x_0 = 0$, and use Proposition 702.

17.1.4 Strictly Quasi-concave Functions.

Definitions.

Definition 705 Consider a C^0 function f . f is strictly quasi-concave iff $\forall x', x'' \in X$, such that $x' \neq x''$, and $\forall \lambda \in (0, 1)$, we have that

$$f((1 - \lambda)x' + \lambda x'') > \min \{f(x'), f(x'')\}.$$

Proposition 706 Consider a C^0 function f . f is strictly quasi-concave $\Rightarrow \forall \alpha \in \mathbb{R}, B(\alpha) := \{x \in X : f(x) \geq \alpha\}$ is strictly convex.

Proof.

Taken an arbitrary α and $x', x'' \in B(\alpha)$, with $x' \neq x''$, we want to show that $\forall \lambda \in (0, 1)$, we have that

$$x^\lambda := (1 - \lambda)x' + \lambda x'' \in \text{Int } B(\alpha)$$

Since f is strictly quasi-concave,

$$f(x^\lambda) > \min \{f(x'), f(x'')\} \geq \alpha$$

Since f is C^0 , there exists $\delta > 0$ such that $\forall x \in B(x^\lambda, \delta)$

$$f(x) > \alpha$$

i.e., $B(x^\lambda, \delta) \subseteq B(\alpha)$, as desired. Of course, we are using the fact that $\{x \in X : f(x) > \alpha\} \subseteq B(\alpha)$.

■

Remark 707 Observe that in Proposition 706, the opposite implication does not hold true: just consider $f : \mathbb{R} \rightarrow \mathbb{R}, f : x \mapsto 1$.

Observe that $\forall \alpha \leq 1, B(\alpha) = \mathbb{R}$, and $\forall \alpha > 1, B(\alpha) = \emptyset$. On the other hand, f is not strictly quasi-concave.

Definition 708 Consider a differentiable function f . f is differentiable-strictly-quasi-concave iff $\forall x', x'' \in X$, such that $x' \neq x''$, we have that

$$f(x'') - f(x') \geq 0 \Rightarrow Df(x')(x'' - x') > 0.$$

Proposition 709 Consider a differentiable function f .

If f is differentiable-strictly-quasi-concave, then f is strictly quasi-concave.

Proof.

The proof is analogous to the case of quasi concave functions.

■

Remark 710 Given a differentiable function, it is not the case that strict-quasi-concavity implies differentiable-strict-quasi-concavity.

$f : \mathbb{R} \rightarrow \mathbb{R}$, $f : x \mapsto x^3$ a. is differentiable and strictly quasi concave and b. it is not differentiable-strictly-quasi-concave.

a. f is strictly increasing and therefore strictly quasi concave - see Fact below.

b. Take $x' = 0$ and $x'' = 1$. Then $f(1) = 1 > 0 = f(0)$. But $Df(x')(x'' - x') = 0 \cdot 1 = 0 \not> 0$.

Remark 711 If we restrict the class of differentiable functions to those with non-zero gradients everywhere in the domain, then differentiable-strict-quasi-concavity and strict-quasi-concavity are equivalent (see Balasko (1988), Math. 7.2.).

Fact. Consider $f : (a, b) \rightarrow \mathbb{R}$. f strictly monotone $\Rightarrow f$ strictly quasi concave.

Proof.

By assumption, $x' \neq x''$, say $x' < x''$ implies that $f(x') < f(x'')$ (or $f(x') > f(x'')$). If $\lambda \in (0, 1)$, then $(1 - \lambda)x' + \lambda x'' > x'$ and therefore $f((1 - \lambda)x' + \lambda x'') > \min\{f(x'), f(x'')\}$.

■

Proposition 712 Consider a C^2 function f . If

$$\forall x \in X, \forall \Delta \in \mathbb{R}^n \setminus \{0\}, \text{ we have that } \langle Df(x) \Delta = 0 \Rightarrow \Delta^T D^2 f(x) \Delta < 0 \rangle,$$

then f is differentiable-strictly-quasi-concave.

Proof.

Suppose otherwise, i.e., there exist $x', x'' \in X$ such that

$$x' \neq x'', f(x'') \geq f(x') \quad \text{and} \quad Df(x')(x'' - x') \leq 0.$$

Since X is an open set, $\exists a \in \mathbb{R}_{++}$ such the following function is well defined:

$$g : [-a, 1] \rightarrow \mathbb{R}, g : h \mapsto f((1 - h)x' + hx'').$$

Since g is continuous, there exists $h_m \in [0, 1]$ which is a global minimum. We now proceed as follows. Step 1. $h_m \notin \{0, 1\}$. Step 2. h_m is a strict local maximum point, a contradiction.

Preliminary observe that

$$g'(h) = Df(x' + h(x'' - x')) \cdot (x'' - x')$$

and

$$g''(h) = (x'' - x')^T \cdot D^2 f(x' + h(x'' - x')) \cdot (x'' - x').$$

Step 1. If $Df(x')(x'' - x') = 0$, then, by assumption,

$$(x'' - x')^T \cdot D^2 f(x' + h(x'' - x')) \cdot (x'' - x') < 0.$$

Therefore, $zero$ is a strict local maximum (see, for example, Theorem 13.10, page 378, in Apostol (1974)). Therefore, there exists $h^* \in \mathbb{R}$ such that $g(h^*) = f(x' + h^*(x'' - x')) < f(x') = g(0)$.

If

$$g'(0) = Df(x')(x'' - x') < 0,$$

then there exists $h^{**} \in \mathbb{R}$ such that

$$g(h^{**}) = f(x' + h^{**}(x'' - x')) < f(x') = g(0).$$

Moreover, $g(1) = f(x'') \geq f(x')$. In conclusion, neither *zero* nor *one* can be global minimum points for g on $[0, 1]$.

Step 2. Since the global minimum point $h_m \in (0, 1)$, we have that

$$0 = g'(h_m) = Df(x' + h_m(x'' - x'))(x'' - x').$$

Then, by assumption,

$$g''(0) = (x'' - x')^T \cdot D^2 f(x' + h_m(x'' - x')) \cdot (x'' - x') < 0,$$

but then h_m is a strict local maximum point, a contradiction.

■

Remark 713 *Differentiable-strict-quasi-concavity does not imply the condition presented in Proposition 712. $f : \mathbb{R} \rightarrow \mathbb{R}$, $f : x \mapsto -x^4$ is differentiable-strictly-quasi-concave (in next section we will show that strict-concavity implies differentiable-strict-quasi-concavity). On the other hand, take $x^* = 0$. Then $Df(x^*) = 0$. Therefore, for any $\Delta \in \mathbb{R}^n \setminus \{0\}$, we have $Df(x^*)\Delta = 0$, but $\Delta^T D^2 f(x^*)\Delta = 0 \not< 0$.*

Some Properties.

Proposition 714 *Consider a differentiable-strictly-quasi-concave function f .*

x^ is a strict global maximum point $\Leftrightarrow Df(x^*) = 0$.*

Proof.

[\Rightarrow] Obvious.

[\Leftarrow] From the contrapositive of the definition of differentiable-strictly-quasi-concave function, we have:

$\forall x^*, x'' \in X$, such that $x^* \neq x''$, it is the case that $Df(x^*)(x'' - x^*) \leq 0 \Rightarrow f(x'') - f(x^*) < 0$ or $f(x^*) > f(x'')$. Since $Df(x^*) = 0$, then the desired result follows.

■

Remark 715 *Obviously, we also have that if f is differentiable-strictly-quasi-concave, it is the case that:*

x^ local maximum point $\Rightarrow x^*$ is a strict maximum point.*

Remark 716 *The above implication is true also for continuous strictly quasi concave functions. (Suppose otherwise, i.e., $\exists x' \in X$ such that $f(x') \geq f(x^*)$. Since f is strictly quasi-concave, $\forall \lambda \in (0, 1)$, $f((1 - \lambda)x^* + \lambda x') > f(x^*)$, which for sufficiently small λ contradicts the fact that x^* is a local maximum point.*

Is there a definition of ?-concavity weaker than concavity and such that:

If f is a ?-concave function, then

x^* is a global maximum point iff $Df(x^*) = 0$.

The answer is given in the next section.

17.1.5 Pseudo-concave Functions.

Definition 717 *Consider a differentiable function f . f is pseudo-concave iff*

$$\forall x', x'' \in X, f(x'') > f(x') \Rightarrow Df(x')(x'' - x') > 0,$$

or

$$\forall x', x'' \in X, Df(x')(x'' - x') \leq 0 \Rightarrow f(x'') \leq f(x').$$

Proposition 718 *If f is a pseudo-concave function, then*

$$x^* \text{ is a global maximum point} \Leftrightarrow Df(x^*) = 0.$$

Proof.

[\Rightarrow] Obvious.

[\Leftarrow] $Df(x^*) = 0 \Rightarrow \forall x \in X, Df(x^*)(x - x^*) \leq 0 \Rightarrow f(x) \leq f(x^*)$.

■

Remark 719 *Observe that the following “definition of pseudo-concavity” will not be useful:*

$$\forall x', x'' \in X, Df(x')(x'' - x') \geq 0 \Rightarrow f(x'') \leq f(x') \quad (17.5)$$

For such a definition the above Proposition would still apply, but it is not weaker than concavity. Simply consider the function $f : \mathbb{R} \rightarrow \mathbb{R}, f : x \mapsto -x^2$. That function is concave, but it does not satisfy condition (17.5). Take $x' = -2$ and $x'' = -1$. Then, $f'(x')(x'' - x') = 4(-1 - (-2)) = 4 > 0$, but $f(x'') = -1 > f(x') = -4$.


We summarize some of the results of this subsection in the following tables.

Class of function	Fundamental properties		
	$C \Rightarrow G \text{ max}$	$L \text{ max} \Rightarrow G \text{ max}$	Uniqueness of G. max
Strictly concave	Yes	Yes	Yes
Concave	Yes	Yes	No
Diff.ble-str.-q.-conc.	Yes	Yes	Yes
Pseudoconcave	Yes	Yes	No
Quasiconcave	No	No	No

where C stands for property of being a critical point, and L and G stand for local and global, respectively. Observe that the first, the second and the last row of the second column apply to the case of C^0 and not necessarily differentiable functions.

17.2 Relationships among Different Kinds of Concavity

The relationships among different definitions of concavity in the case of *differentiable functions* are summarized in the following table.



All the implications which are not implied by those explicitly written do not hold true.

In what follows, we prove the truth of each implication described in the table and we explain why the other implications do not hold.

Recall that

1. $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a linear function iff $\forall x', x'' \in \mathbb{R}^n, \forall a, b \in \mathbb{R} f(ax' + bx'') = af(x') + bf(x'')$;
2. $g : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is an affine function iff there exists a linear function $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $c \in \mathbb{R}^m$ such that $\forall x \in \mathbb{R}^n, g(x) = f(x) + c$.

$SC \Rightarrow C$

Obvious (“ $a > b \Rightarrow a \geq b$ ”).

$$\boxed{C \Rightarrow PC}$$

From the assumption and from Proposition 682, we have that $f(x'') - f(x') \leq Df(x')(x'' - x')$. Then $f(x'') - f(x') > 0 \Rightarrow Df(x')(x'' - x') > 0$.

$$\boxed{PC \Rightarrow QC}$$

Suppose otherwise, i.e., $\exists x', x'' \in X$ and $\exists \lambda^* [0, 1]$ such that

$$f((1 - \lambda^*)x' + \lambda^*x'') < \min\{f(x'), f(x'')\}.$$

Define $x(\lambda) := (1 - \lambda)x' + \lambda x''$. Consider the segment $L(x', x'')$ joining x' to x'' . Take $\bar{\lambda} \in \arg \min_{\lambda} f(x(\lambda))$ s.t. $\lambda \in [0, 1]$. $\bar{\lambda}$ is well defined from the Extreme Value Theorem. Observe that $\bar{\lambda} \neq 0, 1$, because $f(x(\lambda^*)) < \min\{f(x(0)) = f(x'), f(x(1)) = f(x'')\}$.

Therefore, $\forall \lambda \in [0, 1]$ and $\forall \mu \in (0, 1)$,

$$f(x(\bar{\lambda})) \leq f((1 - \mu)x(\bar{\lambda}) + \mu x(\lambda)).$$

$$\begin{array}{ccccc} & & (1 - \mu)x(\bar{\lambda}) + \mu x(\lambda) & & \\ & & \downarrow & & \\ \begin{array}{c} \cdot \\ \uparrow \\ x' \end{array} & \begin{array}{c} \cdot \\ \uparrow \\ x(\lambda) \end{array} & & \begin{array}{c} \cdot \\ \uparrow \\ x(\bar{\lambda}) \end{array} & \begin{array}{c} \cdot \\ \uparrow \\ x'' \end{array} \end{array}$$

Then,

$$\forall \lambda \in [0, 1], \quad 0 \leq \lim_{\mu \rightarrow 0^+} \frac{f((1 - \mu)x(\bar{\lambda}) + \mu x(\lambda)) - f(x(\bar{\lambda}))}{\mu} = Df(x(\bar{\lambda}))(x(\lambda) - x(\bar{\lambda})).$$

Taking $\lambda = 0, 1$ in the above expression, we get:

$$Df(x(\bar{\lambda}))(x' - x(\bar{\lambda})) \geq 0 \quad (1)$$

and

$$Df(x(\bar{\lambda}))(x'' - x(\bar{\lambda})) \geq 0 \quad (2).$$

Since

$$x' - x(\bar{\lambda}) = x' - (1 - \bar{\lambda})x' - \bar{\lambda}x'' = -\bar{\lambda}(x'' - x') \quad (3),$$

and

$$x'' - x(\bar{\lambda}) = x'' - (1 - \bar{\lambda})x' - \bar{\lambda}x'' = (1 - \bar{\lambda})(x'' - x') \quad (4),$$

substituting (3) in (1), and (4) in (2), we get

$$\stackrel{(-)}{-\bar{\lambda}} \cdot [Df(x(\bar{\lambda})) \cdot (x'' - x')] \geq 0,$$

and

$$\stackrel{(+)}{(1 - \bar{\lambda})} \cdot [Df(x(\bar{\lambda})) \cdot (x'' - x')] \geq 0.$$

Therefore,

$$\begin{aligned} 0 &= Df(x(\bar{\lambda})) \cdot (x'' - x') = Df(x(\bar{\lambda})) \cdot (1 - \bar{\lambda}) \cdot (x'' - x') \stackrel{(4)}{=} \\ &= Df(x(\bar{\lambda})) \cdot (x'' - x(\bar{\lambda})). \end{aligned}$$

Then, by pseudo-concavity,

$$f(x'') \leq f(x(\bar{\lambda})) \quad (5).$$

By assumption,

$$f(x(\lambda^*)) < f(x'') \quad (6).$$

(5) and (6) contradict the definition of $\bar{\lambda}$.

$$\boxed{DSQC \Rightarrow PC}$$

Obvious.

$$\boxed{SC \Rightarrow DSQC}$$

Obvious.

$$\boxed{L \Rightarrow C}$$


Obvious.

$$\boxed{C \not\Rightarrow SC}$$

$f : \mathbb{R} \rightarrow \mathbb{R}, f : x \mapsto x$.

$$\boxed{QC \not\Rightarrow PC}$$

$$f : \mathbb{R} \rightarrow \mathbb{R}, f : x \mapsto \begin{cases} 0 & \text{if } x \leq 0 \\ e^{-\frac{1}{x^2}} & \text{if } x > 0 \end{cases}$$



f is clearly nondecreasing and therefore, from Lemma 702, quasi-concave.

f is not pseudo-concave: $0 < f(-1) > f(1) = 0$, but

$$f'(1)(-1 - 1) = 0 \cdot (-2) = 0.$$

$$\boxed{PC \not\Rightarrow C}, \boxed{DSQC \not\Rightarrow C} \text{ and } \boxed{DSQC \not\Rightarrow SC}$$

Take $f : (1, +\infty) \rightarrow \mathbb{R}, f : x \mapsto x^3$.

Take $x' < x''$. Then $f(x'') > f(x')$. Moreover, $Df(x')(x'' - x') \stackrel{(>0)}{>} 0$. Therefore, f is $DSQC$ and therefore PC . Since $f''(x) > 0$, f is strictly convex and therefore it is not concave and, a fortiori, it is not strictly concave.

$$\boxed{PC \not\Rightarrow DSQC}, \boxed{C \not\Rightarrow DSQC}$$

Consider $f : \mathbb{R} \rightarrow \mathbb{R}, f : x \mapsto 1$. f is clearly concave and PC , as well ($\forall x', x'' \in \mathbb{R}, Df(x')(x'' - x') \geq 0$). Moreover, any point in \mathbb{R} is a critical point, but it is not the unique global maximum point. Therefore, from Proposition 714, f is not differentiable - strictly - quasi - concave.

$$\boxed{QC \not\Rightarrow DSQC}$$

If so, we would have $QC \Rightarrow DSQC \Rightarrow PC$, contradicting the fact that $QC \not\Rightarrow PC$.

$$\boxed{C \not\Rightarrow L} \text{ and } \boxed{SC \not\Rightarrow L}$$

$f : \mathbb{R} \rightarrow \mathbb{R}, f : x \mapsto -x^2$.

17.2.1 Hessians and Concavity.

In this subsection, we study the relation between submatrices of a matrix involving the Hessian matrix of a C^2 function and the concavity of that function.

Definition 720 Consider a matrix $A_{n \times n}$. Let $1 \leq k \leq n$.

A k -th order principal submatrix (minor) of A is the (determinant of the) square submatrix of A obtained deleting $(n - k)$ rows and $(n - k)$ columns in the same position. Denote these matrices by \tilde{D}_k^i .

The k -th order leading principal submatrix (minor) of A is the (determinant of the) square submatrix of A obtained deleting the last $(n - k)$ rows and the last $(n - k)$ columns. Denote these matrices by D_k .

Example 721 Consider

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}.$$

Then

$$\tilde{D}_1^1 = a_{11}, \quad \tilde{D}_1^2 = a_{22}, \quad \tilde{D}_1^3 = a_{33}, \quad D_1 = \tilde{D}_1^1 = a_{11};$$

$$\tilde{D}_2^1 = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}, \quad \tilde{D}_2^2 = \begin{bmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{bmatrix}, \quad \tilde{D}_2^3 = \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix},$$

$$D_2 = \tilde{D}_2^1 = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix};$$

$$D_3 = \tilde{D}_3^1 = A.$$

Definition 722 Consider a C^2 function $f : X \subseteq \mathbb{R}^n \rightarrow \mathbb{R}$. The bordered Hessian of f is the following matrix

$$B_f(x) = \begin{bmatrix} 0 & Df(x) \\ [Df(x)]^T & D^2f(x) \end{bmatrix}_{(n+1) \times (n+1)}.$$

Theorem 723 (Simon, (1985), Theorem 1.9.c, page 79 and Sydsaeter (1981), Theorem 5.17, page 259). Consider a C^2 function $f : X \rightarrow \mathbb{R}$.

1. If $\forall x \in X, \forall k \in \{1, \dots, n\}$,
 $\text{sign}(k - \text{leading principal minor of } D^2f(x)) = \text{sign}(-1)^k$,
then f is strictly concave.
2. $\forall x \in X, \forall k \in \{1, \dots, n\}$,
 $\text{sign}(\text{non zero } k - \text{principal minor of } D^2f(x)) = \text{sign}(-1)^k$,
iff f is concave.
3. If $n \geq 2$ and $\forall x \in X, \forall k \in \{3, \dots, n+1\}$,
 $\text{sign}(k - \text{leading principal minor of } Bf(x)) = \text{sign}(-1)^{k-1}$,
then f is pseudo concave and, therefore, quasi-concave.
4. If f is quasi-concave, then $\forall x \in X, \forall k \in \{2, \dots, n+1\}$,
 $\text{sign}(\text{non zero } k - \text{leading principal minors of } Bf(x)) = \text{sign}(-1)^{k-1}$

Remark 724 It can be proved that Conditions in part 1 and 2 of the above Theorem are sufficient for $D^2f(x)$ being negative definite and equivalent to $D^2f(x)$ being negative semidefinite, respectively.

Remark 725 (From Sydsaetter (1981), page 239) It is tempting to conjecture that a function f is concave iff

$$\forall x \in X, \forall k \in \{1, \dots, n\}, \quad \text{sign}(\text{non zero } k - \text{leading principal minor of } D^2f(x)) = \text{sign}(-1)^k, \quad (17.6)$$

That conjecture is false. Consider

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad f : (x_1, x_2, x_3) \mapsto -x_2^2 + x_3^2.$$

Then $Df(x) = (0, -2x_2, 2x_3)$ and

$$D^2f(x) = \begin{bmatrix} 0 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

All the leading principal minors of the above matrix are zero, and therefore Condition 17.6 is satisfied, but f is not a concave function. Take $x' = (0, 0, 0)$ and $x'' = (0, 0, 1)$. Then

$$\forall \lambda \in (0, 1), \quad f((1-\lambda)x' + \lambda x'') = \lambda^2 < (1-\lambda)f(x') + \lambda f(x'') = \lambda$$

Example 726 Consider $f : \mathbb{R}_{++}^2 \rightarrow \mathbb{R}, f : (x, y) \mapsto x^\alpha y^\beta$, with $\alpha, \beta \in \mathbb{R}_{++}$. Observe that $\forall (x, y) \in \mathbb{R}_{++}^2, f(x, y) > 0$. Verify that

1. if $\alpha + \beta < 1$, then f is strictly concave;
2. $\forall \alpha, \beta \in \mathbb{R}_{++}$, f is quasi-concave;
3. $\alpha + \beta \leq 1$ if and only if f is concave.

1.

$$D_x f(x, y) = \alpha x^{\alpha-1} y^\beta = \frac{\alpha}{x} f(x, y);$$

$$D_y f(x, y) = \beta x^\alpha y^{\beta-1} = \frac{\beta}{y} f(x, y);$$

$$D_{x,x}^2 f(x, y) = \alpha(\alpha-1) x^{\alpha-2} y^\beta = \frac{\alpha(\alpha-1)}{x^2} f(x, y);$$

$$D_{y,y}^2 f(x, y) = \beta(\beta-1) x^\alpha y^{\beta-2} = \frac{\beta(\beta-1)}{y^2} f(x, y);$$

$$D_{x,y}^2 f(x, y) = \alpha\beta x^{\alpha-1} y^{\beta-1} = \frac{\alpha}{x} \frac{\beta}{y} f(x, y).$$

$$D^2 f(x, y) = f(x, y) \begin{bmatrix} \frac{\alpha(\alpha-1)}{x^2} & \frac{\alpha}{x} \frac{\beta}{y} \\ \frac{\alpha}{x} \frac{\beta}{y} & \frac{\beta(\beta-1)}{y^2} \end{bmatrix}.$$

$$a. \quad \frac{\alpha(\alpha-1)}{x^2} < 0 \Leftrightarrow \alpha \in (0, 1).$$

$$b. \quad \frac{\alpha(\alpha-1)\beta(\beta-1) - \alpha^2\beta^2}{x^2y^2} = \frac{1}{x^2y^2} (\alpha\beta(\alpha\beta - \alpha - \beta + 1) - \alpha^2\beta^2) =$$

$$= \frac{1}{x^2y^2} \alpha\beta(1 - \alpha - \beta) > 0 \stackrel{\alpha, \beta > 0}{\Leftrightarrow} \alpha + \beta < 1.$$

In conclusion, if $\alpha, \beta \in (0, 1)$ and $\alpha + \beta < 1$, then f is strictly concave.

2.

Observe that

$$f(x, y) = g(h(x, y))$$

where

$$h : \mathbb{R}_{++}^2 \rightarrow \mathbb{R}, \quad (x, y) \mapsto \alpha \ln x + \beta \ln y$$

$$g : \mathbb{R} \rightarrow \mathbb{R}, \quad z \mapsto e^z$$

Since h is strictly concave (why?) and therefore quasi-concave and g is strictly increasing, the desired result follows from Proposition 694.

3.

Obvious from above results.

Chapter 18

Maximization Problems

Let the following objects be given:

1. an open convex set $X \subseteq \mathbb{R}^n$, $n \in \mathbb{N}$;
2. $f : X \rightarrow \mathbb{R}$, $g : X \rightarrow \mathbb{R}^m$, $h : X \rightarrow \mathbb{R}^k$, $m, k \in \mathbb{N}$, with f, g, h at least differentiable.

The goal of this Chapter is to study the problem.

$$\begin{array}{ll} \max_{x \in X} & f(x) \\ \text{s.t.} & g(x) \geq 0 \quad (1) \\ & h(x) = 0 \quad (2) \end{array} \quad (18.1)$$

f is called **objective function**; x **choice variable** vector; (1) and (2) in (18.1) **constraints**; g and h **constraint functions**;

$$C := \{x \in X : g(x) \geq 0 \quad \text{and} \quad h(x) = 0\}$$

is the constraint set.

To solve the problem (18.1) means to describe the following set

$$\{x^* \in C : \forall x \in C, f(x^*) \geq f(x)\}$$

which is called solution set to problem (18.1) and it is also denoted by $\arg \max$ (18.1). We will proceed as follows.

1. We will analyze in detail the problem with inequality constraints, i.e.,

$$\begin{array}{ll} \max_{x \in X} & f(x) \\ \text{s.t.} & g(x) \geq 0 \quad (1) \end{array}$$

2. We will analyze in detail the problem with equality constraints, i.e.,

$$\begin{array}{ll} \max_{x \in X} & f(x) \\ \text{s.t.} & h(x) = 0 \quad (2) \end{array}$$

3. We will describe how to solve the problem with both equality and inequality constraints, i.e.,

$$\begin{array}{ll} \max_{x \in X} & f(x) \\ \text{s.t.} & g(x) \geq 0 \quad (1) \\ & h(x) = 0 \quad (2) \end{array}$$

18.1 The case of inequality constraints: Kuhn-Tucker theorems

Consider the open and convex set $X \subseteq \mathbb{R}^n$ and the differentiable functions $f : X \rightarrow \mathbb{R}$, $g := (g^j)_{j=1}^m : X \rightarrow \mathbb{R}^m$. The problem we want to study is

$$\max_{x \in X} f(x) \quad \text{s.t.} \quad g(x) \geq 0. \quad (18.2)$$

Definition 727 The Kuhn-Tucker system (or conditions) associated with problem 18.2 is

$$\begin{cases} Df(x) + \lambda Dg(x) &= 0 & (1) \\ \lambda &\geq 0 & (2) \\ g(x) &\geq 0 & (3) \\ \lambda g(x) &= 0 & (4) \end{cases} \quad (18.3)$$

Equations (1) are called first order conditions; equations (2), (3) and (4) are called complementary slackness conditions.

Remark 728 $(x, \lambda) \in X \times \mathbb{R}^m$ is a solution to Kuhn-Tucker system iff it is a solution to any of the following systems:

1.
$$\begin{cases} \frac{\partial f(x)}{\partial x_i} + \sum_{j=1}^m \lambda_j \frac{\partial g_j(x)}{\partial x_i} &= 0 & \text{for } i = 1, \dots, n & (1) \\ \lambda_j &\geq 0 & \text{for } j = 1, \dots, m & (2) \\ g_j(x) &\geq 0 & \text{for } j = 1, \dots, m & (3) \\ \lambda_j g_j(x) &= 0 & \text{for } j = 1, \dots, m & (4) \end{cases}$$
2.
$$\begin{cases} Df(x) + \lambda Dg(x) &= 0 & (1) \\ \min \{\lambda_j, g_j(x)\} &= 0 & \text{for } j = 1, \dots, m & (2) \end{cases}$$

Moreover, $(x, \lambda) \in X \times \mathbb{R}^m$ is a solution to Kuhn-Tucker system iff it is a solution to

$$Df(x) + \lambda Dg(x) = 0 \quad (1)$$

and for each $j = 1, \dots, m$, to **one** of the following conditions

$$\begin{array}{ll} \text{either} & (\lambda_j > 0 \quad \text{and} \quad g_j(x) = 0) \\ \text{or} & (\lambda_j = 0 \quad \quad \quad g_j(x) > 0) \\ \text{or} & (\lambda_j = 0 \quad \quad \quad g_j(x) = 0) \end{array}$$

Definition 729 Given $x^* \in X$, we say that j is a binding constraint at x^* if $g_j(x^*) = 0$. Let

$$J^*(x^*) := \{j \in \{1, \dots, m\} : g_j(x^*) = 0\},$$

$$g^* := (g_j)_{j \in J^*(x^*)}, \quad \hat{g} := (g_j)_{j \notin J^*(x^*)}$$

and

$$m^* = \#J^*(x^*).$$

Definition 730 $x^* \in \mathbb{R}^n$ satisfies the constraint qualifications associated with problem 18.2 if it is a solution to

$$\max_{x \in \mathbb{R}^n} Df(x^*)x \quad \text{s.t.} \quad Dg^*(x^*)(x - x^*) \geq 0 \quad (18.4)$$

The above problem is obtained from 18.2

1. replacing g with g^* ;
2. linearizing f and g^* around x^* , i.e., substituting f and g^* with $f(x^*) + Df(x^*)(x - x^*)$ and $g(x^*) + Dg(x^*)(x - x^*)$, respectively;
3. dropping redundant terms, i.e., the term $f(x^*)$ in the objective function, and the term $g^*(x^*) = 0$ in the constraint.

Theorem 731 Suppose x^* is a solution to problem 18.2 and to problem 18.4, then there exists $\lambda^* \in \mathbb{R}^m$ such that (x^*, λ^*) satisfies Kuhn-Tucker conditions.

The proof of the above theorem requires the following lemma.

Lemma 732 (Farkas) *Given a matrix $A_{m \times n}$ and a vector $a \in \mathbb{R}^n$, either 1. there exists $\lambda \in \mathbb{R}_+^m$ such that $a = \lambda A$, or 2. there exists $y \in \mathbb{R}^n$ such that $Ay \geq 0$ and $ay < 0$, but not both.*

Proof. It follows immediately from Proposition 677. ■

Proof. of Theorem 731

(main steps: 1. use the fact x^* is a solution to problem 18.4; 2. apply Farkas Lemma; 3. choose $\lambda^* = (\lambda \text{ from Farkas}, 0)$).

Since x^* is a solution to problem 18.4, for any $x \in \mathbb{R}^n$ such that $Dg^*(x^*)(x - x^*) \geq 0$ it is the case that $Df(x^*)x^* \geq Df(x^*)x$ or

$$Dg^*(x^*)(x - x^*) \geq 0 \Rightarrow [-Df(x^*)(x - x^*)] \geq 0. \quad (18.5)$$

Applying Farkas Lemma identifying

$$a \quad \text{with} \quad -Df(x^*)$$

and

$$A \quad \text{with} \quad Dg^*(x^*)$$

we have that either

1. there exists $\lambda \in \mathbb{R}_+^m$ such that

$$-Df(x^*) = \lambda Dg^*(x^*) \quad (18.6)$$

or 2. there exists $y \in \mathbb{R}^n$ such that

$$Dg^*(x^*)y \geq 0 \quad \text{and} \quad -Df(x^*)y < 0 \quad (18.7)$$

but not both 1 and 2.

Choose $x = y + x^*$ and therefore you have $y = x - x^*$. Then, 18.7 contradicts 18.5. Therefore,

1. above holds.

Now, choose $\lambda^* := (\lambda, 0) \in \mathbb{R}^{m^*} \times \mathbb{R}^{m-m^*}$, we have that

$$Df(x^*) + \lambda^* Dg(x^*) = Df(x^*) + (\lambda, 0) \begin{pmatrix} Dg^*(x^*) \\ D\hat{g}(x^*) \end{pmatrix} = Df(x^*) + \lambda Dg(x^*) = 0$$

where the last equality follows from 18.6;

$\lambda^* \geq 0$ by Farkas Lemma;

$g(x^*) \geq 0$ from the assumption that x^* solves problem 18.2;

$\lambda^* g(x^*) = (\lambda, 0) \begin{pmatrix} g^*(x^*) \\ \hat{g}(x^*) \end{pmatrix} = \lambda g^*(x) = 0$, where the last equality follows from the definition

of g^* . ■

Theorem 733 *If x^* is a solution to problem (18.2) and*

either for $j = 1, \dots, m$, g_j is pseudo-concave and $\exists x^{++} \in X$ such that $g(x^{++}) \gg 0$,

or $\text{rank } Dg^(x^*) = m^* := \#J^*(x^*)$,*

then x^ solves problem (18.4).*

Proof. We prove the conclusion of the theorem under the first set of conditions.

Main steps: 1. suppose otherwise: $\exists \tilde{x} \dots$; 2. use the two assumptions; 3. move from x^* in the direction $x^\theta := (1 - \theta)\tilde{x} + \theta x^{++}$.

Suppose that the conclusion of the theorem is false. Then there exists $\tilde{x} \in \mathbb{R}^n$ such that

$$Dg^*(x^*)(\tilde{x} - x^*) \geq 0 \quad \text{and} \quad Df(x^*)(\tilde{x} - x^*) > 0 \quad (18.8)$$

Moreover, from the definition of g^* and x^{++} , we have that

$$g^*(x^{++}) \gg 0 = g^*(x^*)$$

Since for $j = 1, \dots, m$, g_j is pseudo-concave we have that

$$Dg^*(x^*)(x^{++} - x^*) \gg 0 \quad (18.9)$$

Define

$$x^\theta := (1 - \theta)\tilde{x} + \theta x^{++}$$

with $\theta \in (0, 1)$. Observe that

$$x^\theta - x^* = (1 - \theta)\tilde{x} + \theta x^{++} - (1 - \theta)x^* - \theta x^* = (1 - \theta)(\tilde{x} - x^*) + \theta(x^{++} - x^*)$$

Therefore,

$$Dg^*(x^*)(x^\theta - x^*) = (1 - \theta)Dg^*(x^*)(\tilde{x} - x^*) + \theta Dg^*(x^*)(x^{++} - x^*) \gg 0 \quad (18.10)$$

where the last equality come from 18.8 and 18.9.

Moreover,

$$Df(x^*)(x^\theta - x^*) = (1 - \theta)Df(x^*)(\tilde{x} - x^*) + \theta Df(x^*)(x^{++} - x^*) \gg 0 \quad (18.11)$$

where the last equality come from 18.8 and a choice of θ sufficiently small.¹

Observe that from Remark 611, 18.10 and 18.11 we have that

$$(g^*)'(x^*, x^\theta) \gg 0$$

and

$$f'(x^*, x^\theta) > 0$$

Therefore, using the fact that X is open, and that $\hat{g}(x^*) \gg 0$, there exists γ such that

$$\begin{aligned} x^* + \gamma(x^\theta - x^*) &\in X \\ g^*(x^* + \gamma(x^\theta - x^*)) &\gg g^*(x^*) = 0 \\ f^*(x^* + \gamma(x^\theta - x^*)) &> f(x^*) \\ \hat{g}(x^* + \gamma(x^\theta - x^*)) &\gg 0 \end{aligned} \quad (18.12)$$

But then 18.12 contradicts the fact that x^* solves problem (18.2). ■

From Theorems 731 and 733, we then get the following corollary.

Theorem 734 Suppose x^* is a solution to problem 18.2, and one of the following constraint qualifications hold:

- a. for $j = 1, \dots, m$, g_j is pseudo-concave and there exists $x^{++} \in X$ such that $g(x^{++}) \gg 0$
- b. $\text{rank } Dg^*(x^*) = \#J^*$,

Then there exists $\lambda^* \in \mathbb{R}^m$ such that (x^*, λ^*) solves the system 18.3.

Theorem 735 If f is pseudo-concave, and for $j = 1, \dots, m$, g_j is quasi-concave, and (x^*, λ^*) solves the system 18.3, then x^* solves problem 18.2.

Proof. Main steps: 1. suppose otherwise and use the fact that f is pseudo-concave; 2. for $j \in J^*(x^*)$, use the quasi-concavity of g_j ; 3. for $j \in J^*(x^*)$, use (second part of) kuhn-Tucker conditions; 4. Observe that 2. and 3. above contradict the first part of Kuhn-Tucker conditions.)

Suppose otherwise, i.e., there exists $\hat{x} \in X$ such that

$$g(\hat{x}) \geq 0 \quad \text{and} \quad f(\hat{x}) > f(x^*) \quad (18.13)$$

¹ Assume that $\theta \in (0, 1)$, $\alpha \in \mathbb{R}_{++}$ and $\beta \in \mathbb{R}$. We want to show that there exist $\theta^* \in (0, 1)$ such that

$$(1 - \theta)\alpha + \theta\beta > 0$$

i.e.,

$$\alpha > \theta(\alpha - \beta)$$

If $(\alpha - \beta) = 0$, the claim is true.

If $(\alpha - \beta) > 0$, any $\theta < \frac{\alpha}{\alpha - \beta}$ will work (observe that $\frac{\alpha}{\alpha - \beta} > 0$).

If $(\alpha - \beta) < 0$, the claim is clearly true because $0 < \alpha$ and $\theta(\alpha - \beta) < 0$.

From 18.13 and the fact that f pseudo-concave, we get

$$Df(x^*)(\hat{x} - x^*) > 0 \quad (18.14)$$

From 18.13, the fact that $g^*(x^*) = 0$ and that g_j is quasi-concave, we get that

$$\text{for } j \in J^*(x^*), \quad Dg^j(x^*)(\hat{x} - x^*) \geq 0$$

and since $\lambda^* \geq 0$,

$$\text{for } j \in J^*(x^*), \quad \lambda_j^* Dg^j(x^*)(\hat{x} - x^*) \geq 0 \quad (18.15)$$

For $j \in \hat{J}(x^*)$, from Kuhn-Tucker conditions, we have that $g_j(x^*) > 0$ and $\lambda_j^* = 0$, and therefore

$$\text{for } j \in \hat{J}(x^*), \quad \lambda_j^* Dg^j(x^*)(\hat{x} - x^*) = 0 \quad (18.16)$$

But then from 18.14, 18.15 and 18.16, we have

$$Df(x^*)(\hat{x} - x^*) + \lambda^* Dg(x^*)(\hat{x} - x^*) > 0$$

contradicting Kuhn-Tucker conditions. ■

We can summarize the above results as follows. Call (M) the problem

$$\max_{x \in X} f(x) \quad \text{s.t.} \quad g(x) \geq 0 \quad (18.17)$$

and define

$$M := \arg \max (M) \quad (18.18)$$

$$S := \{x \in X : \exists \lambda \in \mathbb{R}^m \text{ such that } (x, \lambda) \text{ is a solution to Kuhn-Tucker system (18.3)}\} \quad (18.19)$$

1. Assume that one of the following conditions hold:

- (a) for $j = 1, \dots, m$, g_j is pseudo-concave and there exists $x^{++} \in X$ such that $g(x^{++}) \gg 0$
- (b) $\text{rank } Dg^*(x^*) = \#J^*$.

Then

$$x^* \in M \Rightarrow x^* \in S$$

2. Assume that both the following conditions hold:

- (a) f is pseudo-concave, and
- (b) for $j = 1, \dots, m$, g_j is quasi-concave.

Then

$$x^* \in S \Rightarrow x^* \in M.$$

18.1.1 On uniqueness of the solution

The following proposition is a useful tool to show uniqueness.

Proposition 736 *The solution to problem*

$$\max_{x \in X} f(x) \quad \text{s.t.} \quad g(x) \geq 0 \quad (P)$$

either does not exist or it is unique if one of the following conditions holds

1. f is strictly quasi-concave, and
for $j \in \{1, \dots, m\}$, g_j is quasi-concave;
2. f is quasi-concave and locally non-satiated (i.e., $\forall x \in X, \forall \varepsilon > 0$, there exists $x' \in B(x, \varepsilon)$ such that $f(x') > f(x)$), and
for $j \in \{1, \dots, m\}$, g_j is strictly quasi-concave.

Proof. 1.

Since g_j is quasi concave $V^j := \{x \in X : g_j(x) \geq 0\}$ is convex. Since the intersection of convex sets is convex $V = \bigcap_{j=1}^m V^j$ is convex.

Suppose that both x' and x'' are solutions to problem (P) and $x' \neq x''$. Then for any $\lambda \in (0, 1)$,

$$(1 - \lambda)x' + \lambda x'' \in V \quad (18.20)$$

because V is convex, and

$$f((1 - \lambda)x' + \lambda x'') > \min\{f(x'), f(x'')\} = f(x') = f(x'') \quad (18.21)$$

because f is strictly-quasi-concave.

But (18.20) and (18.21) contradict the fact that x' and x'' are solutions to problem (P).

2.

Observe that V is strictly convex because each V^j is strictly convex. Suppose that both x' and x'' are solutions to problem (P) and $x' \neq x''$. Then for any $\lambda \in (0, 1)$,

$$x(\lambda) := (1 - \lambda)x' + \lambda x'' \in \text{Int } V$$

i.e., $\exists \varepsilon > 0$ such that $B(x(\lambda), \varepsilon) \subseteq V$. Since f is locally non-satiated, there exists $x' \in B(x(\lambda), \varepsilon) \subseteq V$ such that

$$f(\hat{x}) > f(x(\lambda)) \quad (18.22)$$

Since f is quasi-concave,

$$f(x(\lambda)) \geq f(x') = f(x'') \quad (18.23)$$

(18.22) and (18.23) contradict the fact that x' and x'' are solutions to problem (P). ■

Remark 737 1. If f is strictly increasing (i.e., $\forall x', x'' \in X$ such that $x' > x''$, we have that $f(x') > f(x'')$) or strictly decreasing, then f is locally non-satiated.

2. If f is affine and not constant, then f is quasi-concave and Locally NonSatiated.

Proof of 2.

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ affine and not constant means that there exists $a \in \mathbb{R}$ and $b \in \mathbb{R} \setminus \{0\}$ such that $f : x \mapsto a + b^T x$. Take an arbitrary \bar{x} and $\varepsilon > 0$. For $i \in \{1, \dots, n\}$, define $\alpha_i := \frac{\varepsilon}{k} \cdot (\text{sign } b_i)$ and $\tilde{x} := \bar{x} + (\alpha_i)_{i=1}^n$, with $k \neq 0$ and which will be computed below. Then

$$f(\tilde{x}) = a + b\bar{x} + \sum_{i=1}^n \frac{\varepsilon}{k} |b_i| > f(\bar{x});$$

$$\|\tilde{x} - \bar{x}\| = \left\| \frac{\varepsilon}{k} \cdot ((\text{sign } b_i) \cdot b_i)_{i=1}^n \right\| = \frac{\varepsilon}{k} \cdot \left\| \sqrt{\sum_{i=1}^n (b_i)^2} \right\| = \frac{\varepsilon}{k} \cdot \|b\| < \varepsilon \text{ if } k > \frac{1}{\|b\|}.$$

Remark 738 In part 2 of the statement of the Proposition f **has** to be both quasi-concave and Locally NonSatiated.

a. Example of f quasi-concave (and g_j strictly-quasi-concave) with more than one solution:

$$\max_{x \in \mathbb{R}} 1 \quad \text{s.t.} \quad x + 1 \geq 0 \quad 1 - x \geq 0$$

The set of solution is $[-1, +1]$

a. Example of f Locally NonSatiated (and g_j strictly-quasi-concave) with more than one solution:

$$\max_{x \in \mathbb{R}} x^2 \quad \text{s.t.} \quad x + 1 \geq 0 \quad 1 - x \geq 0$$

The set of solutions is $\{-1, +1\}$.

18.2 The Case of Equality Constraints: Lagrange Theorem.

Consider the C^1 functions

$$f : X \rightarrow \mathbb{R}, \quad f : x \mapsto f(x),$$

$$g : X \rightarrow \mathbb{R}^m, \quad g : x \mapsto g(x) := (g_j(x))_{j=1}^m$$

with $\underline{m \leq n}$. Consider also the following “” maximization problem:

$$(P) \quad \max_{x \in X} f(x) \quad s.t. \quad g(x) = 0 \quad (18.24)$$

$$\mathcal{L} : X \times \mathbb{R}^m \rightarrow \mathbb{R}, \quad \mathcal{L} : (x, \lambda) \mapsto f(x) + \lambda g(x)$$

is called Lagrange function associated with problem (15.15).

We recall below the statement of Theorem 658.

Theorem 739 (*Necessary Conditions*)

Assume that $\text{rank}[Dg(x^*)] = m$.

Under the above condition, we have that

x^* is a local maximum for (P)

\Rightarrow

there exists $\lambda^* \in \mathbb{R}^m$, such that

$$\begin{cases} Df(x^*) + \lambda^* Dg(x^*) = 0 \\ g(x^*) = 0 \end{cases} \quad (18.25)$$

Remark 740 The full rank condition in the above Theorem cannot be dispensed. The following example shows a case in which x^* is a solution to maximization problem (18.24), $Dg(x^*)$ does not have full rank and there exists no λ^* satisfying Condition 18.25. Consider

$$\max_{(x,y) \in \mathbb{R}^2} x \quad s.t. \quad \begin{aligned} x^3 - y &= 0 \\ x^3 + y &= 0 \end{aligned}$$

The constraint set is $\{(0,0)\}$ and therefore the solution is just $(x^*, y^*) = (0,0)$. The Jacobian matrix of the constraint function is

$$\begin{bmatrix} 3x^2 & -1 \\ 3x^2 & 1 \end{bmatrix}_{|(x^*, y^*)} = \begin{bmatrix} 0 & -1 \\ 0 & 1 \end{bmatrix}$$

which does not have full rank.

$$\begin{aligned} (0,0) &= Df(x^*, y^*) + (\lambda_1, \lambda_2) Dg(x^*, y^*) = \\ &= (1,0) + (\lambda_1, \lambda_2) \begin{bmatrix} 0 & -1 \\ 0 & 1 \end{bmatrix} = (1, -\lambda_1 + \lambda_2), \end{aligned}$$

from which it follows that there exists no λ^* solving the above system.

Theorem 741 (*Sufficient Conditions*)

Assume that

1. f is pseudo-concave,

2. for $j = 1, \dots, m$, g_j is quasi concave.

Under the above conditions, we have what follows.

[there exist $(x^*, \lambda^*) \in X \times \mathbb{R}^m$ such that

3. $\lambda^* \geq 0$,

4. $Df(x^*) + \lambda^* Dg(x^*) = 0$, and

5. $g(x^*) = 0$]

\Rightarrow

x^* solves (P).

Proof.

Suppose otherwise, i.e., there exists $\hat{x} \in X$ such that

$$\text{for } j = 1, \dots, m, \quad g_j(\hat{x}) = g_j(x^*) = 0 \quad (1), \text{ and}$$

$$f(\hat{x}) > f(x^*) \quad (2).$$

Quasi-concavity of g_j and (1) imply that

$$Dg^j(x^*)(\hat{x} - x^*) \geq 0 \quad (3).$$

Pseudo concavity of f and (2) imply that

$$Df(x^*)(\hat{x} - x^*) > 0 \quad (4).$$

But then

$$0 \stackrel{\text{Assumption}}{=} [Df(x^*) + \lambda Dg(x^*)](\hat{x} - x^*) = Df(x^*)(\hat{x} - x^*) + \frac{(\geq 0)}{\lambda} Dg(x^*)(\hat{x} - x^*) > 0,$$

a contradiction.

■

18.3 The Case of Both Equality and Inequality Constraints.

Consider

the open and convex set $X \subseteq \mathbb{R}^n$ and the differentiable functions $f : X \rightarrow \mathbb{R}$, $g : X \rightarrow \mathbb{R}^m$, $h : X \rightarrow \mathbb{R}^l$.

Consider the problem

$$\max_{x \in X} f(x) \quad s.t. \quad \begin{array}{l} g(x) \geq 0 \\ h(x) = 0. \end{array} \quad (18.26)$$

Observe that

$$h(x) = 0 \Leftrightarrow$$

$$\Leftrightarrow \text{for } k = 1, \dots, l, \quad h^k(x) = 0 \Leftrightarrow$$

$$\Leftrightarrow \text{for } k = 1, \dots, l, \quad h^{k1}(x) := h^k(x) \geq 0 \text{ and } h^{k2}(x) := -h^k(x) \geq 0.$$

Defined $h^1(x) := (h^{k1}(x))_{k=1}^l$ and $h^2(x) := (h^{k2}(x))_{k=1}^l$, problem 18.26 with associated multipliers can be rewritten as

$$\max_{x \in X} f(x) \quad s.t. \quad \begin{array}{l} g(x) \geq 0 \\ h^1(x) \geq 0 \\ h^2(x) \geq 0 \end{array} \quad \begin{array}{l} \lambda \\ \mu_1 \\ \mu_2 \end{array} \quad (18.27)$$

The Lagrangian function of the above problem is

$$\mathcal{L}(x; \lambda, \mu_1, \mu_2) = f(x) + \lambda^T g(x) + (\mu_1 - \mu_2)^T h(x) =$$

$$= f(x) + \sum_{j=1}^m \lambda_j g_j(x) + \sum_{k=1}^l (\mu_1^k - \mu_2^k)^T h(x),$$

and the Kuhn-Tucker Conditions are:

$$\begin{array}{l} Df(x) + \lambda^T Dg(x) + (\mu_1 - \mu_2)^T Dh(x) = 0 \\ g_j(x) \geq 0, \lambda_j \geq 0, \lambda_j g_j(x) = 0, \\ h^k(x) = 0, (\mu_1^k - \mu_2^k) := \mu \geq 0, \end{array} \quad \begin{array}{l} \text{for } j = 1, \dots, m, \\ \text{for } k = 1, \dots, l. \end{array} \quad (18.28)$$

Theorem 742 Assume that f, g and h are C^2 functions and that

$$\text{either rank} \begin{bmatrix} Dg^*(x^*) \\ Dh(x^*) \end{bmatrix} = m^* + l,$$

or for $j = 1, \dots, m$, $-g_j$ is pseudoconcave, and $\forall k$, h_k and $-h_k$ are pseudoconcave

Under the above conditions,

if x^* solves 18.26, then $\exists (x^*, \lambda^*, \mu^*) \in X \times \mathbb{R}^m \times \mathbb{R}^l$ which satisfies the associated Kuhn-Tucker conditions.

Proof. The above conditions are called “Weak reverse convex constraint qualification” (Mangasarian (1969)) or “Reverse constraint qualification” (Bazaraa and Shetty (1976)). The needed result is presented and proved in

Mangasarian², - see 4, page 172 and Theorem 6, page 173, and Bazaraa and Shetty (1976) - see 7 page 148, and theorems 6.2.3, page 148 and Theorem 6.2.4, page 150.

See also El-Hodiri (1991), Theorem 1, page 48 and Simon (1985), Theorem 4.4. (iii), page 104.

■

■

Remark 743 For other conditions, see Theorem 5.8, page 124, in Jahn (1996).

Theorem 744 Assume that

f is pseudo-concave, and

for $j = 1, \dots, m$, g_j is quasi-concave, and for $k = 1, \dots, l$, h^k is quasi-concave and $-h^k$ is quasi-concave.

Under the above conditions,

if $(x^*, \lambda^*, \mu^*) \in X \times \mathbb{R}^m \times \mathbb{R}^l$ satisfies the Kuhn-Tucker conditions associated with 18.26, then x^* solves 18.26.

Proof.

This follows from Theorems proved in the case of inequality constraints.

■

Similarly, to what we have done in previous sections, we can summarize what said above as follows.

Call (M_2) the problem

$$\max_{x \in X} f(x) \quad s.t. \quad \begin{aligned} g(x) &\geq 0 \\ h(x) &= 0. \end{aligned} \quad (18.29)$$

and define

$$M_2 := \arg \max (M_2)$$

$$S_2 := \{x \in X : \exists \lambda \in \mathbb{R}^m \text{ such that } (x, \lambda) \text{ is a solution to Kuhn-Tucker system (18.28)}\}$$

1. Assume that one of the following conditions hold:

- (a) $\text{rank} \begin{bmatrix} Dg^*(x^*) \\ Dh(x^*) \end{bmatrix} = m^* + l$, or
- (b) for $j = 1, \dots, m$, g_j is linear, and $h(x)$ is affine.

Then

$$M_2 \subseteq S_2$$

2. Assume that both the following conditions hold:

- (a) f is pseudo-concave, and
- (b) for $j = 1, \dots, m$, g_j is quasi-concave, and for $k = 1, \dots, l$, h^k is quasi-concave and $-h^k$ is quasi-concave.

Then

$$M_2 \supseteq S_2$$

²What Mangasarian calls a linear function is what we call an affine function.

18.4 Main Steps to Solve a (Nice) Maximization Problem

We have studied the problem

$$\max_{x \in X} f(x) \quad s.t. \quad g(x) \geq 0 \quad (M)$$

which we call a maximization problem in the “canonical form”, i.e., a maximization problem with constraints in the form of “ \geq ”, and we have defined

$$M := \arg \max (M)$$

$$C := \{x \in X : g(x) \geq 0\}$$

$$S := \{x \in X : \exists \lambda \in \mathbb{R}^m \text{ such that } (x, \lambda) \text{ satisfies Kuhn-Tucker Conditions (18.3)}\}$$

Recall that X is an open, convex subset of \mathbb{R}^n , $f : X \rightarrow \mathbb{R}$, $\forall j \in \{1, \dots, m\}$, $g_j : X \rightarrow \mathbb{R}$ and $g := (g_j)_{j=1}^m : X \rightarrow \mathbb{R}^m$.

In many cases, we have to study the following problem

$$\max_x f(x) \quad s.t. \quad g(x) \geq 0, \quad (M')$$

in which the set X is not specified.

We list the main steps to try to solve (M') .

1. Canonical form.

Write the problem in the (in fact, our definition of) canonical form. Sometimes the problem contains a parameter $\pi \in \Pi$ an open subset of \mathbb{R}^k . Then we should write: for given $\pi \in \Pi$

$$\max_x f(x, \pi) \quad s.t. \quad g(x, \pi) \geq 0.$$

2. The set X and the functions f and g .

a. Define the functions \tilde{f} , \tilde{g} naturally arising from the problem with domain equal to their definition set, where the definition set of a function φ is the largest set \mathcal{D}_φ which can be the domain of that function.

b. Determine X . A possible choice for X is the intersection of the “definition set” of each function, i.e.,

$$X = \mathcal{D}_f \cap \mathcal{D}_{g_1} \cap \dots \cap \mathcal{D}_{g_m}$$

c. Check if X is open and convex.

d. To apply the analysis described in the previous sections, show, if possible, that f and g are of class C^2 or at least C^1 .

3. Existence.

Try to apply the Extreme Value Theorem. If f is at least C^1 , then f is continuous and therefore we have to check if the constraint set C is non-empty and compact. Recall that a set S in \mathbb{R}^n is compact if and only if S is closed (in \mathbb{R}^n) and bounded.

Boundedness has to be shown “brute force”, i.e., using the specific form of the maximization problem.

If $X = \mathbb{R}^n$, then $C := \{x \in X : g(x) \geq 0\}$ is closed, because of the following well-known argument:

$C = \bigcap_{j=1}^m g_j^{-1}([0, +\infty))$; since g_j is C^2 (or at least C^1) and therefore continuous, and $[0, +\infty)$ closed, $g_j^{-1}([0, +\infty))$ is closed in $X = \mathbb{R}^n$; then C is closed because intersection of closed sets.

A problem may arise if X is an open proper subset of \mathbb{R}^n . In that case the above argument shows that C is a closed set in $X \neq \mathbb{R}^n$ and therefore it is not necessarily closed in \mathbb{R}^n . A possible way out is the following one.

Verify that while the definition set of f is X , the definition set of g is \mathbb{R}^n . If $\mathcal{D}_{\tilde{g}} = \mathbb{R}^n$, then from the above argument

$$\tilde{C} := \{x \in \mathbb{R}^n : \tilde{g}(x) \geq 0\}$$

is closed (in \mathbb{R}^n). Observe that

$$C = \tilde{C} \cap X.$$

Then, we are left with showing that $\tilde{C} \subseteq X$ and therefore $\tilde{C} \cap X = \tilde{C}$ and then

$$C = \tilde{C}$$

If \tilde{C} is compact, C is compact as well.³

4. Number of solutions.

See subsection 18.1.1. In fact, summarizing what said there, we know that the solution to (M) , if any, is unique if

1. f is strictly-quasi-concave, and for $j \in \{1, \dots, m\}$, g_j is quasi-concave; **or**
2. for $j \in \{1, \dots, m\}$, g_j is strictly-quasi-concave and either a. f is quasi-concave and locally non-satiated, or b. f is affine and non-constant, or c. f is quasi-concave and strictly monotone, or d. f is quasi-concave and $\forall x \in X, Df(x) >> 0$, or e. f is quasi-concave and $\forall x \in X, Df(x) << 0$.

5. Necessity of K-T conditions.

Check if the conditions which insure that $M \subseteq S$ hold, i.e.,

- either a. for $j = 1, \dots, m$, g_j is pseudo-concave and there exists $x^{++} \in X$ such that $g(x^{++}) \gg 0$,
- or b. $\text{rank } Dg^*(x^*) = \#J^*$.

If those conditions holds, each property we show it holds for elements of S does hold *a fortiori* for elements of M .

6. Sufficiency of K-T conditions.

Check if the conditions which insure that $M \supseteq S$ hold, i.e., that

f is pseudo-concave and for $j = 1, \dots, m$, g_j is quasi-concave.

If those conditions holds, each property we show it does *not* hold for elements of S does not hold *a fortiori* for elements of M .

7. K-T conditions.

Write the Lagrangian function and then the Kuhn-Tucker conditions.

8. Solve the K-T conditions.

Try to solve the system of Kuhn-Tucker conditions in the unknown variables (x, λ) . To do that; either, analyze all cases,

or, try to get a “good conjecture” and check if the conjecture is correct.

Example 745 *Discuss the problem*

$$\max_{(x_1, x_2)} \frac{1}{2} \log(1 + x_1) + \frac{1}{3} \log(1 + x_2) \quad s.t. \quad \begin{array}{rcl} x_1 & \geq & 0 \\ x_2 & \geq & 0 \\ x_1 + x_2 & \leq & w \end{array}$$

with $w > 0$.

1. Canonical form.

For given $w \in \mathbb{R}_{++}$,

$$\max_{(x_1, x_2)} \frac{1}{2} \log(1 + x_1) + \frac{1}{3} \log(1 + x_2) \quad s.t. \quad \begin{array}{rcl} x_1 & \geq & 0 \\ x_2 & \geq & 0 \\ w - x_1 - x_2 & \geq & 0 \end{array} \quad (18.30)$$

2. The set X and the functions f and g .

³Observe that the above argument does not apply to the case in which

$$C = \{x \in \mathbb{R}_{++}^2 : w - x_1 - x_2 \geq 0\} :$$

in that case,

$$\tilde{C} = \{x \in \mathbb{R}^2 : w - x_1 - x_2 \geq 0\} \not\subseteq \mathbb{R}_{++}^2.$$

a.

$$\begin{aligned}
\tilde{f} : (-1, +\infty)^2 &\rightarrow \mathbb{R}, & (x_1, x_2) &\mapsto \frac{1}{2} \log(1+x_1) + \frac{1}{3} \log(1+x_2) \\
\tilde{g}_1 : \mathbb{R}^2 &\rightarrow \mathbb{R} & (x_1, x_2) &\mapsto x_1 \\
\tilde{g}_2 : \mathbb{R}^2 &\rightarrow \mathbb{R} & (x_1, x_2) &\mapsto x_2 \\
\tilde{g}_3 : \mathbb{R}^2 &\rightarrow \mathbb{R} & (x_1, x_2) &\mapsto w - x_1 - x_2
\end{aligned}$$

b.

$$X = (-1, +\infty)^2$$

and therefore f and g are just \tilde{f} and \tilde{g} restricted to X .

c. X is open and convex because Cartesian product of open intervals which are open, convex sets.

d. Let's try to compute the Hessian matrices of f, g_1, g_2, g_3 . Gradients are

$$\begin{aligned}
Df(x_1, x_2) &= \left(\frac{1}{2(x_1+1)}, \frac{1}{3(x_2+1)} \right) \\
D\tilde{g}_1(x_1, x_2) &= (1, 0) \\
D\tilde{g}_2(x_1, x_2) &= (0, 1) \\
D\tilde{g}_3(x_1, x_2) &= (-1, -1)
\end{aligned}$$

Hessian matrices are

$$\begin{aligned}
D^2f(x_1, x_2) &= \begin{bmatrix} -\frac{1}{2(x_1+1)^2} & 0 \\ 0 & -\frac{1}{3(x_2+1)^2} \end{bmatrix} \\
D^2\tilde{g}_1(x_1, x_2) &= 0 \\
D^2\tilde{g}_2(x_1, x_2) &= 0 \\
D^2\tilde{g}_3(x_1, x_2) &= 0
\end{aligned}$$

In fact, g_1, g_2, g_3 are affine functions. In conclusion, f and g_1, g_2, g_3 are C^2 . In fact, g_1 and g_2 are linear and g_3 is affine.

3. Existence.

C is clearly bounded: $\forall x \in C$,

$$(0, 0) \leq (x_1, x_2) \leq (w, w)$$

In fact, the first two constraint simply say that $(x_1, x_2) \geq (0, 0)$. Moreover, from the third constraint $x_1 \leq w - x_2 \leq w$, simply because $x_2 \geq 0$; similar argument can be used to show that $x_2 \leq w$.

To show closedness, use the strategy proposed above.

$$\tilde{C} := \{x \in \mathbb{R}^n : g(x) \geq 0\}$$

is obviously closed. Since $\tilde{C} \subseteq \mathbb{R}_+^2$, because of the first two constraints, $\tilde{C} \subseteq X := (-1, +\infty)^2$ and therefore $C = \tilde{C} \cap X = \tilde{C}$ is closed.

We can then conclude that C is compact and therefore $\arg \max (18.30) \neq \emptyset$.

4. Number of solutions.

From the analysis of the Hessian and using Theorem 723, parts 1 and 2, we have that f is strictly concave:

$$\begin{aligned}
&-\frac{1}{2(x_1+1)^2} < 0 \\
\det \begin{bmatrix} -\frac{1}{2(x_1+1)^2} & 0 \\ 0 & -\frac{1}{3(x_2+1)^2} \end{bmatrix} &= \frac{1}{2(x_1+1)^2} \cdot \frac{1}{3(x_2+1)^2} > 0
\end{aligned}$$

Moreover g_1, g_2, g_3 are affine and therefore concave. From Proposition 736, part 1, the solution is unique.

5. Necessity of K-T conditions.

Since each g_j is affine and therefore pseudo-concave, we are left with showing that there exists $x^{++} \in X$ such that $g(x^{++}) \gg 0$. Just take $(x_1^{++}, x_2^{++}) = \frac{w}{4}(1, 1)$:

$$\begin{aligned}
\frac{w}{4} &> 0 \\
\frac{w}{4} &> 0 \\
w - \frac{w}{4} - \frac{w}{4} = \frac{w}{2} &> 0
\end{aligned}$$

Therefore

$$M \subseteq S$$

6. Sufficiency of K-T conditions.

f is strictly concave and therefore pseudo-concave, and each g_j is linear and therefore quasi-concave. Therefore

$$M \supseteq S$$

7. K-T conditions.

$$\mathcal{L}(x_1, x_2, \lambda_1, \lambda_2, \mu; w) = \frac{1}{2} \log(1 + x_1) + \frac{1}{3} \log(1 + x_2) + \lambda_1 x_1 + \lambda_2 x_2 + \mu(w - x_1 - x_2)$$

$$\begin{cases} \frac{1}{2(x_1+1)} + \lambda_1 - \mu & = 0 \\ \frac{1}{3(x_2+1)} + \lambda_2 - \mu & = 0 \\ \min\{x_1, \lambda_1\} & = 0 \\ \min\{x_2, \lambda_2\} & = 0 \\ \min\{w - x_1 - x_2, \mu\} & = 0 \end{cases}$$

8. Solve the K-T conditions.

Conjecture: $x_1 > 0$ and therefore $\lambda_1 = 0$; $x_2 > 0$ and therefore $\lambda_2 = 0$; $w - x_1 - x_2 = 0$. The Kuhn-Tucker system becomes:

$$\begin{cases} \frac{1}{2(x_1+1)} - \mu & = 0 \\ \frac{1}{3(x_2+1)} - \mu & = 0 \\ w - x_1 - x_2 & = 0 \\ \mu & \geq 0 \\ x_1 > 0, x_2 > 0 \\ \lambda_1 = 0, \lambda_2 = 0 \end{cases}$$

Then,

$$\begin{cases} \frac{1}{2(x_1+1)} & = \mu \\ \frac{1}{3(x_2+1)} & = \mu \\ w - x_1 - x_2 & = 0 \\ \mu & > 0 \\ x_1 > 0, x_2 > 0 \\ \lambda_1 = 0, \lambda_2 = 0 \end{cases}$$

$$\begin{cases} x_1 & = \frac{1}{2\mu} - 1 \\ x_2 & = \frac{1}{3\mu} - 1 \\ w - \left(\frac{1}{2\mu} - 1\right) - \left(\frac{1}{3\mu} - 1\right) & = 0 \\ \mu & > 0 \\ x_1 > 0, x_2 > 0 \\ \lambda_1 = 0, \lambda_2 = 0 \end{cases}$$

$$0 = w - \left(\frac{1}{2\mu} - 1\right) - \left(\frac{1}{3\mu} - 1\right) = w - \frac{5}{6\mu} + 2; \text{ and } \mu = \frac{5}{6(w+2)} > 0. \text{ Then } x_1 = \frac{1}{2\mu} - 1 = \frac{6(w+2)}{2 \cdot 5} - 1 = \frac{3w+6-5}{5} = \frac{3w+1}{5} \text{ and } x_2 = \frac{1}{3\mu} - 1 = \frac{6(w+2)}{3 \cdot 5} - 1 = \frac{2w+4-5}{5} = \frac{2w-1}{5}.$$

Summarizing

$$\begin{cases} x_1 & = \frac{3w+1}{5} > 0 \\ x_2 & = \frac{2w-1}{5} > 0 \\ \mu & = \frac{5}{6(w+2)} > 0 \\ \lambda_1 = 0, \lambda_2 = 0 \end{cases}$$

Observe that while $x_1 > 0$ for any value of w , $x_2 > 0$ iff $w > \frac{1}{2}$. Therefore, for $w \in (0, \frac{1}{2}]$, the above one is not a solution, and we have to come up with another conjecture;

$x_1 = w$ and therefore $\lambda_1 = 0$; $x_2 = 0$ and $\lambda_2 \geq 0$; $w - x_1 - x_2 = 0$ and $\mu \geq 0$. The Kuhn-Tucker conditions become

$$\begin{cases} \frac{1}{2(w+1)} - \mu = 0 \\ \frac{1}{3} + \lambda_2 - \mu = 0 \\ \lambda_1 = 0 \\ x_2 = 0 \\ \lambda_2 \geq 0 \\ x_1 = w \\ \mu \geq 0 \end{cases}$$

and


$$\begin{cases} \mu = \frac{1}{2(w+1)} > 0 \\ \lambda_2 = \frac{1}{2(w+1)} - \frac{1}{3} = \frac{3-2w-2}{6(w+1)} = \frac{1-2w}{6(w+1)} \\ \lambda_1 = 0 \\ x_2 = 0 \\ \lambda_2 \geq 0 \\ x_1 = w \end{cases}$$

$\lambda_2 = \frac{1-2w}{6(w+1)} = 0$ if $w = \frac{1}{2}$, and $\lambda_2 = \frac{1-2w}{6(w+1)} > 0$ if $w \in (0, \frac{1}{2})$

Summarizing, the unique solution x^* to the maximization problem is


$$\begin{array}{lll} \text{if } w \in (0, \frac{1}{2}), & \text{then } x_1^* = w, & \lambda_1^* = 0 \text{ and } x_2^* = 0, \quad \lambda_2^* > 0 \\ \text{if } w = \frac{1}{2}, & \text{then } x_1^* = w, & \lambda_1^* = 0 \quad x_2^* = 0, \quad \lambda_2^* = 0 \\ \text{if } w \in (\frac{1}{2}, +\infty), & \text{then } x_1^* = \frac{3w+1}{5} > 0 & \lambda_1^* = 0 \text{ and } x_2^* = \frac{2w-1}{5} > 0 \quad \lambda_2^* = 0 \end{array}$$

The graph of x_1^* as a function of w is presented below (please, complete the picture)




The graph below shows constraint sets for different “important” values of w and some significant level curve of the objective function.


$w \in (0, \frac{1}{2})$:



$w = \frac{1}{2}$:



$w > \frac{1}{2}$:



Observe that in the example, we get that if $\lambda_2^* = 0$, the associated constraint $x_2 \geq 0$ is not significant. See Subsection 18.6.2, for a discussion of that statement.

Of course, several problems may arise in applying the above procedure. Below, we describe some commonly encountered problems and some possible (partial) solutions.

18.4.1 Some problems and some solutions

1. The set X .

X is not open.

Rewrite the problem in terms of an open set X' and some added constraints. A standard example is the following one.

$$\max_{x \in \mathbb{R}_+^n} f(x) \quad s.t. \quad g(x) \geq 0$$

which can be rewritten as

$$\max_{x \in \mathbb{R}^n} f(x) \quad s.t. \quad \begin{aligned} g(x) &\geq 0 \\ x &\geq 0 \end{aligned}$$

2. Existence.

a. The constraint set is not compact. If the constraint set is not compact, it is sometimes possible to find another maximization problem such that

- i. its constraint set is compact and nonempty, and
- ii. whose solution set is contained in the solution set of the problem we are analyzing.

A way to try to achieve both i. and ii. above is to “restrict the constraint set (to make it compact) without eliminating the solution of the original problem”. Sometimes, a problem with

the above properties is the following one.

$$\max_{x \in X} f(x) \quad s.t. \quad \begin{array}{ll} g(x) & \geq 0 \\ f(x) - f(\hat{x}) & \geq 0 \end{array} \quad (P1)$$

where \hat{x} is an element of X such that $g(\hat{x}) \geq 0$.

Observe that $(P - \text{cons})$ is an example of $(P1)$.

In fact, while, condition 1. above., i.e., the compactness of the constraint set of $(P1)$ depends upon the specific characteristics of X, f and g , condition ii. above is satisfied by problem $(P1)$, as shown in detail below.

Define

$$M := \arg \max (P) \quad M^1 := \arg \max (P1)$$

and V and V^1 the constraint sets of Problems (P) and $(P1)$, respectively. Observe that

$$V^1 \subseteq V \quad (18.31)$$

If V^1 is compact, then $M^1 \neq \emptyset$ and the only thing left to show is that $M^1 \subseteq M$, which is always insured as proved below.

Proposition 746 $M^1 \subseteq M$.

Proof. If $M^1 = \emptyset$, we are done.

Suppose that $M^1 \neq \emptyset$, and that the conclusion of the Proposition is false, i.e., there exists $x^1 \in M^1$ such that

a. $x^1 \in M^1$, and b. $x^1 \notin M$, or

a. $\forall x \in X$ such that $g(x) \geq 0$ and $f(x) \geq f(\hat{x})$, we have $f(x^1) \geq f(x)$;

and

b. either i. $x^1 \notin V$,

or ii. $\exists \tilde{x} \in X$ such that

$$g(\tilde{x}) \geq 0 \quad (18.32)$$

and

$$f(\tilde{x}) > f(x^1) \quad (18.33)$$

Let's show that i. and ii. cannot hold.

i.

It cannot hold simply because $V^1 \subseteq V$, from 18.31.

ii.

Since $x^1 \in V^1$,

$$f(x^1) \geq f(\hat{x}) \quad (18.34)$$

From (18.33) and (18.34), it follows that

$$f(\tilde{x}) > f(\hat{x}) \quad (18.35)$$

But (18.32), (18.35) and (18.33) contradict the definition of x^1 , i.e., a. above ■

b. Existence without the Extreme Value Theorem If you are not able to show existence, but

i. sufficient conditions to apply Kuhn-Tucker conditions hold, and

ii. you are able to find a solution to the Kuhn-Tucker conditions, then a solution exists.

18.5 The Implicit Function Theorem and Comparative Statics Analysis

The Implicit Function Theorem can be used to study how solutions ($x \in X \subseteq \mathbb{R}^n$) to maximization problems and, if needed, associated Lagrange or Kuhn-Tucker multipliers ($\lambda \in \mathbb{R}^m$) change when parameters ($\pi \in \Pi \subseteq \mathbb{R}^k$) change. That analysis can be done if the solutions to the maximization problem (and the multipliers) are solution to a system of equation of the form

$$F_1(x, \pi) = 0$$

with $(\# \text{ choice variables}) = (\# \text{ dimension of the codomain of } F_1)$, or

$$F_2(\xi, \pi) = 0$$

where $\xi := (x, \lambda)$, and $(\# \text{ choice variables and multipliers}) = (\# \text{ dimension of the codomain of } F_2)$,

To apply the Implicit Function Theorem, it must be the case that the following conditions do hold.

1. $(\# \text{ choice variables } x) = (\# \text{ dimension of the codomain of } F_1)$, or
 $(\# \text{ choice variables and multipliers}) = (\# \text{ dimension of the codomain of } F_2)$.
2. F_i has to be at least C^1 . That condition is insured if the above systems are obtained from maximization problems characterized by functions f, g which are at least C^2 : usually the above systems contain some form of first order conditions, which are written using first derivatives of f and g .
3. $F_1(x^*, \pi_0) = 0$ or $F_2(\xi^*, \pi_0) = 0$. The existence of a solution to the system is usually the result of the strategy to describe how to solve a maximization form - see above Section 18.4.
4. $\det [D_x F_1(x^*, \pi_0)]_{n \times n} \neq 0$ or $\det [D_\xi F_2(\xi^*, \pi_0)]_{(n+m) \times (n+m)} \neq 0$. That condition has to be verified directly on the problem.

If the above conditions are verified, the Implicit Function Theorem allow to conclude what follows (in reference to F_2).

There exist an open neighborhood $N(\xi^*) \subseteq X$ of ξ^* , an open neighborhood $N(\pi_0) \subseteq \Pi$ of π_0 and a unique C^1 function $g : N(\pi_0) \subseteq \Pi \subseteq \mathbb{R}^p \rightarrow N(\xi^*) \subseteq X \subseteq \mathbb{R}^n$ such that $\forall \pi \in N(\pi_0)$, $F(g(\pi), \pi) = 0$ and

$$Dg(\pi) = - \left[D_\xi F(\xi, \pi)|_{\xi=g(\pi)} \right]^{-1} \cdot \left[D_\pi F(\xi, \pi)|_{\xi=g(\pi)} \right]$$

Therefore, using the above expression, we may be able to say if the increase in any value of any parameter implies an increase in the value of any choice variable (or multiplier).

Three significant cases of application of the above procedure are presented below. We are going to consider C^2 functions defined on open subsets of Euclidean spaces.

18.5.1 Maximization problem without constraint

Assume that the problem to study is

$$\max_{x \in X} f(x, \pi)$$

and that

1. f is concave;
 2. There exists a solution x^* to the above problem associated with π_0 .
- Then, from Proposition 686, we know that x^* is a solution to

$$Df(x, \pi_0) = 0$$

Therefore, we can try to apply the Implicit Function Theorem to

$$F_1(x, \pi) = Df(x, \pi_0)$$

An example of application of the strategy illustrated above is presented in Section 19.3.

18.5.2 Maximization problem with equality constraints

Consider a maximization problem

$$\max_{x \in X} f(x, \pi) \quad s.t. \quad g(x, \pi) = 0,$$

Assume that necessary and sufficient conditions to apply Lagrange Theorem hold and that there exists a vector (x^*, λ^*) which is a solution (not necessarily unique) associated with the parameter π_0 . Therefore, we can try to apply the Implicit Function Theorem to

$$F_2(\xi, \pi) = \begin{pmatrix} Df(x^*, \pi_0) + \lambda^* Dg(x^*, \pi_0) \\ g(x^*, \pi_0) \end{pmatrix} \quad (18.36)$$

18.5.3 Maximization problem with Inequality Constraints

Consider the following maximization problems with inequality constraints. For given $\pi \in \Pi$,

$$\max_{x \in X} f(x, \pi) \quad s.t. \quad g(x, \pi) \geq 0 \quad (18.37)$$

Moreover, assume that the set of solutions of that problem is nonempty and characterized by the set of solutions of the associated Kuhn-Tucker system, i.e., using the notation of Subsection 18.1,

$$M = S \neq \emptyset.$$

We have seen that we can write Kuhn-Tucker conditions in one of the two following ways, beside some other ones,

$$\begin{cases} Df(x, \pi) + \lambda Dg(x, \pi) = 0 & (1) \\ \lambda \geq 0 & (2) \\ g(x, \pi) \geq 0 & (3) \\ \lambda g(x, \pi) = 0 & (4) \end{cases} \quad (18.38)$$

or

$$\begin{cases} Df(x, \pi) + \lambda Dg(x, \pi) = 0 & (1) \\ \min\{\lambda_j, g_j(x, \pi)\} = 0 & \text{for } j \in \{1, \dots, m\} \end{cases} \quad (18.39)$$

The Implicit Function Theorem *cannot* be applied to either system (18.38) or system (18.39): system (18.38) contains *inequalities*; system (18.39) involves functions which are *not differentiable*. We present below conditions under which the Implicit Function Theorem can be anyway applied to allow to make comparative statics analysis. Take a solution (x^*, λ^*, π_0) to the above system(s). Assume that

$$\text{for each } j, \text{ either } \lambda_j^* > 0 \text{ or } g_j(x^*, \pi_0) > 0.$$

In other words, there is no j such that $\lambda_j = g_j(x^*, \pi_0) = 0$. Consider a partition J^*, \hat{J} of $\{1, \dots, m\}$, and the resulting Kuhn-Tucker conditions.

$$\begin{cases} Df(x^*, \pi_0) + \lambda^* Dg(x^*, \pi_0) = 0 \\ \lambda_j^* > 0 & \text{for } j \in J^* \\ g_j(x^*, \pi_0) = 0 & \text{for } j \in J^* \\ \lambda_j^* = 0 & \text{for } j \in \hat{J} \\ g_j(x^*, \pi_0) > 0 & \text{for } j \in \hat{J} \end{cases} \quad (18.40)$$

Define

$$\begin{aligned} g^*(x^*, \pi_0) &:= (g_j(x^*, \pi_0))_{j \in J^*} \\ \hat{g}(x^*, \pi_0) &:= (g_j(x^*, \pi_0))_{j \in \hat{J}} \\ \lambda^{**} &:= (\lambda_j^*)_{j \in J^*} \\ \hat{\lambda}^* &:= (\lambda_j^*)_{j \in \hat{J}} \end{aligned}$$

Write the system of equations obtained from system (18.40) eliminating strict inequality constraints and substituting in the zero variables:

$$\begin{cases} Df(x^*, \pi_0) + \lambda^{**} Dg^*(x^*, \pi_0) = 0 \\ g^*(x^*, \pi_0) = 0 \end{cases} \quad (18.41)$$

Observe that the number of equations is equal to the number of “remaining” unknowns and they are

$$n + \#J^*$$

i.e., Condition 1 presented at the beginning of the present Section 18.5 is satisfied. Assume that the needed rank condition does hold and we therefore can apply the Implicit Function Theorem to

$$F_2(\xi, \pi) = \begin{pmatrix} Df(x^*, \pi_0) + \lambda^{**} Dg^*(x^*, \pi_0) \\ g^*(x^*, \pi_0) \end{pmatrix} = 0$$

Then, we can conclude that there exists a unique C^1 function φ defined in an open neighborhood N_1 of π_0 such that

$$\forall \pi \in N_1, \quad \varphi(\pi) := (x^*(\pi), \lambda^{**}(\pi))$$

is a solution to system (18.41) at π .

Therefore, by definition of φ ,

$$\begin{cases} Df(x^*(\pi), \pi) + \lambda^{**}(\pi)^T Dg^*(x^*(\pi), \pi) & = 0 \\ g^*(x^*(\pi), \pi) & = 0 \end{cases} \quad (18.42)$$

Since φ is continuous and $\lambda^{**}(\pi_0) > 0$ and $\hat{g}(x^*(\pi_0), \pi_0) > 0$, there exist an open neighborhood $N_2 \subseteq N_1$ of π_0 such that $\forall \pi \in N_2$

$$\begin{cases} \lambda^{**}(\pi) & > 0 \\ \hat{g}(x(\pi), \pi) & > 0 \end{cases} \quad (18.43)$$

Take also $\forall \pi \in N_2$

$$\begin{cases} \hat{\lambda}^*(\pi) & = 0 \end{cases} \quad (18.44)$$

Then, systems (18.42), (18.43) and (18.44) say that $\forall \pi \in N_2$, $(x(\pi), \lambda^*(\pi), \hat{\lambda}(\pi))$ satisfy Kuhn-Tucker conditions for problem (18.37) and therefore, since $C = M$, they are solutions to the maximization problem.

The above conclusion *does not hold true* if Kuhn-Tucker conditions are of the following form

$$\begin{cases} Df(x, \pi) + \lambda^T Dg(x, \pi) & = 0 \\ \lambda_j = 0, & g_j(x, \pi) = 0 & \text{for } j \in J' \\ \lambda_j > 0, & g_j(x, \pi) = 0 & \text{for } j \in J'' \\ \lambda_j = 0, & g_j(x, \pi) > 0 & \text{for } j \in \hat{J} \end{cases} \quad (18.45)$$

where $J' \neq \emptyset$, J'' and \hat{J} is a partition of J .

In that case, applying the same procedure described above, i.e., eliminating strict inequality constraints and substituting in the zero variables, leads to the following systems in the unknowns $x \in \mathbb{R}^n$ and $(\lambda_j)_{j \in J''} \in \mathbb{R}^{\#J''}$:

$$\begin{cases} Df(x, \pi) + (\lambda_j)_{j \in J''} D(g_j)_{j \in J''}(x, \pi) & = 0 \\ g_j(x, \pi) = 0 & \text{for } j \in J' \\ g_j(x, \pi) = 0 & \text{for } j \in J'' \end{cases}$$

and therefore the number of equation is $n + \#J'' + \#J' > n + \#J''$, simply because we are considering the case $J' \neq \emptyset$. Therefore the crucial condition

$$(\# \text{ choice variables and multipliers}) = (\# \text{ dimension of the codomain of } F_2)$$

is violated.

Even if the Implicit Function Theorem could be applied to the equations contained in (18.45), in an open neighborhood of π_0 we could have

$$\lambda_j(\pi) < 0 \text{ and/or } g_j(x(\pi), \pi) < 0 \text{ for } j \in J'$$

Then $\varphi(\pi)$ would be solutions to a set of equations and inequalities which are not Kuhn-Tucker conditions of the maximization problem under analysis, and therefore $x(\pi)$ would not be a solution to the that maximization problem.

An example of application of the strategy illustrated above is presented in Section 19.1.

18.6 The Envelope Theorem and the meaning of multipliers

18.6.1 The Envelope Theorem

Consider the problem (M) : for given $\pi \in \Pi$,

$$\max_{x \in X} f(x, \pi) \quad s.t. \quad g(x, \pi) = 0$$

Assume that for every π , the above problem admits a unique solution characterized by Lagrange conditions and that the Implicit function theorem can be applied. Then, there exists an open set $\mathcal{O} \subseteq \Pi$ such that

$$\begin{aligned} x : \mathcal{O} &\rightarrow X, \quad x : \pi \mapsto \arg \max (P) , \\ v : \mathcal{O} &\rightarrow \mathbb{R}, \quad v : \pi \mapsto \max (P) \\ \lambda : \mathcal{O} &\rightarrow \mathbb{R}^m, \quad \pi \mapsto \text{unique Lagrange multiplier vector} \end{aligned} \quad \text{and}$$

are differentiable functions.

Theorem 747 *For any $\pi^* \in \mathcal{O}$ and for any pair of associated $(x^*, \lambda^*) := (x(\pi^*), \lambda(\pi^*))$, we have*

$$D_\pi v(\pi^*) = D_\pi \mathcal{L}(x^*, \lambda^*, \pi^*)$$

i.e.,

$$D_\pi v(\pi^*) = D_\pi f(x^*, \pi^*) + \lambda^* D_\pi g(x^*, \pi^*)$$

Remark 748 *Observe that the above analysis applies also to the case of inequality constraints, as long as the set of binding constraints does not change.*

Proof. of Theorem 747 By definition of $v(\cdot)$ and $x(\cdot)$, we have that

$$\forall \pi \in \mathcal{O}, \quad v(\pi) = f(x(\pi), \pi). \quad (1)$$

Consider an arbitrary value π^* and the unique associate solution $x^* = x(\pi^*)$ of problem (P) . Differentiating both sides of (1) with respect to π and computing at π^* , we get

$$[D_\pi v(\pi^*)]_{1 \times k} = [D_x f(x, \pi)_{|(x^*, \pi^*)}]_{1 \times n} \cdot [D_\pi x(\pi)_{|\pi=\pi^*}]_{n \times k} + [D_\pi f(x, \pi)_{|(x^*, \pi^*)}]_{1 \times k} \quad (2)$$

From Lagrange conditions

$$D_x f(x, \pi)_{|(x^*, \pi^*)} = -\lambda^* D_x g(x, \pi)_{|(x^*, \pi^*)} \quad (3),$$

where λ^* is the unique value of the Lagrange multiplier. Moreover

$$\forall \pi \in \mathcal{O}, \quad g(x(\pi), \pi) = 0. \quad (4)$$

Differentiating both sides of (4) with respect to π and computing at π^* , we get

$$[D_x g(x, \pi)_{|(x^*, \pi^*)}]_{m \times n} \cdot [D_\pi x(\pi)_{|\pi=\pi^*}]_{n \times k} + [D_\pi g(x, \pi)_{|(x^*, \pi^*)}]_{m \times k} = 0 \quad (5).$$

Finally,

$$\begin{aligned} [D_\pi v(\pi^*)]_{1 \times k} &\stackrel{(2),(3)}{=} -\lambda^* D_x g(x, \pi)_{|(x^*, \pi^*)} D_\pi x(\pi)_{|\pi=\pi^*} + D_\pi f(x, \pi)_{|(x^*, \pi^*)} \stackrel{(5)}{=} \\ &= D_\pi f(x, \pi)_{|(x^*, \pi^*)} + \lambda^* D_\pi g(x, \pi)_{|(x^*, \pi^*)} \end{aligned}$$

■

18.6.2 On the meaning of the multipliers

The main goal of this subsection is to try to formalize the following statements.

1. The fact that $\lambda_j = 0$ indicates that the associated constraint $g_j(x) \geq 0$ is not significant - see Proposition 749 below.
2. The fact that $\lambda_j > 0$ indicates that a way to increase the value of the objective function is to violate the associated constraint $g_j(x) \geq 0$ - see Proposition 750 below.

For simplicity, consider the case $m = 1$. Let (CP) be the problem

$$\max_{x \in X} f(x) \quad \text{s.t.} \quad g(x) \geq 0$$

and (UP) the problem

$$\max_{x \in X} f(x)$$

with f strictly quasi-concave, g is quasi-concave and solutions to both problem exist. Define $x^* := \arg \max (CP)$ with associated multiplier λ^* , and $x^{**} := \arg \max (UP)$.

Proposition 749 *If $\lambda^* = 0$, then $x^* = \arg \max (UP) \Leftrightarrow x^* = \arg \max (CP)$.*

Proof. By the assumptions of this section, the solution to (CP) exists, is unique, it is equal to x^* and there exists λ^* such that

$$\begin{aligned} Df(x^*) + \lambda^* Dg(x^*) &= 0 \\ \min\{g(x^*), \lambda^*\} &= 0. \end{aligned}$$

Moreover, the solution to (UP) exists, is unique and it is the solution to

$$Df(x) = 0.$$

Since $\lambda^* = 0$, the desired result follows. ■

Take $\varepsilon > 0$ and $k \in (-\varepsilon, +\infty)$. Let (CPk) be the problem

$$\max_{x \in X} f(x) \quad \text{s.t.} \quad g(x) \geq k$$

Let

$$\hat{x} : (-\varepsilon, +\infty) \rightarrow X, \quad k \mapsto \arg \max (CPk)$$

$$\hat{v} : (-\varepsilon, +\infty) \rightarrow \mathbb{R}, \quad k \mapsto \max (CPk) := f(\hat{x}(k))$$

Let $\hat{\lambda}(k)$ be such that $(\hat{x}(k), \hat{\lambda}(k))$ is the solution to the associated Kuhn-Tucker conditions.

Observe that

$$x^* = \hat{x}(0), \quad \lambda^* = \hat{\lambda}(0) \tag{18.46}$$

Proposition 750 *If $\lambda^* > 0$, then $\hat{v}'(0) < 0$.*

Proof. From the envelope theorem,

$$\forall k \in (-\varepsilon, +\infty), \hat{v}'(k) = \frac{\partial (f(x) + \lambda(g(x) - k))}{\partial k} \Big|_{(\hat{x}(k), \hat{\lambda}(k))} = -\hat{\lambda}(k)$$

and from (18.46)

$$\hat{v}'(0) = -\hat{\lambda}(0) = -\lambda^* < 0.$$

■

Remark 751 *Consider the following problem. For given $a \in \mathbb{R}$,*

$$\max_{x \in X} f(x) \quad \text{s.t.} \quad g(x) - a \geq 0 \tag{18.47}$$

Assume that the above problem is “well-behaved” and that $x(a) = \arg \max (18.47)$, $v(a) = f(x(a))$ and $(x(a), \lambda(a))$ is the solution of the associated Kuhn-Tucker conditions. Then, applying the Envelope Theorem we have

$$v'(a) = \lambda(a)$$

Chapter 19

Applications to Economics

19.1 The Walrasian Consumer Problem

The utility function of a household is

$$u : \mathbb{R}_{++}^C \rightarrow \mathbb{R} \quad : x \mapsto u(x).$$

Assumption. u is a \mathcal{C}^2 function; u is differentially strictly increasing, i.e., $\forall x \in \mathbb{R}_{++}^C, Du(x) \gg 0$; u is differentially strictly quasi-concave, i.e., $\forall x \in \mathbb{R}_{++}^C, \Delta x \neq 0$ and $Du(x) \Delta x = 0 \Rightarrow \Delta x^T D^2 u(x) \Delta x < 0$; for any $\underline{u} \in \mathbb{R}$, $\{x \in \mathbb{R}_{++}^C : u(x) \geq \underline{u}\}$ is closed in \mathbb{R}^C .

The maximization problem for household h is

$$(P1) \quad \max_{x \in \mathbb{R}_{++}^C} u(x) \quad s.t. \quad px - w \leq 0.$$

The budget set of the above problem is clearly not compact. But, in the Appendix, we show that the solution of (P1) are the same as the solutions of (P2) and (P3) below. Observe that the constraint set of (P3) is compact.

$$(P2) \quad \max_{x \in \mathbb{R}_{++}^C} u(x) \quad s.t. \quad px - w = 0;$$

$$(P3) \quad \max_{x \in \mathbb{R}_{++}^C} u(x) \quad s.t. \quad \begin{aligned} px - w &\leq 0; \\ u(x) &\geq u(e^*), \end{aligned}$$

where $e^* \in \{x \in \mathbb{R}_{++}^C : px \leq w\}$.

Theorem 752 Under the Assumptions (smooth 1-5), $\xi_h(p, w_h)$ is a \mathcal{C}^1 function.

Proof.

Observe that, from it can be easily shown that, ξ is a function.

We want to show that (P2) satisfies necessary and sufficient conditions to Lagrange Theorem, and then apply the Implicit Function Theorem to the First Order Conditions of that problem.

The necessary condition is satisfied because $D_x [px - w] = p \neq 0$;

Define also

$$\mu : \mathbb{R}_{++}^C \times \mathbb{R}_{++}^{C-1} \rightarrow \mathbb{R}_{++}^C,$$

$$\mu : (p, w) \mapsto \text{Lagrange multiplier for (P2)}.$$

The sufficient conditions are satisfied because: from Assumptions (smooth 4), u is differentially strictly quasi-concave; the constraint is linear; the Lagrange multiplier μ is strictly positive -see below.

The Lagrangian function for problem (P2) and the associated First Order Conditions are described below.

$$\mathcal{L}(x, \mu, p, w) = u(x) + \mu \cdot (-px + w)$$

$$(FOC) \quad \begin{aligned} (1) \quad Du(x) - \mu p &= 0 \\ (2) \quad -px + w &= 0 \end{aligned}$$

Define

$$F : \mathbb{R}_{++}^C \times \mathbb{R}_{++} \times \mathbb{R}_{++}^C \times \mathbb{R}_{++} \rightarrow \mathbb{R}^C \times \mathbb{R},$$

$$F : (x, \mu, p, w) \mapsto \begin{pmatrix} Du(x) - \mu p \\ -px + w \end{pmatrix}.$$

As an application of the Implicit Function Theorem, it is enough to show that $D_{(x,\mu)} F(x, \mu, p, w)$ has full row rank $(C+1)$.

Suppose $D_{(x,\mu)} F$ does not have full rank; then there would exist

$\Delta x \in \mathbb{R}^C$ and $\Delta \mu \in \mathbb{R}$ such that $\Delta := (\Delta x, \Delta \mu) \neq 0$ and $D_{(x,\mu)} F \cdot (\Delta x, \Delta \mu) = 0$, or

$$\begin{bmatrix} D^2 u(x) & -p^T \\ -p & 0 \end{bmatrix} \begin{bmatrix} \Delta x \\ \Delta \mu \end{bmatrix} = 0,$$

or

$$\begin{aligned} (a) \quad D^2 u(x) \Delta x - p^T \Delta \mu &= 0 \\ (b) \quad -p \Delta x &= 0 \end{aligned}$$

The idea of the proof is to contradict Assumption u3.

Claim 1. $\Delta x \neq 0$.

By assumption it must be $\Delta \neq 0$ and therefore, if $\Delta x = 0$, $\Delta \mu \neq 0$. Since $p \in \mathbb{R}_{++}^C$, $p^T \Delta \mu \neq 0$. Moreover, if $\Delta x = 0$, from (a), we would have $p^T \Delta \mu = 0$, a contradiction.

Claim 2. $Du \cdot \Delta x = 0$.

From (FOC1), we have $Du \cdot \Delta x - \mu_h p \cdot \Delta x = 0$; using (b) the desired result follows.

Claim 3. $\Delta x^T D^2 u \cdot \Delta x = 0$.

Premultiplying (a) by Δx^T , we get $\Delta x^T D^2 u(x) \Delta x - \Delta x^T p^T \Delta \mu = 0$. Using (b), the result follows.

Claims 1, 2 and 3 contradict Assumption u3.

■

The above result gives also a way of computing $D_{(p,w)} x(p, w)$, as an application of the Implicit Function Theorem.

Since

$$\begin{array}{cccccc} & x & \mu & p & w & \\ & Du(x) - \mu p & D^2 u & -p^T & -\mu I_C & 0 \\ & -px + w & -p & 0 & -x & 1 \end{array}$$

$$[D_{(p,w)}(x, \mu)(p, w)]_{(C+1) \times (C+1)} = \begin{bmatrix} D_p x & D_w x \\ D_p \mu & D_w \mu \end{bmatrix} =$$

$$= - \begin{bmatrix} D^2 u & -p^T \\ -p & 0 \end{bmatrix}_{(C+1) \times (C+1)}^{-1} \begin{bmatrix} -\mu I_C & 0 \\ -x & 1 \end{bmatrix}_{(C+1) \times (C+1)}$$

To compute the inverse of the above matrix, we can use the following fact about the inverse of partitioned matrix (see for example, Goldberger, (1963), page 26)

Let A be an $n \times n$ nonsingular matrix partitioned as

$$A = \begin{bmatrix} E & F \\ G & H \end{bmatrix},$$

where $E_{n_1 \times n_1}$, $F_{n_1 \times n_2}$, $G_{n_2 \times n_1}$, $H_{n_2 \times n_2}$ and $n_1 + n_2 = n$. Suppose that E and $D := H - GE^{-1}F$ are non singular. Then

$$A^{-1} = \begin{bmatrix} E^{-1}(I + FD^{-1}GE^{-1}) & -E^{-1}FD^{-1} \\ -D^{-1}GE^{-1} & D^{-1} \end{bmatrix}.$$

If we assume that $D^2 u$ is negative definite and therefore invertible, we have

$$\begin{bmatrix} D^2 u & -p^T \\ -p & 0 \end{bmatrix}^{-1} = \begin{bmatrix} (D^2)^{-1} (I + \delta^{-1} p^T p (D^2)^{-1}) & \delta^{-1} (D^2)^{-1} p^T \\ \delta^{-1} p (D^2)^{-1} & \delta^{-1} \end{bmatrix}$$

where $\delta = -p (D^2)^{-1} p^T \in \mathbb{R}_{++}$.

And

$$\begin{aligned}
[D_p x(p, w)]_{C \times C} &= - \begin{bmatrix} (D^2)^{-1} (I + \delta^{-1} p^T p (D^2)^{-1}) & \delta^{-1} (D^2)^{-1} p^T \end{bmatrix} \begin{bmatrix} \mu I_C \\ x \end{bmatrix} = \\
&= -\mu (D^2)^{-1} (I + \delta^{-1} p^T p (D^2)^{-1}) - \delta^{-1} (D^2)^{-1} p^T x = -\delta^{-1} (D^2)^{-1} \left[\mu (\delta I + p^T p (D^2)^{-1}) + p^T x \right] \\
[D_w x(p, w)]_{C \times 1} &= - \begin{bmatrix} (D^2)^{-1} (I + \delta^{-1} p^T p (D^2)^{-1}) & \delta^{-1} (D^2)^{-1} p^T \end{bmatrix} \begin{bmatrix} 0 \\ -1 \end{bmatrix} = \delta^{-1} (D^2)^{-1} p^T \\
[D_p \mu(p, w)]_{1 \times C} &= - \begin{bmatrix} \delta^{-1} p (D^2)^{-1} & \delta^{-1} \end{bmatrix} \begin{bmatrix} \mu I_C \\ x \end{bmatrix} = -\delta^{-1} (\mu p (D^2)^{-1} + x). \\
[D_w \mu(p, w)]_{1 \times 1} &= - \begin{bmatrix} \delta^{-1} p (D^2)^{-1} & \delta^{-1} \end{bmatrix} \begin{bmatrix} 0 \\ -1 \end{bmatrix} = \delta^{-1}.
\end{aligned}$$

As a simple application of the Envelope Theorem, we also have that, defined the indirect utility function as

$$v : \mathbb{R}_{++}^{C+1} \rightarrow \mathbb{R}, \quad v : (p, w) \mapsto u(x(p, w)),$$

we have that

$$D_{(p,w)} v(p, w) = \lambda \begin{bmatrix} -x^T & 1 \end{bmatrix}.$$

19.2 Production

Definition 753 A production vector (or input-output or netput vector) is a vector $y := (y^c)_{c=1}^C \in \mathbb{R}^C$ which describes the net outputs of C commodities from a production process. Positive numbers denote outputs, negative numbers denote inputs, zero numbers denote commodities neither used nor produced.

Observe that, given the above definition, py is the profit of the firm.

Definition 754 The set of all feasible production vectors is called the production set $Y \subseteq \mathbb{R}^C$. If $y \in Y$, then y can be obtained as a result of the production process; if $y \notin Y$, that is not the case.

Definition 755 The Profit Maximization Problem (PMP) is

$$\max_y py \quad \text{s.t.} \quad y \in Y.$$

It is convenient to describe the production set Y using a function $F : \mathbb{R}^C \rightarrow \mathbb{R}$ called the transformation function. That is done as follows:

$$Y = \{y \in \mathbb{R}^C : F(y) \geq 0\}.$$

We list below a smooth version of the assumptions made on Y , using the transformation function.

Some assumption on $F(\cdot)$.

- (1) $\exists y \in \mathbb{R}^C$ such that $F(y) \geq 0$.
- (2) F is C^2 .
- (3) (No Free Lunch) If $y \geq 0$, then $F(y) < 0$.
- (4) (Possibility of Inaction) $F(0) = 0$.
- (5) (F is differentially strictly decreasing) $\forall y \in \mathbb{R}^C, DF(y) \ll 0$
- (6) (Irreversibility) If $y \neq 0$ and $F(y) \geq 0$, then $F(-y) < 0$.
- (7) (F is differentially strictly concave) $\forall \Delta \in \mathbb{R}^C \setminus \{0\}, \Delta^T D^2 F(y) \Delta < 0$.

Definition 756 Consider a function $F(\cdot)$ satisfying the above properties and a strictly positive real number N . The Smooth Profit Maximization Problem (SPMP) is

$$\max_y py \quad \text{s.t.} \quad F(y) \geq 0 \text{ and } \|y\| \leq N. \quad (19.1)$$

Remark 757 For any solution to the above problem it must be the case that $F(y) = 0$. Suppose there exists a solution y' to (SPMP) such that $F(y') > 0$. Since F is continuous, in fact C^2 , there exists $\varepsilon > 0$ such that $z \in B(y', \varepsilon) \Rightarrow F(z) > 0$. Take $z' = y' + \frac{\varepsilon \cdot 1}{C}$. Then, $d(y', z') := \left(\sum_{c=1}^C \left(\frac{\varepsilon}{C} \right)^2 \right)^{\frac{1}{2}} = \left(C \left(\frac{\varepsilon}{C} \right)^2 \right)^{\frac{1}{2}} = \left(\frac{\varepsilon^2}{C} \right)^{\frac{1}{2}} = \frac{\varepsilon}{\sqrt{C}} < \varepsilon$. Therefore $z' \in B(y', \varepsilon)$ and

$$F(z') > 0 \quad (1).$$

But,

$$pz' = py' + p \frac{\varepsilon \cdot 1}{C} > py' \quad (2).$$

(1) and (2) contradict the fact that y' solves (SPMP).

Proposition 758 If a solution with $\|y\| < N$ to (SPMP) exists. Then $y : \mathbb{R}_{++}^C \rightarrow \mathbb{R}^C$, $p \mapsto \arg \max (19.1)$ is a well defined C^1 function.

Proof.

Let's first show that $y(p)$ is single valued.

Suppose there exist $y, y' \in y(p)$ with $y \neq y'$. Consider $y^\lambda := (1 - \lambda)y + \lambda y'$. Since $F(\cdot)$ is strictly concave, it follows that $F(y^\lambda) > (1 - \lambda)F(y) + \lambda F(y') \geq 0$, where the last inequality comes from the fact that $y, y' \in y(p)$. But then $F(y^\lambda) > 0$. Then following the same argument as in Remark 757, there exists $\varepsilon > 0$ such that $z' = y^\lambda + \frac{\varepsilon \cdot 1}{C}$ and $F(z') > 0$. But $pz' > py^\lambda = (1 - \lambda)py + \lambda py' = py$, contradicting the fact that $y \in y(p)$.

Let's now show that y is C^1 .

From Remark 757 and from the assumption that $\|y\| < N$, (SPMP) can be rewritten as $\max_y py$ s.t. $F(y) = 0$. We can then try to apply Lagrange Theorem.

Necessary conditions: $DF(y) \ll 0$;

sufficient conditions: py is linear and therefore pseudo-concave; $F(\cdot)$ is differentially strictly concave and therefore quasi-concave; the Lagrange multiplier λ is strictly positive -see below.

Therefore, the solutions to (SPMP) are characterized by the following First Order Conditions, i.e., the derivative of the Lagrangian function with respect to y and λ equated to zero:

$$\mathcal{L}(y, p) = py + \lambda F(y). \quad \begin{matrix} y \\ p + \lambda DF(y) = 0 \end{matrix} \quad \begin{matrix} p \\ F(y) = 0 \end{matrix}$$

Observe that $\lambda = -\frac{p \cdot 1}{D_{y^1} F(y)} > 0$.

As usual to show differentiability of the choice function we take derivatives of the First Order Conditions.

$$\begin{matrix} y & \lambda \\ p + \lambda DF(y) = 0 & D^2 F(y) \quad [DF(y)]^T \\ F(y) = 0 & DF(y) \quad 0 \end{matrix}^T$$

We want to show that the above matrix has full rank. By contradiction, assume that there exists $\Delta := (\Delta y, \Delta \lambda) \in \mathbb{R}^C \times \mathbb{R}$, $\Delta \neq 0$ such that

$$\begin{bmatrix} D^2 F(y) & [DF(y)]^T \\ DF(y) & 0 \end{bmatrix} \begin{bmatrix} \Delta y \\ \Delta \lambda \end{bmatrix} = 0,$$

i.e.,

$$D^2 F(y) \cdot \Delta y + [DF(y)]^T \cdot \Delta \lambda = 0 \quad (a),$$

$$DF(y) \cdot \Delta y = 0 \quad (b).$$

Premultiplying (a) by Δy^T , we get $\Delta y^T \cdot D^2 F(y) \cdot \Delta y + \Delta y^T \cdot [DF(y)]^T \cdot \Delta \lambda = 0$. From (b), it follows that $\Delta y^T \cdot D^2 F(y) \cdot \Delta y = 0$, contradicting the differentially strict concavity of $F(\cdot)$.

(3)

From the Envelope Theorem, we know that if $(\bar{y}, \bar{\lambda})$ is the unique pair of solution-multiplier associated with \bar{p} , we have that

$$D_p \pi(p)|_{\bar{p}} = D_p(py)|_{(\bar{p}, \bar{y})} + \bar{\lambda} D_p F(y)|_{(\bar{p}, \bar{y})}.$$

Since $D_p(py)|_{(\bar{p}, \bar{y})} = \bar{y}$, $D_p F(y) = 0$ and, by definition of \bar{y} , $\bar{y} = y(\bar{p})$, we get $D_p \pi(p)|_{\bar{p}} = y(\bar{p})$, as desired.

(4)

From (3), we have that $D_p y(p) = D_p^2 \pi(p)$. Since $\pi(\cdot)$ is convex -see Proposition ?? (2)- the result follows.

(5)

From Proposition ?? (4) and the fact that $y(p)$ is single valued, we know that $\forall \alpha \in \mathbb{R}_{++}$, $y(p) - y(\alpha p) = 0$. Taking derivatives with respect to α , we have $D_p y(p)|_{(\alpha p)} \cdot p = 0$. For $\alpha = 1$, the desired result follows.

■

19.3 The demand for insurance

Consider an individual whose wealth is

$$\begin{array}{lll} W - d & \text{with probability} & \pi, \text{ and} \\ W & \text{with probability} & 1 - \pi, \end{array}$$

where $W > 0$ and $d > 0$.

Let the function

$$u : A \rightarrow \mathbb{R}, \quad u : c \rightarrow u(c)$$

be the individual's Bernoulli function.

Assumption 1. $\forall c \in \mathbb{R}$, $u'(c) > 0$ and $u''(c) < 0$.

Assumption 2. u is bounded above.

An insurance company offers a contract with following features: the potentially insured individual pays a premium p in each state and receives d if the accident occurs. The (potentially insured) individual can buy a quantity $a \in \mathbb{R}$ of the contract. In the case, she pays a premium $(a \cdot p)$ in each state and receives a reimbursement $(a \cdot d)$ if the accident occurs. Therefore, if the individual buys a quantity a of the contract, she get a wealth described as follows

$$\begin{array}{lll} W_1 := W - d - ap + ad & \text{with probability} & \pi, \text{ and} \\ W_2 := W - ap & \text{with probability} & 1 - \pi. \end{array} \quad (19.2)$$

Remark 759 *It is reasonable to assume that $p \in (0, d)$.*

Define

$$U : \mathbb{R} \rightarrow \mathbb{R}, \quad U : a \mapsto \pi u(W - d - ap + ad) + (1 - \pi) u(W - ap).$$

Then the individual solves the following problem. For given, $W \in \mathbb{R}_{++}$, $d \in \mathbb{R}_{++}$, $p \in (0, d)$, $\pi \in (0, 1)$

$$\max_{a \in \mathbb{R}} U(a) \quad (M) \quad (19.3)$$

To show existence of a solution, we introduce the problem presented below. For given $W \in \mathbb{R}_{++}$, $d \in \mathbb{R}_{++}$, $p \in (0, d)$, $\pi \in (0, 1)$

$$\max_{a \in \mathbb{R}} U(a) \quad \text{s.t.} \quad U(a) \geq U(0) \quad (M')$$

Defined $A^* := \arg \max(M)$ and $A' := \arg \max(M')$, the existence of solution to (M) , follows from the Proposition below.

Proposition 760 1. $A' \subset A^*$. 2. $A' \neq \emptyset$.

Proof.

Exercise

■

To show that the solution is unique, observe that

$$U'(a) = \pi u'(W - d + a(d - p))(d - p) + (1 - \pi) u'(W - ap)(-p) \quad (19.4)$$

and therefore

$$U''(a) = \overset{(+)}{\pi} u''(W - d + a(d - p)) \overset{(-)}{(d - p)} \overset{(+)}{(d - p)^2} + (1 - \pi) \overset{(+)}{u''(W - ap)} \overset{(-)}{p} \overset{(+)}{p^2} < 0.$$

Summarizing, the unique solution of problem (M) is the unique solution of the equation:

$$U'(a) = 0.$$

Definition 761 $a^* : \mathbb{R}_{++} \times (0, 1) \times (0, d) \times \mathbb{R}_{++} \rightarrow \mathbb{R}$, $a^* : (d, \pi, p, W) \mapsto \arg \max (M).$ $U^* : \Theta \rightarrow \mathbb{R}$, $U^* : \theta \mapsto \pi u(W - d + a^*(\theta)(d - p)) + (1 - \pi) u(W - a^*(\theta)p)$ **Proposition 762** *The signs of the derivatives of a^* and U^* with respect to θ are presented in the following table¹:*

		d	π	p	W
a^*		> 0 if $a^* \in [0, 1]$	> 0	≥ 0	≤ 0 if $a^* \leq 1$
U^*		≤ 0 if $a^* \in [0, 1]$	≤ 0 if $a^* \in [0, 1]$	≤ 0 if $a^* \geq 0$	> 0

Proof. Exercise. ■

19.4 Exercises on part IV

See Tito Pietra's file (available on line): Exercises 15.1 \rightarrow 15.6.

¹ Conditions on $a^*(\theta)$ contained in the table can be expressed in terms of exogenous variables.

Part V

Problem Sets

Chapter 20

Exercises

20.1 Linear Algebra

1.

Show that the set of pair of real numbers is **not** a vector space with respect to the following operations:

- (i). $(a, b) + (c, d) = (a + c, b + d)$ and $k(a, b) = (ka, b)$;
- (ii) $(a, b) + (c, d) = (a + c, b)$ and $k(a, b) = (ka, kb)$.

2.

Show that W is *not* a vector subspace of \mathbb{R}^3 on \mathbb{R} if

- (i) $W = \{(x, y, z) \in \mathbb{R}^3 : z \geq 0\}$;
- (ii) $W = \{x \in \mathbb{R}^3 : \|x\| \leq 1\}$,
- (iii) $W = \mathbb{Q}^3$.

3.

Let V be the vector space of all functions $f : \mathbb{R} \rightarrow \mathbb{R}$. Show the W is a vector subspace of V if

- (i) $W = \{f \in V : f(1) = 0\}$;
- (ii) $W = \{f \in V : f(1) = f(2)\}$.

4.

Show that

- (i).

$$V = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1 + x_2 + x_3 = 0\}$$

is a vector subspace of \mathbb{R}^3 ;

- (ii).

$$S = \{(1, -1, 0), (0, 1, -1)\}$$

is a basis for V .

5.

Show the following fact.

Proposition. Let a matrix $A \in \mathbb{M}(n, n)$, with $n \in \mathbb{N}$ be given. The set

$$\mathcal{C}_A := \{B \in \mathbb{M}(n, n) : BA = AB\}$$

is a vector subspace of $\mathbb{M}(n, n)$ (with respect to the field \mathbb{R}).

6.

Let U and V be vector subspaces of a vector space W . Show that

$$U + V := \{w \in W : \exists u \in U \text{ and } v \in V \text{ such that } w = u + v\}$$

is a vector subspace of W .

7.

Show that the following set of vectors is linearly independent:

$$\{(1, 1, 1), (0, 1, 1), (0, 0, 1)\}.$$

8. Using the definition, find the change-of-basis matrix from

$$S = \{u_1 = (1, 2), u_2 = (3, 5)\}$$

to

$$E = \{e_1 = (1, 0), e_2 = (0, 1)\}$$

and from E to S . Check the conclusion of Proposition ??, i.e., that one matrix is the inverse of the other one.

9. Find the determinant of

$$C = \begin{bmatrix} 6 & 2 & 1 & 0 & 5 \\ 2 & 1 & 1 & -2 & 1 \\ 1 & 1 & 2 & -2 & 3 \\ 3 & 0 & 2 & 3 & -1 \\ -1 & -1 & -3 & 4 & 2 \end{bmatrix}$$

10. Say for which values of $k \in \mathbb{R}$ the following matrix has rank a. 4, b. 3:

$$A := \begin{bmatrix} k+1 & 1 & -k & 2 \\ -k & 1 & 2-k & k \\ 1 & 0 & 1 & -1 \end{bmatrix}$$

11.

Show that

$$V = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1 - x_2 = 0\}$$

is a vector subspace of \mathbb{R}^3 and find a basis for V .

12.

Given

$$l : \mathbb{R}^4 \rightarrow \mathbb{R}^4, \quad l(x_1, x_2, x_3, x_4) = (x_1, x_1 + x_2, x_1 + x_2 + x_3, x_1 + x_2 + x_3 + x_4)$$

show it is linear, compute the associated matrix with respect to canonical bases, and compute $\ker l$ and $\operatorname{Im} l$.

13.

Complete the text below.

Proposition. Assume that $l \in L(V, U)$ and $\ker l = \{0\}$. Then,

$$\forall u \in \operatorname{Im} l, \quad \text{there exists a unique } v \in V \text{ such that } l(v) = u.$$

Proof.

Since, by definition, there exists $v \in V$ such that

$$l(v) = u. \tag{20.1}$$

Take $v' \in V$ such that $l(v') = u$. We want to show that

$$\dots \tag{20.2}$$

Observe that

$$l(v) - l(v') \stackrel{(a)}{=} \dots\dots\dots (20.3)$$

where (a) follows from

Moreover,

$$l(v) - l(v') \stackrel{(b)}{=} \dots\dots\dots, (20.4)$$

where (b) follows from

Therefore,

$$l(v - v') = 0,$$

and, by definition of $\ker l$,

$$\dots\dots\dots (20.5)$$

Since,, from (20.5), it follows that

$$v - v' = 0.$$

14.

Let the following sets be given:

$$V = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 : x_1 - x_2 + x_3 - x_4 = 0\}$$

and

$$W = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 : x_1 + x_2 + x_3 + x_4 = 0\}$$

If possible, find a basis of $V \cap W$.

15.

Say if the following statement is true or false.

Let V and U be vector spaces on \mathbb{R} , W a vector subspace of U and $l \in \mathcal{L}(V, U)$. Then $l^{-1}(W)$ is a vector subspace of V .

16.

Let the following full rank matrices

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}$$

be given. Say for which values of $k \in \mathbb{R}$, the following linear system has solutions.

$$\begin{bmatrix} 1 & a_{11} & a_{12} & 0 & 0 & 0 \\ 2 & a_{21} & a_{22} & 0 & 0 & 0 \\ 3 & 5 & 6 & b_{11} & b_{12} & 0 \\ 4 & 7 & 8 & b_{21} & b_{22} & 0 \\ 1 & a_{11} & a_{12} & 0 & 0 & k \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} k \\ 1 \\ 2 \\ 3 \\ k \end{bmatrix}$$

17.

Consider the following Proposition contained in Section 8.1 in the class Notes:

Proposition $\forall v \in V$,

$$[l]_{\mathbf{v}}^{\mathbf{u}} \cdot [v]_{\mathbf{v}} = [l(v)]_{\mathbf{u}} \quad (20.6)$$

Verify the above equality in the case in which

a.

$$l : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad (x_1, x_2) \mapsto \begin{pmatrix} x_1 + x_2 \\ x_1 - x_2 \end{pmatrix}$$

b. the basis \mathbf{v} of the domain of l is

$$\left\{ \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right\},$$

c. the basis \mathbf{u} of the codomain of l is

$$\left\{ \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right\},$$

d.

$$v = \begin{pmatrix} 3 \\ 4 \end{pmatrix}.$$

18.

Complete the following proof.

Proposition. Let

$n, m \in \mathbb{N}$ such that $m > n$, and

a vector subspace L of \mathbb{R}^m such that $\dim L = n$

be given. Then, there exists $l \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ such that $\text{Im } l = L$.

Proof. Let $\{v^i\}_{i=1}^n$ be a basis of $L \subseteq \mathbb{R}^m$. Take $l \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ such that

$$\forall i \in \{1, \dots, n\}, \quad l_2(e_n^i) = v^i,$$

where e_n^i is the i -th element in the canonical basis in \mathbb{R}^n . Such function does exists and, in fact, it is unique as a consequence of a Proposition in the Class Notes that we copy below:

.....

Then, from the Dimension theorem

$$\dim \text{Im } l = \dots\dots\dots$$

Moreover,

$$L = \dots\dots\dots \{v^i\}_{i=1}^n \subseteq \dots\dots\dots$$

Summarizing,

$$L \subseteq \text{Im } l, \dim L = n \text{ and } \dim \text{Im } l \leq n,$$

and therefore

$$\dim \text{Im } l = n.$$

Finally, from Propositionin the class Notes since $L \subseteq \text{Im } l$, $\dim L = n$ and $\dim \text{Im } l = n$, we have that $\text{Im } l = L$, as desired.

Proposition in the class Notes says what follows:

.....

19.

Say for which value of the parameter $a \in \mathbb{R}$ the following system has one, infinite or no solutions

$$\begin{cases} ax_1 + x_2 = 1 \\ x_1 + x_2 = a \\ 2x_1 + x_2 = 3a \\ 3x_1 + 2x_2 = a \end{cases}$$

20.

Say for which values of k , the system below admits one, none or infinite solutions.

$$A(k) \cdot x = b(k)$$

where $k \in \mathbb{R}$, and

$$A(k) \equiv \begin{bmatrix} 1 & 0 \\ 1-k & 2-k \\ 1 & k \\ 1 & k-1 \end{bmatrix}, \quad b(k) \equiv \begin{bmatrix} k-1 \\ k \\ 1 \\ 0 \end{bmatrix}.$$

21.

Let $\mathcal{V} = \{v^1, v^2, \dots, v^n\}$ be a set of vectors in \mathbb{R}^n such that for any $i, j \in \{1, \dots, n\}$,

$$v^i \cdot v^j = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j. \end{cases} \quad (20.7)$$

Show that \mathcal{V} is a basis of \mathbb{R}^n .

22.

Given a vector space V on a field F , sets $A, B \subseteq V$ and $k \in F$, we define

$$A + B := \{v \in V : \text{there exist } a \in A \text{ and } b \in B \text{ such that } v = a + b\},$$

$$kA := \{v \in V : \text{there exist } a \in A \text{ such that } v = ka\}.$$

Given a vector space V on a field F , a linear function $T \in \mathcal{L}(V, V)$ and W vector subspace of V , W is said to be T -invariant if

$$T(W) \subseteq W.$$

Let W be both S -invariant and T -invariant and let $k \in F$. Show that

- W is $S + T$ -invariant;
- W is $S \circ T$ -invariant;
- W is kT -invariant.

23.

Show that the set of all 2×2 symmetric real matrices is a vector subspace of $\mathbb{M}(2, 2)$ and compute its dimension.

24.

Let V be a vector space on a field F and W a vector subspace of V . Show that

- $W + W = W$, and
- for any $\alpha \in F \setminus \{0\}$, $\alpha W = W$.

25.

Let $\mathcal{P}_n(\mathbb{R})$ be the set polynomials of degree smaller or equal than $n \in \mathbb{N}_+$ on the set of real numbers, i.e.,

$$\mathcal{P}_n(\mathbb{R}) = \left\{ f : \mathbb{R} \rightarrow \mathbb{R} \text{ such that } \exists a_0, a_1, \dots, a_n \in \mathbb{R} \text{ such that for any } t \in \mathbb{R}, f(t) = \sum_{i=0}^n a_i t^i \right\}.$$

Show that $\mathcal{P}_n(\mathbb{R})$ is isomorphic to \mathbb{R}^{n+1} .

20.2 Some topology in metric spaces

20.2.1 Basic topology in metric spaces

1.

Do Exercise 363: Let d be a metric on a non-empty set X . Show that

$$d' : X \times X \rightarrow \mathbb{R}, d'(x, y) = \frac{d(x, y)}{1 + d(x, y)}$$

is a metric on X .

2.

Let X be the set of continuous real valued functions with domain $[0, 1] \subseteq \mathbb{R}$ and

$$d(f, g) = \int_0^1 f(x) - g(x) dx,$$

where the integral is the Riemann Integral (that one you learned in Calculus 1). Show that (X, d) is not a metric space.

3.

Do Exercise 380 for $n = 2$: $\forall n \in \mathbb{N}, \forall i \in \{1, \dots, n\}, \forall a_i, b_i \in \mathbb{R}$ with $a_i < b_i$,

$$\times_{i=1}^n (a_i, b_i)$$

is (\mathbb{R}^n, d_2) open.

4.

Show the second equality in Remark 388:

$$\cap_{n=1}^{+\infty} \left(-\frac{1}{n}, \frac{1}{n} \right) = \{0\}$$

5.

Say if the following set is (\mathbb{R}, d_2) open or closed:

$$S := \left\{ x \in \mathbb{R} : \exists n \in \mathbb{N} \text{ such that } x = (-1)^n \frac{1}{n} \right\}$$

6.

Say if the following set is (\mathbb{R}, d_2) open or closed:

$$A := \cup_{n=1}^{+\infty} \left(\frac{1}{n}, 10 - \frac{1}{n} \right).$$

7.

Do Exercise 398: show that $\mathcal{F}(S) = \mathcal{F}(S^C)$.

8.

Do Exercise 399: show that $\mathcal{F}(S)$ is a closed set.

9.

Let the metric space (\mathbb{R}, d_2) be given. Find $\text{Int } S, \text{Cl } (S), \mathcal{F}(S), D(S), \text{Is } (S)$ and say if S is open or closed for $S = \mathbb{Q}$, $S = (0, 1)$ and $S = \{x \in \mathbb{R} : \exists n \in \mathbb{N}_+ \text{ such that } x = \frac{1}{n}\}$.

10.

Show that the following statements are **false**:

- $\text{Cl } (\text{Int } S) = S$,
- $\text{Int } \text{Cl } (S) = S$.

11.

Given $S \subseteq \mathbb{R}$, say if the following statements are true or false.

- a. S is an open bounded interval $\Rightarrow S$ is an open set;
- b. S is an open set $\Rightarrow S$ is an open bounded interval;
- c. $x \in \mathcal{F}(S) \Rightarrow x \in D(S)$;
- d. $x \in D(S) \Rightarrow x \in \mathcal{F}(S)$.

12.

Using the definition of convergent sequences, show that the following sequences do converge:

- a. $(x_n)_{n \in \mathbb{N}} \in \mathbb{R}^\infty$ such that $\forall n \in \mathbb{N}, x_n = 1$;
- b. $(x_n)_{n \in \mathbb{N}} \in \mathbb{R}^\infty$ such that $\forall n \in \mathbb{N}, x_n = \frac{1}{n}$.

13.

Using Proposition 425, show that $[0, 1]$ is (\mathbb{R}, d_2) closed.

14.

Show the following result: A subset of a discrete space, i.e., a metric space with the discrete metric, is compact if and only if it is finite.

15.

Say if the following statement is true: An open set is not compact.

16.

Using the definition of compactness, show the following statement: Any open ball in (\mathbb{R}^2, d_2) is not compact.

17.

Show that $f(A \cup B) = f(A) \cup f(B)$.

18.

Show that $f(A \cap B) \neq f(A) \cap f(B)$.

19.

Using the characterization of continuous functions in terms of open sets, show that for any metric space (X, d) the constant function is continuous.

20.

- a. Say if the following sets are (\mathbb{R}^n, d_2) compact:

- i.

$$\mathbb{R}_+^n,$$

- ii.

$$\forall x \in \mathbb{R}^n \text{ and } \forall r \in \mathbb{R}_{++}, \text{ } Cl B(x, r).$$

- b. Say if the following set is (\mathbb{R}, d_2) compact:

$$\left\{ x \in \mathbb{R} : \exists n \in \mathbb{N} \text{ such that } x = \frac{1}{n} \right\}.$$

21.

Given the continuous functions

$$g : \mathbb{R}^n \rightarrow \mathbb{R}^m$$

show that the following set is closed

$$\{x \in \mathbb{R}^n : g(x) \geq 0\}$$

22.

Assume that $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is continuous. Say if

$$X = \{x \in \mathbb{R}^m : f(x) = 0\}$$

is (a) closed, (b) is compact.

23.

Using the characterization of continuous functions in terms of open sets, show that the following function is not continuous

$$f : \mathbb{R} \rightarrow \mathbb{R}, \quad f(x) = \begin{cases} x & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$

24.

Using the Extreme Value Theorem, say if the following maximization problems have solutions (with $\| \cdot \|$ being the Euclidean norm).

$$\begin{aligned} \max_{x \in \mathbb{R}^n} \sum_{i=1}^n x_i \quad & \text{s.t.} \quad \|x\| \leq 1 \\ \max_{x \in \mathbb{R}^n} \sum_{i=1}^n x_i \quad & \text{s.t.} \quad \|x\| < 1 \\ \max_{x \in \mathbb{R}^n} \sum_{i=1}^n x_i \quad & \text{s.t.} \quad \|x\| \geq 1 \end{aligned}$$

25.

Let $(E, \|\cdot\|_E)$, $(F, \|\cdot\|_F)$ be normed vector spaces. A function $f : (E, \|\cdot\|_E) \rightarrow (F, \|\cdot\|_F)$ is bounded if

$$\exists M \in \mathbb{R}_{++} \text{ such that } \forall x \in E \quad \|f(x)\|_F \leq M.$$

Show that given a linear function $l : E \rightarrow F$,

$$l \text{ is bounded} \Leftrightarrow l = 0.$$

26.

$f : (X, d) \rightarrow \mathbb{R}$ is upper semicontinuous at $x_0 \in X$ if

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } d(x - x_0) < \delta \Rightarrow f(x) < f(x_0) + \varepsilon.$$

f is upper semicontinuous if is upper semicontinuous at any $x_0 \in X$.

Show that the following statements are equivalent:

- f is upper semicontinuous;
- for any $\alpha \in \mathbb{R}$, $\{x \in X : f(x) < \alpha\}$ is (X, d) open;
- for any $\alpha \in \mathbb{R}$, $\{x \in X : f(x) \geq \alpha\}$ is (X, d) closed.

27.

Let A be a subset of (\mathbb{R}^n, d) where d is the Euclidean distance. Show that if A is (\mathbb{R}^n, d) open, then for any $x \in X$, $\{x\} + A$ is (\mathbb{R}^n, d) open.

Hint: use the fact that for any $x, y \in \mathbb{R}^n$, $d(x, y) = \|x - y\|$ and therefore, for any $a \in \mathbb{R}^n$, $d(a + x, a + y) = \|a + x - a - y\| = \|x - y\| = d(x, y)$.

28.

Let (X, d) be a metric space. Show that if K_1 and K_2 are compact subsets of X , then $K_1 + K_2$ is compact.

29.

Given two metric spaces (E, d_1) and (F, d_2) , a function $f : E \rightarrow F$ is an isometry with respect to d_1 and d_2 if $\forall x_1, x_2 \in E$,

$$d_2(f(x_1), f(x_2)) = d_1(x_1, x_2).$$

Show that if $f : E \rightarrow F$ is an isometry then

- f is one-to-one;
- $\hat{f} : E \rightarrow f(E)$ is invertible;
- f is continuous.

20.2.2 Correspondences

To solve the following exercises on correspondences, we need some preliminary definitions.¹

A set $C \subseteq \mathbb{R}^n$ is convex if $\forall x_1, x_2 \in C$ and $\forall \lambda \in [0, 1]$, $(1 - \lambda)x_1 + \lambda x_2 \in C$.

A set $C \subseteq \mathbb{R}^n$ is strictly convex if $\forall x_1, x_2 \in C$ and $\forall \lambda \in (0, 1)$, $(1 - \lambda)x_1 + \lambda x_2 \in \text{Int } C$.

Consider an open and convex set $X \subseteq \mathbb{R}^n$ and a continuous function $f : X \rightarrow \mathbb{R}$, f is quasi-concave iff $\forall x', x'' \in X$, $\forall \lambda \in [0, 1]$,

$$f((1 - \lambda)x' + \lambda x'') \geq \min\{f(x'), f(x'')\}.$$

f is strictly quasi-concave

Definition 763 iff $\forall x', x'' \in X$, such that $x' \neq x''$, and $\forall \lambda \in (0, 1)$, we have that

$$f((1 - \lambda)x' + \lambda x'') > \min\{f(x'), f(x'')\}.$$

We define the budget correspondence as

Definition 764

$$\beta : \mathbb{R}_{++}^C \times \mathbb{R}_{++} \rightarrow \mathbb{R}^C, \beta(p, w) = \{x \in \mathbb{R}_+^C : px \leq w\}.$$

The Utility Maximization Problem (UMP) is

Definition 765

$$\max_{x \in \mathbb{R}_+^C} u(x) \quad \text{s.t.} \quad px \leq w, \text{ or } x \in \beta(p, w)$$

$\xi : \mathbb{R}_{++}^C \times \mathbb{R}_{++} \rightarrow \mathbb{R}^C$, $\xi(p, w) = \arg \max(\text{UMP})$ is the demand correspondence.

The Profit Maximization Problem (PMP) is

$$\max_y py \quad \text{s.t.} \quad y \in Y.$$

Definition 766 The supply correspondence is

$$y : \mathbb{R}_{++}^C \rightarrow \mathbb{R}^C, y(p) = \arg \max(\text{PMP}).$$

We can now solve some exercises. (the numbering has to be changed)

1.

Show that ξ is non-empty valued.

2.

Show that for every $(p, w) \in \mathbb{R}_{++}^C \times \mathbb{R}_{++}$,

(a) if u is quasiconcave, ξ is convex valued;

(b) if u is strictly quasiconcave, ξ is single valued, i.e., it is a function.

3.

Show that β is closed.

4.

If a solution to (PMP) exists, show the following properties hold.

(a) If Y is convex, $y(\cdot)$ is convex valued;

(b) If Y is strictly convex (i.e., $\forall \lambda \in (0, 1)$, $y^\lambda := (1 - \lambda)y' + \lambda y'' \in \text{Int } Y$), $y(\cdot)$ is single valued.

5

Consider $\phi_1, \phi_2 : [0, 2] \rightarrow \mathbb{R}$,

$$\phi_1(x) = \begin{cases} [-1 + 0.25 \cdot x, x^2 - 1] & \text{if } x \in [0, 1) \\ [-1, 1] & \text{if } x = 1 \\ [-1 + 0.25 \cdot x, x^2 - 1] & \text{if } x \in (1, 2] \end{cases},$$

and

$$\phi_2(x) = \begin{cases} [-1 + 0.25 \cdot x, x^2 - 1] & \text{if } x \in [0, 1) \\ [-0.75, -0.25] & \text{if } x = 1 \\ [-1 + 0.25 \cdot x, x^2 - 1] & \text{if } x \in (1, 2] \end{cases}.$$

¹The definition of quasi-concavity and strict quasi-concavity will be studied in detail in Chapter .

Say if ϕ_1 and ϕ_2 are LHC, UHC, closed, convex valued, compact valued.

6.

Consider $\phi : \mathbb{R}_+ \rightarrow \mathbb{R}$,

$$\phi(x) = \begin{cases} \{\sin \frac{1}{x}\} & \text{if } x > 0 \\ [-1, 1] & \text{if } x = 0 \end{cases},$$

Say if ϕ is LHC, UHC, closed.

7.

Consider $\phi : [0, 1] \rightarrow \mathbb{R}$,

$$\phi(x) = \begin{cases} [0, 1] & \text{if } x \in \mathbb{Q} \cap [0, 1] \\ [-1, 0] & \text{if } x \in [0, 1] \setminus \mathbb{Q} \end{cases}.$$

Say if ϕ is LHC, UHC, closed.

8.

Consider $\phi_1, \phi_2 : [0, 3] \rightarrow \mathbb{R}$,

$$\phi_1(x) = [x^2 - 2, x^2],$$

and

$$\phi_2(x) = [x^2 - 3, x^2 - 1],$$

$$\phi_3(x) := (\phi_1 \cap \phi_2)(x) := \phi_1(x) \cap \phi_2(x).$$

Say if ϕ_1, ϕ_2 and ϕ_3 are LHC, UHC, closed.

20.3 Differential Calculus in Euclidean Spaces

1.

Using the definition, compute the partial derivative of the following function in an arbitrary point (x_0, y_0) :

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f(x, y) = 2x^2 - xy + y^2.$$

2.

If possible, compute partial derivatives of the following functions.

- $f(x, y) = x \cdot \arctan \frac{y}{x}$;
- $f(x, y) = x^y$;
- $f(x, y) = (\sin(x + y))^{\sqrt{x+y}}$ in $(0, 3)$

3.

Given the function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$f(x, y) = \begin{cases} \frac{xy}{x^2+y^2} & \text{if } x+y \neq 0 \\ 0 & \text{otherwise} \end{cases},$$

show that it admits both partial derivatives in $(0, 0)$ and it is not continuous in $(0, 0)$.

4.

Using the definition, compute the directional derivative $f'((1, 1); (\alpha_1, \alpha_2))$ with $\alpha_1, \alpha_2 \neq 0$ for $f : \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$f(x, y) = \frac{x+y}{x^2+y^2+1}.$$

5.

Using the definition, show that the following function is differentiable: $f : \mathbb{R}^2 \rightarrow \mathbb{R}$,

$$f(x, y) = x^2 - y^2 + xy \tag{20.8}$$

Comment: this exercise requires some tricky computations. Do not spend too much time on it. Do this exercise after having studied Proposition 641.

6 .

Using the definition, show that the following functions are differentiable.

- a. $l \in L(\mathbb{R}^n, \mathbb{R}^m)$;
- b. the projection function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $f : (x^i)_{i=1}^n \mapsto x^1$.

7.

Show the following result which was used in the proof of Proposition 606. A linear function $l : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is continuous.

8 .

Compute the Jacobian matrix of $f : \mathbb{R}^2 \rightarrow \mathbb{R}^3$,

$$f(x, y) = (\sin x \cos y, \quad \sin x \sin y, \quad \cos x \cos y)$$

9 .

Given differentiable functions $g, h : \mathbb{R} \rightarrow \mathbb{R}$ and $y \in \mathbb{R} \setminus \{0\}$, compute the Jacobian matrix of $f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, $f(x, y, z) = \left(g(x) \cdot h(z), \quad \frac{g(h(x))}{y}, \quad e^{x \cdot g(h(x))} \right)$

10 .

Compute total derivative and directional derivative at x_0 in the direction u .

a.

$$f : \mathbb{R}_{++}^3 \rightarrow \mathbb{R}, \quad f(x_1, x_2, x_3) = \frac{1}{3} \log x_1 + \frac{1}{6} \log x_2 + \frac{1}{2} \log x_3$$

$$x_0 = (1, 1, 2), \quad u = \frac{1}{\sqrt{3}}(1, 1, 1);$$

b.

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad f(x_1, x_2, x_3) = x_1^2 + 2x_2^2 - x_3^2 - 2x_1x_2 - 6x_2x_3$$

$$x_0 = (1, 0, -1), \quad u = \left(-\frac{1}{\sqrt{2}}, 0, \frac{1}{\sqrt{2}}\right);$$

c.

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad f(x_1, x_2) = x_1 \cdot e^{x_1 x_2}$$

$$x_0 = (0, 0), \quad u = (2, 3).$$

11 .

Given

$$f(x, y, z) = (x^2 + y^2 + z^2)^{-\frac{1}{2}},$$

show that if $(x, y, z) \neq 0$, then

$$\frac{\partial^2 f(x, y, z)}{\partial x^2} + \frac{\partial^2 f(x, y, z)}{\partial y^2} + \frac{\partial^2 f(x, y, z)}{\partial z^2} = 0$$

12 .

Given the C^2 functions $g, h : \mathbb{R} \rightarrow \mathbb{R}_{++}$, compute the Jacobian matrix of

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^3, \quad f(x, y, z) = \left(\frac{g(x)}{h(z)}, \quad g(h(x)) + xy, \quad \ln(g(x) + h(x)) \right)$$

13 .

Given the functions

$$f : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad f(x, y) = \begin{pmatrix} e^x + y \\ e^y + x \end{pmatrix}$$

$$g : \mathbb{R} \rightarrow \mathbb{R}, \quad x \mapsto g(x)$$

$$h : \mathbb{R} \rightarrow \mathbb{R}^2, \quad h(x) = f(x, g(x))$$

Assume that g is C^2 . a. compute the differential of h in 0; b. check the conclusion of the Chain Rule.

14 .

Let the following differentiable functions be given.

$$f : \mathbb{R}^3 \rightarrow \mathbb{R} \quad (x_1, x_2, x_3) \mapsto f(x_1, x_2, x_3)$$

$$g : \mathbb{R}^3 \rightarrow \mathbb{R} \quad (x_1, x_2, x_3) \mapsto g(x_1, x_2, x_3)$$

$$a : \mathbb{R}^3 \rightarrow \mathbb{R}^3, \quad (x_1, x_2, x_3) \mapsto \begin{pmatrix} f(x_1, x_2, x_3) \\ g(x_1, x_2, x_3) \\ x_1 \end{pmatrix}$$

$$b : \mathbb{R}^3 \rightarrow \mathbb{R}^2, \quad (y_1, y_2, y_3) \mapsto \begin{pmatrix} g(y_1, y_2, y_3) \\ f(y_1, y_2, y_3) \end{pmatrix}$$

Compute the directional derivative of the function $b \circ a$ in the point $(0, 0, 0)$ in the direction $(1, 1, 1)$.

15 .

Using the theorems of Chapter 16, show that the function in (20.8) is differentiable.

16 .

Given

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^1, \quad (x, y, z) \mapsto z + x + y^3 + 2x^2y^2 + 3xyz + z^3 - 9,$$

say if you can apply the Implicit Function Theorem to the function in $(x_0, y_0, z_0) = (1, 1, 1)$ and, if possible, compute $\frac{\partial x}{\partial z}$ and $\frac{\partial y}{\partial z}$ in $(1, 1, 1)$.

17 .

Using the notation of the statement of the Implicit Function Theorem presented in the Class Notes, say if that Theorem can be applied to the cases described below; if it can be applied, compute the Jacobian of g . (Assume that a solution to the system $f(x, t) = 0$ does exist).

a. $f : \mathbb{R}^4 \rightarrow \mathbb{R}^2$,

$$f(x_1, x_2, t_1, t_2) \mapsto \begin{pmatrix} x_1^2 - x_2^2 + 2t_1 + 3t_2 \\ x_1x_2 + t_1 - t_2 \end{pmatrix}$$

b. $f : \mathbb{R}^4 \rightarrow \mathbb{R}^2$,

$$f(x_1, x_2, t_1, t_2) \mapsto \begin{pmatrix} 2x_1x_2 + t_1 + t_2^2 \\ x_1^2 + x_2^2 + t_1^2 - 2t_1t_2 + t_2^2 \end{pmatrix}$$

c. $f : \mathbb{R}^4 \rightarrow \mathbb{R}^2$,

$$f(x_1, x_2, t_1, t_2) \mapsto \begin{pmatrix} t_1^2 - t_2^2 + 2x_1 + 3x_2 \\ t_1t_2 + x_1 - x_2 \end{pmatrix}$$

18.

Say under which conditions, if $z^3 - xz - y = 0$, then

$$\frac{\partial^2 z}{\partial x \partial y} = -\frac{3z^2 + x}{(3z^2 - x)^3}$$

19. Do Exercise 656: Let the utility function $u : \mathbb{R}_{++}^2 \rightarrow \mathbb{R}_{++}$, $(x, y) \mapsto u(x, y)$ be given. Assume that it satisfies the following properties i. u is C^2 , ii. $\forall (x, y) \in \mathbb{R}_{++}^2$, $Du(x, y) >> 0$, iii.

$\forall (x, y) \in \mathbb{R}_{++}^2$, $D_{xx}u(x, y) < 0$, $D_{yy}u(x, y) < 0$, $D_{xy}u(x, y) > 0$. Compute the Marginal Rate of Substitution in (x_0, y_0) and say if the graph of each indifference curve is concave.

20.

Let the function $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$ be given and assume that

for every $x_0 \in \mathbb{R}^n$ and every $u \in \mathbb{R}^n$, the directional derivatives $f'(x_0; u)$ and $g'(x_0; u)$ do exist.

Define $h : \mathbb{R}^n \rightarrow \mathbb{R}$, $x \mapsto f(x) \cdot g(x)$. If possible, compute $h'(x_0; u)$ for every $x_0 \in \mathbb{R}^n$ and every $u \in \mathbb{R}^n$.

21.

A function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, $x \mapsto f(x)$ is homogenous of degree $n \in \mathbb{N}_+$ if

$$\text{for every } x = (x_1, x_2) \in \mathbb{R}^2 \text{ and every } a \in \mathbb{R}_+, f(ax_1, ax_2) = a^n f(x_1, x_2).$$

Show that if f is homogenous of degree n and $f \in C^1(\mathbb{R}^2, \mathbb{R})$, then

$$\text{for every } x = (x_1, x_2) \in \mathbb{R}^2, \quad x_1 \cdot D_{x_1}f(x_1, x_2) + x_2 \cdot D_{x_2}f(x_1, x_2) = nf(x_1, x_2).$$

20.4 Nonlinear Programming

1. ² Determine, if possible, the nonnegative parameter values for which the following functions $f : X \rightarrow \mathbb{R}$, $f : (x_i)_{i=1}^n := x \mapsto f(x)$ are concave, pseudo-concave, quasi-concave, strictly concave.

(a) $X = \mathbb{R}_{++}$, $f(x) = \alpha x^\beta$;

(b) $X = \mathbb{R}_{++}^n$, $n \geq 2$, $f(x) = \sum_{i=1}^n \alpha_i (x_i)^{\beta_i}$ (for pseudo-concavity and quasi-concavity consider only the case $n = 2$).

(c) $X = \mathbb{R}$, $f(x) = \min\{\alpha, \beta x - \gamma\}$.

2.

a. Discuss the following problem. For given $\pi \in (0, 1)$, $a \in (0, +\infty)$,

$$\begin{aligned} \max_{(x_1, x_2)} \pi \cdot u(x) + (1 - \pi) u(y) \quad \text{s.t.} \quad & y \leq a - \frac{1}{2}x \\ & y \leq 2a - 2x \\ & x \geq 0 \\ & y \geq 0 \end{aligned}$$

where $u : \mathbb{R} \rightarrow \mathbb{R}$ is a C^2 function such that $\forall z \in \mathbb{R}$, $u'(z) > 0$ and $u''(z) < 0$.

b. Say if there exist values of (π, a) such that $(x, y, \lambda_1, \lambda_2, \lambda_3, \lambda_4) = (\frac{2}{3}a, \frac{2}{3}a, \lambda_1, 0, 0, 0)$, with $\lambda_1 > 0$, is a solution to Kuhn-Tucker conditions, where for $j \in \{1, 2, 3, 4\}$, λ_j the multiplier associated with constraint j .

c. "Assuming" that the first, third and fourth constraint hold with a strict inequality, and the multiplier associated with the second constraint is strictly positive, describe in detail how to compute the effect of a change of a or π on a solution of the problem.

3.

a. Discuss the following problem. For given $\pi \in (0, 1)$, $w_1, w_2 \in \mathbb{R}_{++}$,

$$\begin{aligned} \max_{(x, y, m) \in \mathbb{R}_{++}^2 \times \mathbb{R}} \pi \log x + (1 - \pi) \log y \quad \text{s.t.} \quad & \\ & w_1 - m - x \geq 0 \\ & w_2 + m - y \geq 0 \end{aligned}$$

b. Compute the effect of a change of w_1 on the component x^* of the solution.

c. Compute the effect of a change of π on the objective function computed at the solution of the problem.

4.

²Exercise 1 is taken from David Cass' problem sets for his Microeconomics course at the University of Pennsylvania.

a. Discuss the following problem.

$$\begin{aligned} \min_{(x,y) \in \mathbb{R}^2} \quad & x^2 + y^2 - 4x - 6y \quad s.t. \quad \begin{aligned} & x + y \leq 6 \\ & y \leq 2 \\ & x \geq 0 \\ & y \geq 0 \end{aligned} \end{aligned}$$

Let (x^*, y^*) be a solution to the the problem.

b. Can it be $x^* = 0$?

c. Can it be $(x^*, y^*) = (2, 2)$?

5.

Characterize the solutions to the following problems.

(a) (consumption-investment)

$$\begin{aligned} \max_{(c_1, c_2, k) \in \mathbb{R}^3} \quad & u(c_1) + \delta u(c_2) \\ s.t. \quad & c_1 + k \leq e \\ & c_2 \leq f(k) \\ & c_1, c_2, k \geq 0, \end{aligned}$$

where $u : \mathbb{R} \rightarrow \mathbb{R}$, $u' > 0, u'' < 0$; $f : \mathbb{R}_+ \rightarrow \mathbb{R}$, $f' > 0, f'' < 0$ and such that $f(0) = 0$; $\delta \in (0, 1), e \in \mathbb{R}_{++}$. After having written Kuhn Tucker conditions, consider just the case in which $c_1, c_2, k > 0$.

(b) (labor-leisure)

$$\begin{aligned} \max_{(x, l) \in \mathbb{R}^2} \quad & u(x, l) \\ s.t. \quad & px + wl \leq w\bar{l} \\ & l \leq \bar{l} \\ & x, l \geq 0, \end{aligned}$$

where $u : \mathbb{R}^2$ is C^2 , $\forall (x, l) \quad Du(x, l) \gg 0$, u is differentially strictly quasi-concave, i.e., $\forall (x, l)$, if $\Delta \neq 0$ and $Du(x, l) \cdot \Delta = 0$, then $\Delta^T D^2 u \Delta < 0$; $p > 0$, $w > 0$ and $\bar{l} > 0$.

Describe solutions for which $x > 0$ and $0 < l < \bar{l}$,

6.

(a) Consider the model described in Exercise 6. (a). What would be the effect on consumption (c_1, c_2) of an increase in initial endowment e ?

What would be the effect on (the value of the objective function computed at the solution of the problem) of an increase in initial endowment e ?

Assume that $f(k) = ak^\alpha$, with $a \in \mathbb{R}_{++}$ and $\alpha \in (0, 1)$. What would be the effect on consumption (c_1, c_2) of an increase in a ?

(b) Consider the model described in Exercise 6. (b). What would be the effect on leisure l of an increase in the wage rate w ? in the price level p ?

What would be the effect on (the value of the objective function computed at the solution of the problem) of an increase in the wage rate w ? in the price level p ?

7.

Show that if $f : \mathbb{R}^2 \rightarrow \mathbb{R}$, $x \mapsto f(x)$ is homogenous of degree 1, then

$$f \text{ is concave} \Leftrightarrow \text{for any } x, y \in \mathbb{R}^2, f(x + y) \geq f(x) + f(y).$$

Chapter 21

Solutions

21.1 Linear Algebra

1.

We want to prove that the following sets are **not** vector spaces. Thus, it is enough to find a counter-example violating one of the conditions defining vector spaces.

(i) The definition violates the so-called M2 distributive assumption since for $k = 1$ and $a = b = 1$,

$$(1 + 1) \cdot (1, 1) = 2 \cdot (1, 1) = (2, 1) : \text{ while } : 1 \cdot (1, 1) + 1 \cdot (1, 1) = (2, 2)$$

(ii) The definition violates the so-called A4 commutative property since for $a = b = c = 1$ and $d = 0$,

$$(1, 1) + (1, 0) = (2, 1) \neq (1, 0) + (1, 1) = (2, 0)$$

2.

(i) Take any $w := (x, y, z) \in W$ with $z > 0$, and $\alpha \in \mathbb{R}$ with $\alpha < 0$; then $\alpha w = (\alpha x, \alpha y, \alpha z)$ with $\alpha z < 0$ and therefore $w \notin W$.

(ii) Take any nonzero $w \in W$ and define $\alpha = 2/\|w\|$. Observe that $\|\alpha w\| = 2 > 1$ and therefore $\alpha w \notin W$.

(iii) Multiplication of any nonzero element of \mathbb{Q}^3 by $\alpha \in \mathbb{R} \setminus \mathbb{Q}$ will give an element of $\mathbb{R}^3 \setminus \mathbb{Q}^3$ instead of \mathbb{Q}^3 .

3.

We use Proposition 139. Therefore, we have to check that

a. $0 \in W$; b. $\forall u, v \in W, \forall \alpha, \beta \in F, \alpha u + \beta v \in W$.

Define simply by 0 the function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $\forall x \in \mathbb{R}, f(x) = 0$.

(i) a. Since $0(1) = 0, 0 \in W$.

b.

$$\alpha u(1) + \beta v(1) = \alpha \cdot 0 + \beta \cdot 0 = 0,$$

where the first equality follows from the assumption that $u, v \in W$. Then, indeed, $\alpha u + \beta v \in W$.

(ii) a. Since $0(1) = 0 = 0(2)$, we have that $0 \in W$.

b.

$$\alpha u(1) + \beta v(1) = \alpha u(2) + \beta v(2),$$

where the equality follows from the assumption that $u, v \in W$ and therefore $u(1) = u(2)$ and $v(1) = v(2)$.

4.

Again we use Proposition 139 and we have to check that

a. $0 \in W$; b. $\forall u, v \in W, \forall \alpha, \beta \in F, \alpha u + \beta v \in W$.

a. $(0, 0, 0) \in W$ simply because $0 + 0 + 0 = 0$.

b. Given $u = (u_1, u_2, u_3), v = (v_1, v_2, v_3) \in V$, i.e., such that

$$u_1 + u_2 + u_3 = 0 \quad \text{and} \quad v_1 + v_2 + v_3 = 0, \tag{21.1}$$

we have

$$\alpha u + \beta v = (\alpha u_1 + \beta v_1, \alpha u_2 + \beta v_2, \alpha u_3 + \beta v_3).$$

Then,

$$\alpha u_1 + \beta v_1 + \alpha u_2 + \beta v_2 + \alpha u_3 + \beta v_3 = \alpha(u_1 + u_2 + u_3) + \beta(v_1 + v_2 + v_3) \stackrel{(21.1)}{=} 0.$$

(ii) We have to check that a. S is linearly independent and b. $\text{span } S = V$.

a.

$$\alpha(1, -1, 0) + \beta(0, 1, -1) = (\alpha - \alpha + \beta - \beta) = 0$$

implies that $\alpha = \beta = 0$.

b. Taken $(x_1, x_2, x_3) \in V$, we want to find $\alpha, \beta \in \mathbb{R}$ such that $(x_1, x_2, x_3) = \alpha(1, -1, 0) + \beta(0, 1, -1) = (\alpha - \alpha + \beta - \beta)$, i.e., we want to find $\alpha, \beta \in \mathbb{R}$ such that

$$\begin{cases} x_1 &= \alpha \\ x_2 &= -\alpha + \beta \\ x_3 &= -\beta \\ x_1 + x_2 + x_3 &= 0 \end{cases} \quad \begin{cases} -x_2 - x_3 &= \alpha \\ x_2 &= -\alpha + \beta \\ x_3 &= -\beta \end{cases}$$

Then, $\alpha = -x_2 - x_3$, $\beta = -x_3$ is the (unique) solution to the above system.

5.

1. $0 \in \mathbb{M}(n, n) : A0 = 0A = 0$.
2. $\forall \alpha, \beta \in \mathbb{R}$ and $\forall B, B' \in \mathcal{C}_A$,

$$(\alpha B + \beta B')A = \alpha BA + \beta B'A = \alpha AB + \beta AB' = A\alpha B + A\beta B' = A(\alpha B + \beta B').$$

6.

- i. $0 \in U + V$, because $0 \in U$ and $0 \in V$.
- ii. Take $\alpha, \beta \in F$ and $w^1, w^2 \in U + V$. Then there exists $u^1, u^2 \in U$ and $v^1, v^2 \in V$ such that $w^1 = u^1 + v^1$ and $w^2 = u^2 + v^2$. Therefore,

$$\alpha w^1 + \beta w^2 = \alpha(u^1 + v^1) + \beta(u^2 + v^2) = (\alpha u^1 + \beta u^2) + (\alpha v^1 + \beta v^2) \in U + V,$$

because U and V are vector spaces and therefore $\alpha u^1 + \beta u^2 \in U$ and $\alpha v^1 + \beta v^2 \in V$.

7.

We want to show that if $\sum_{i=1}^3 \beta_i v_i = 0$, then $\beta_i = 0$ for all i . Note that $\sum_{i=1}^3 \beta_i v_i = (\beta_1, \beta_1 + \beta_2, \beta_1 + \beta_2 + \beta_3) = 0$, which implies the desired result.

8.

We want to apply Definition ?? : Consider a vector space V of dimension n and two bases $\mathbf{v} = \{v^i\}_{i=1}^n$ and $\mathbf{u} = \{u^k\}_{k=1}^n$ of V . Then,

$$P = \begin{bmatrix} [u^1]_{\mathbf{v}} & \dots & [u^k]_{\mathbf{v}} & \dots & [u^n]_{\mathbf{v}} \end{bmatrix} \in \mathbb{M}(n, n),$$

is called the change-of-basis matrix from the basis \mathbf{v} to the basis \mathbf{u} . Then in our case, the change-of-basis matrix from S to E is

$$P = \begin{bmatrix} [e^1]_S & [e^2]_S \end{bmatrix} \in \mathbb{M}(2, 2).$$

Moreover, using also Proposition ??, the change-of-basis matrix from S to E is

$$Q = \begin{bmatrix} [v^1]_E & [v^2]_E \end{bmatrix} = P^{-1}.$$

Computation of $[e^1]_S$. We want to find α and β such that $e_1 = \alpha u_1 + \beta u_2$, i.e., $(1, 0) = \alpha(1, 2) + \beta(3, 5) = (\alpha + 3\beta \quad 2\alpha + 5\beta)$, i.e.,

$$\begin{cases} \alpha + 3\beta &= 1 \\ 2\alpha + 5\beta &= 0 \end{cases}$$

whose solution is $\alpha = -5, \beta = 2$.

Computation of $[e^2]_S$. We want to find α and β such that $e_2 = \alpha u_1 + \beta u_2$, i.e., $(0, 1) = \alpha(1, 2) + \beta(3, 5) = \begin{pmatrix} \alpha + 3\beta & 2\alpha + 5\beta \end{pmatrix}$, i.e.,

$$\begin{cases} \alpha + 3\beta = 0 \\ 2\alpha + 5\beta = 1 \end{cases}$$

$$\begin{cases} \alpha + 3\beta = 0 \\ 2\alpha + 5\beta = 1 \end{cases}$$

whose solution is $\alpha = 3, \beta = -1$. Therefore,

$$P = \begin{bmatrix} -5 & 3 \\ 2 & -1 \end{bmatrix}.$$

Since E is the canonical basis, we have

$$S = \{u_1 = (1, 2), u_2 = (3, 5)\}$$

$$Q = \begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix}.$$

Finally

$$\begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix}^{-1} = \begin{bmatrix} -5 & 3 \\ 2 & -1 \end{bmatrix}$$

as desired.

9.

Easiest way: use row and column operations to change C to a triangular matrix.

$$\begin{aligned} \det C &= \det \begin{bmatrix} 6 & 2 & 1 & 0 & 5 \\ 2 & 1 & 1 & -2 & 1 \\ 1 & 1 & 2 & -2 & 3 \\ 3 & 0 & 2 & 3 & -1 \\ -1 & -1 & -3 & 4 & 2 \end{bmatrix} = [R^1 \leftrightarrow R^3] = -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 2 & 1 & 1 & -2 & 1 \\ 6 & 2 & 1 & 0 & 5 \\ 3 & 0 & 2 & 3 & -1 \\ -1 & -1 & -3 & 4 & 2 \end{bmatrix} \\ &= -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 2 & 1 & 1 & -2 & 1 \\ 6 & 2 & 1 & 0 & 5 \\ 3 & 0 & 2 & 3 & -1 \\ -1 & -1 & -3 & 4 & 2 \end{bmatrix} = \begin{bmatrix} -2R^1 + R^2 \rightarrow R^2 \\ -6R^1 + R^3 \rightarrow R^3 \\ -3R^1 + R^4 \rightarrow R^4 \\ R^1 + R^5 \rightarrow R^5 \end{bmatrix} = -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 0 & -1 & -3 & 2 & -5 \\ 0 & -4 & -11 & 12 & -13 \\ 0 & -3 & -4 & 9 & -10 \\ 0 & 0 & -1 & 2 & 5 \end{bmatrix} \\ &= -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 0 & -1 & -3 & 2 & -5 \\ 0 & -4 & -11 & 12 & -13 \\ 0 & -3 & -4 & 9 & -10 \\ 0 & 0 & -1 & 2 & 5 \end{bmatrix} = \begin{bmatrix} 4R^2 + R^3 \rightarrow R^3 \\ 3R^2 + R^4 \rightarrow R^4 \end{bmatrix} = -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 0 & -1 & -3 & 2 & -5 \\ 0 & 0 & 1 & 4 & 7 \\ 0 & 0 & 5 & 3 & 5 \\ 0 & 0 & -1 & 2 & 5 \end{bmatrix} \\ &= -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 0 & -1 & -3 & 2 & -5 \\ 0 & 0 & 1 & 4 & 7 \\ 0 & 0 & 5 & 3 & 5 \\ 0 & 0 & -1 & 2 & 5 \end{bmatrix} = \begin{bmatrix} -5R^3 + R^4 \rightarrow R^4 \\ R^3 + R^5 \rightarrow R^5 \end{bmatrix} = -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 0 & -1 & -3 & 2 & -5 \\ 0 & 0 & 1 & 4 & 7 \\ 0 & 0 & 0 & -17 & -30 \\ 0 & 0 & 0 & 6 & 12 \end{bmatrix} \\ &= -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 0 & -1 & -3 & 2 & -5 \\ 0 & 0 & 1 & 4 & 7 \\ 0 & 0 & 0 & -17 & -30 \\ 0 & 0 & 0 & 6 & 12 \end{bmatrix} = \begin{bmatrix} 6 \\ 17 \end{bmatrix} R^4 + R^5 \rightarrow R^5 = -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 0 & -1 & -3 & 2 & -5 \\ 0 & 0 & 1 & 4 & 7 \\ 0 & 0 & 0 & -17 & -30 \\ 0 & 0 & 0 & 0 & \frac{24}{17} \end{bmatrix} \end{aligned}$$

$$= -\det \begin{bmatrix} 1 & 1 & 2 & -2 & 3 \\ 0 & -1 & -3 & 2 & -5 \\ 0 & 0 & 1 & 4 & 7 \\ 0 & 0 & 0 & -17 & -30 \\ 0 & 0 & 0 & 0 & \frac{24}{17} \end{bmatrix} = -(1)(-1)(1)(-17)\left(\frac{24}{17}\right) \implies \det C = -24$$

10.

Observe that it cannot be 4 as $\text{rank}(A) \leq \min\{\#rows, \#columns\}$. It's easy to check that $\text{rank}(A) = 3$ by using elementary operations on rows and columns of A :

$$-2R^3 + R^1 \rightarrow R^1, C^4 + C^1 \rightarrow C^1, C^4 + C^3 \rightarrow C^3, C^1 + C^3 \rightarrow C^3, -C^2 + C^3 \rightarrow C^3,$$

to get

$$\begin{bmatrix} k-1 & 1 & 0 & 0 \\ 0 & 0 & 1 & k \\ 0 & 0 & 0 & -1 \end{bmatrix}$$

which has the last three columns independent for any k .

11.

Defined

$$l : \mathbb{R}^3 \rightarrow \mathbb{R}, \quad (x_1, x_2, x_3) \mapsto x_1 - x_2,$$

it is easy to check that l is linear and $V = \ker l$, a vector space. Moreover,

$$[l] = \begin{bmatrix} 1 & -1 & 0 \end{bmatrix}.$$

Therefore, $\dim \ker l = 3 - \text{rank } [l] = 3 - 1 = 2$. $u_1 = (1, 1, 0)$ and $u_2 = (0, 0, 1)$ are independent elements of V . Therefore, from Remark 200, u_1, u_2 are a basis for V .

12.

Linearity is easy. By definition, l is linear if $\forall u, v \in \mathbb{R}^4$ and $\forall \alpha, \beta \in \mathbb{R}$, $l(\alpha u + \beta v) = \alpha l(u) + \beta l(v)$. Then,

$$\begin{aligned} \alpha l(u) + \beta l(v) &= \\ = \alpha(u_1, u_1 + u_2, u_1 + u_2 + u_3, u_1 + u_2 + u_3 + u_4) + \beta(v_1, v_1 + v_2, v_1 + v_2 + v_3, v_1 + v_2 + v_3 + v_4) &= \\ = (\alpha u_1 + \beta v_1, \alpha u_1 + \alpha u_2 + \beta v_1 + \beta v_2, \alpha u_1 + \alpha u_2 + \alpha u_3 + \beta v_1 + \beta v_2 + \beta v_3, \alpha u_1 + \alpha u_2 + \alpha u_3 + \alpha u_4 + \beta v_1 + \beta v_2 + \beta v_3 + \beta v_4) &= \\ = l(\alpha u + \beta v) \end{aligned}$$

and then l is linear.

$$[l] = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 1 \end{bmatrix}.$$

Therefore, $\dim \ker l = 4 - \text{rank } [l] = 4 - 4 = 0$. Moreover, $\dim \text{Im } l = 4$ and the column vectors of $[l]$ are a basis of $\text{Im } l$.

13.

Proposition. Assume that $l \in L(V, U)$ and $\ker l = \{0\}$. Then,

$$\forall u \in \text{Im } l, \quad \text{there exists a unique } v \in V \text{ such that } l(v) = u.$$

Proof.

Since $u \in \text{Im } l$, by definition, there exists $v \in V$ such that

$$l(v) = u. \tag{21.2}$$

Take $v' \in V$ such that $l(v') = u$. We want to show that

$$v = v'. \tag{21.3}$$

Observe that

$$l(v) - l(v') \stackrel{(a)}{=} u - u = 0, \quad (21.4)$$

where (a) follows from (21.2) and (21.3).

Moreover,

$$l(v) - l(v') \stackrel{(b)}{=} l(v - v'), \quad (21.5)$$

where (b) follows from the assumption that $l \in \mathcal{L}(V, U)$.

Therefore,

$$l(v - v') = 0,$$

and, by definition of $\ker l$,

$$v - v' \in \ker l. \quad (21.6)$$

Since, by assumption, $\ker l = \{0\}$, from (21.6), it follows that

$$v - v' = 0.$$

14.

Both V and W are \ker of linear function; therefore V , W and $V \cap W$ are vector subspaces of \mathbb{R}^4 . Moreover

$$V \cap W = \left\{ (x_1, x_2, x_3, x_4) \in \mathbb{R}^4 : \begin{cases} x_1 - x_2 + x_3 - x_4 = 0 \\ x_1 + x_2 + x_3 + x_4 = 0 \end{cases} \right\}$$

$$\text{rank} \begin{bmatrix} 1 & -1 & 1 & -1 \\ 1 & 1 & 1 & 1 \end{bmatrix} = 2$$

Therefore, $\dim \ker l = \dim V \cap W = 4 - 2 = 2$.

Let's compute a basis of $V \cap W$:

$$\begin{cases} x_1 - x_2 = -x_3 + x_4 \\ x_1 + x_2 = -x_3 - x_4 \end{cases}$$

After taking sum and subtraction we get following expression

$$\begin{cases} x_1 = -x_3 \\ x_2 = -x_4 \end{cases}$$

A basis consists two linearly independent vectors. For example,

$$\{(-1, 0, 1, 0), (0, -1, 0, 0)\}.$$

15.

By proposition 139, we have to show that

1. $0 \in l^{-1}(W)$,
2. $\forall \alpha, \beta \in \mathbb{R}$ and $v^1, v^2 \in l^{-1}(W)$ we have that $\alpha v^1 + \beta v^2 \in l^{-1}(W)$.

1.

$$l(0) \stackrel{l \in \mathcal{L}(V, U)}{=} 0 \stackrel{W \text{ vector space}}{\in} W$$

,

2. Since $v^1, v^2 \in l^{-1}(W)$,

$$l(v^1), l(v^2) \in W. \quad (21.7)$$

Then

$$l(\alpha v^1 + \beta v^2) \stackrel{l \in \mathcal{L}(V, U)}{=} \alpha l(v^1) + \beta l(v^2) \stackrel{(a)}{\in} W$$

where (a) follows from (21.7) and the fact that W is a vector space.

16.

Observe that

$$\det \begin{bmatrix} a_{11} & a_{12} & 0 & 0 & 0 \\ a_{21} & a_{22} & 0 & 0 & 0 \\ 5 & 6 & b_{11} & b_{12} & 0 \\ 7 & 8 & b_{21} & b_{22} & 0 \\ a_{11} & a_{12} & 0 & 0 & k \end{bmatrix} = \det A \cdot \det B \cdot k.$$

Then, if $k \neq 0$, then the rank of both matrix of coefficients and augmented matrix is 5 and the set of solution to the system is an affine subspace of \mathbb{R}^6 of dimension 1. If $k = 0$, then the system is

$$\begin{bmatrix} 1 & a_{11} & a_{12} & 0 & 0 & 0 \\ 2 & a_{21} & a_{22} & 0 & 0 & 0 \\ 3 & 5 & 6 & b_{11} & b_{12} & 0 \\ 4 & 7 & 8 & b_{21} & b_{22} & 0 \\ 1 & a_{11} & a_{12} & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 3 \\ 0 \end{bmatrix},$$

which is equivalent to the system

$$\begin{bmatrix} 1 & a_{11} & a_{12} & 0 & 0 & 0 \\ 2 & a_{21} & a_{22} & 0 & 0 & 0 \\ 3 & 5 & 6 & b_{11} & b_{12} & 0 \\ 4 & 7 & 8 & b_{21} & b_{22} & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 2 \\ 3 \end{bmatrix},$$

whose set of solution is an affine subspace of \mathbb{R}^6 of dimension 2.

17.

$$[l]_{\mathbf{v}}^{\mathbf{u}} := \left[\left[l \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right]_{\mathbf{u}}, \left[l \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right]_{\mathbf{u}} \right] = \left[\begin{bmatrix} 1 \\ 1 \end{bmatrix}_{\mathbf{u}}, \begin{bmatrix} 1 \\ -1 \end{bmatrix}_{\mathbf{u}} \right] = \begin{bmatrix} 1 & -3 \\ 0 & 2 \end{bmatrix},$$

$$[v]_{\mathbf{v}} = \begin{bmatrix} 3 \\ 4 \end{bmatrix}$$

$$[l(v)]_{\mathbf{u}} = \left[\begin{pmatrix} 7 \\ -1 \end{pmatrix} \right]_{\mathbf{u}} = \begin{bmatrix} -9 \\ 8 \end{bmatrix}$$

$$[l]_{\mathbf{v}}^{\mathbf{u}} \cdot [v]_{\mathbf{v}} = \begin{bmatrix} 1 & -3 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 3 \\ 4 \end{bmatrix} = \begin{bmatrix} -9 \\ 8 \end{bmatrix}$$

18.

Let

$n, m \in \mathbb{N}$ such that $m > n$, and

a vector subspace L of \mathbb{R}^m such that $\dim L = n$

be given. Then, there exists $l \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ such that $\text{Im } l = L$.

Proof. Let $\{v^i\}_{i=1}^n$ be a basis of $L \subseteq \mathbb{R}^m$. Take $l \in \mathcal{L}(\mathbb{R}^n, \mathbb{R}^m)$ such that

$$\forall i \in \{1, \dots, n\}, l_2(e_n^i) = v^i,$$

where e_n^i is the i -th element in the canonical basis in \mathbb{R}^n . Such function does exist and, in fact, it is unique as a consequence of a Proposition in the Class Notes that we copy below:

Let V and U be finite dimensional vector spaces such that $S = \{v^1, \dots, v^n\}$ is a basis of V and $\{u^1, \dots, u^n\}$ is a set of arbitrary vectors in U . Then there exists a unique linear function $l : V \rightarrow U$ such that $\forall i \in \{1, \dots, n\}, l(v^i) = u^i$ - see Proposition 273, page 82.

Then, from the Dimension theorem

$$\dim \text{Im } l = n - \dim \ker l \leq n.$$

Moreover, $L = \text{span } \{v^i\}_{i=1}^n \subseteq \text{Im } l$. Summarizing,

$$L \subseteq \text{Im } l, \dim L = n \text{ and } \dim \text{Im } l \leq n,$$

and therefore

$$\dim \operatorname{Im} l = n.$$

Finally, from Proposition 179 in the class Notes since $L \subseteq \operatorname{Im} l$, $\dim L = n$ and $\dim \operatorname{Im} l = n$, we have that $\operatorname{Im} l = L$, as desired.

Proposition 179 in the class Notes says what follows: Proposition. Let W be a subspace of an n -dimensional vector space V . Then, 1. $\dim W \leq n$; 2. If $\dim W = n$, then $W = V$.

19.

$$[A \mid b] = \begin{bmatrix} a & 1 & 1 \\ 1 & 1 & a \\ 2 & 1 & 3a \\ 3 & 2 & a \end{bmatrix} \Rightarrow [R^1 \leftrightarrow R^2] \Rightarrow \begin{bmatrix} 1 & 1 & a \\ a & 1 & 1 \\ 2 & 1 & 3a \\ 3 & 2 & a \end{bmatrix}$$

$$\begin{bmatrix} 1 & 1 & a \\ a & 1 & 1 \\ 2 & 1 & 3a \\ 3 & 2 & a \end{bmatrix} \Rightarrow \begin{matrix} -aR^1 + R^2 \rightarrow R^2 \\ -2R^1 + R^3 \rightarrow R^3 \\ -3R^1 + R^4 \rightarrow R^4 \end{matrix} \Rightarrow \begin{bmatrix} 1 & 1 & a \\ 0 & 1-a & 1-a^2 \\ 0 & -1 & a \\ 0 & -1 & -2a \end{bmatrix} := [A'(a) \mid b'(a)]$$

Since

$$\det \begin{bmatrix} 1 & 1 & a \\ 0 & -1 & a \\ 0 & -1 & -2a \end{bmatrix} = 3a,$$

We have that if $a \neq 0$, then $\operatorname{rank} A \leq 2 < 3 = \operatorname{rank} [A'(a) \mid b'(a)]$, and the system has no solutions. If $a = 0$, $[A'(a) \mid b'(a)]$ becomes

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & -1 & 0 \\ 0 & -1 & 0 \end{bmatrix}$$

whose rank is 3 and again the system has no solutions.

20.

$$[A(k) \mid b(k)] \equiv \begin{bmatrix} 1 & 0 & k-1 \\ 1-k & 2-k & k \\ 1 & k & 1 \\ 1 & k-1 & 0 \end{bmatrix}$$

$$\det \begin{bmatrix} 1 & 0 & k-1 \\ 1 & k & 1 \\ 1 & k-1 & 0 \end{bmatrix} = 2-2k$$

If $k \neq 1$, the system has no solutions. If $k = 1$,

$$[A(1) \mid b(1)] \equiv \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 0 & 0 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

Then, if $k = 1$, there exists a unique solution.

21.

The following Proposition is contained in the class Notes.

Proposition 767 *Let V be a vector space of dimension n .*

1. *If $S = \{u^1, \dots, u^n\} \subseteq V$ is a linearly independent set, then it is a basis of V ;*
2. *If $\text{span}(u^1, \dots, u^n) = V$, then $\{u^1, \dots, u^n\}$ is a basis of V .*

From that Proposition, it suffices to show that \mathcal{V} is linearly independent, i.e., given $(\alpha_i)_{i=1}^n \in \mathbb{R}^n$,

if

$$\sum_{i=1}^n \alpha_i v^i = 0 \quad (21.8)$$

then

$$\boxed{(\alpha_i)_{i=1}^n = 0.}$$

Now, for any $j \in \{1, \dots, n\}$, we have

$$0 \stackrel{(1)}{=} \left(\sum_{i=1}^n \alpha_i v^i \right) v^j \stackrel{(2)}{=} \sum_{i=1}^n \alpha_i v^i v^j \stackrel{(3)}{=} \alpha_j,$$

where (1) follows from (21.8),

(2) follows from properties of the scalar product;

(3) follows from (20.7).

22.

a. Let $w \in W$. By assumption, $S(w) \in W$ and $T(w) \in W$ and since W is a subspace, $S(w) + T(w) \in W$. Therefore, $(S + T)(w) = S(w) + T(w) \in W$, as desired.

b. Let $w \in W$. By assumption, $T(w) \in W$. Then $(S \circ T)(w) = S(T(w)) \in W$ since W is S -invariant.

c. Let $w \in W$. By assumption and recalling Proposition 138 in Villanacci(20 September 2012), $(kT)(w) = kT(w) \in W$.

23.

The set of 2×2 symmetric real matrices is

$$\mathcal{S} = \{A \in \mathbb{M}(2, 2) : A = A^T\}.$$

We want to show that

i. $0 \in \mathcal{S}$ and

ii. for any $\alpha, \beta \in \mathbb{R}$, for any $A, B \in \mathcal{S}$, $\alpha A + \beta B \in \mathcal{S}$.

i. $0 = 0^T -$

2. $(\alpha A + \beta B)^T = \alpha A^T + \beta B^T = \alpha A + \beta B$.

We want to show that

$$\mathcal{B} := \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right\}$$

is a basis of \mathcal{S} and therefore $\dim \mathcal{S} = 3$. \mathcal{B} is clearly linearly independent. Moreover,

$$\mathcal{S} = \left\{ A \in \mathbb{M}(2, 2) : \exists a, b, c \in \mathbb{R} \text{ such that } A = \begin{bmatrix} a & b \\ b & c \end{bmatrix} \right\},$$

and

$$\begin{bmatrix} a & b \\ b & c \end{bmatrix} = a \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + c \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix},$$

i.e., $\text{span}(\mathcal{B}) = \mathcal{S}$, as desired.

24.

a. $[\supseteq]$ Taken $w \in W$, we want to find $w^1, w^2 \in W$ such that $w = w^1 + w^2$. take $w^1 = w$ and $w^2 = 0 \in W$.

$[\subseteq]$ Take $w^1, w^2 \in W$. Then $w^1 + w^2 \in W$ by definition of vector space.

¹see Proposition 138 in Villanacci(20 September, 2012)

- b. $[\supseteq]$ Let $w \in W$. Then $\frac{1}{\alpha}w \in W$ and $\alpha\left(\frac{1}{\alpha}w\right)w \in \alpha W$.
 $[\subseteq]$ It follows from the definition of vector space.

25.

The isomorphism is $\varphi : \mathcal{P}_n(\mathbb{R}) \rightarrow \mathbb{R}^{n+1}$

$$f(t) = \sum_{i=0}^n a_i t^i \mapsto (a_i)_{i=0}^n.$$

Indeed, φ is linear

...

and defined $\psi : \mathbb{R}^{n+1} \rightarrow \mathcal{P}_n(\mathbb{R})$,

$$(a_i)_{i=0}^n \mapsto f(t) = \sum_{i=0}^n a_i t^i,$$

we have that $\varphi \circ \psi = id_{\mathcal{P}_n(\mathbb{R})}$ and $\psi \circ \varphi = id_{\mathbb{R}^{n+1}}$

....

21.2 Some topology in metric spaces

21.2.1 Basic topology in metric spaces

1.

To prove that d' is a metric, we have to check the properties listed in Definition 349.

a. $d'(x, y) \geq 0$, $d'(x, y) = 0 \Leftrightarrow x = y$

By definition of $d'(x, y)$, it is always going to be positive as $d(x, y) \geq 0$. Furthermore, $d'(x, y) = 0 \Leftrightarrow d(x, y) = 0 \Leftrightarrow x = y$.

b. $d'(x, y) = d'(y, x)$

Applying the definition

$$d'(x, y) = d'(y, x) \Leftrightarrow \frac{d(x, y)}{1 + d(x, y)} = \frac{d(y, x)}{1 + d(y, x)}$$

but $d(x, y) = d(y, x)$ so we have

$$\frac{d(x, y)}{1 + d(x, y)} = \frac{d(x, y)}{1 + d(x, y)}$$

c. $d'(x, z) \leq d'(x, y) + d'(y, z)$

Applying the definition

$$d'(x, z) \leq d'(x, y) + d'(y, z) \Leftrightarrow \frac{d(x, z)}{1 + d(x, z)} \leq \frac{d(x, y)}{1 + d(x, y)} + \frac{d(y, z)}{1 + d(y, z)}$$

Multiplying both sides by $[1 + d(x, z)][1 + d(x, y)][1 + d(y, z)]$

$$d(x, z)[1 + d(x, y)][1 + d(y, z)] \leq d(x, y)[1 + d(x, z)][1 + d(y, z)] + d(y, z)[1 + d(x, z)][1 + d(x, y)]$$

Simplifying we obtain

$$\boxed{d(x, z) \leq d(x, y) + d(y, z)} + [[1 + d(x, z)][1 + d(x, y)][1 + d(y, z)] + 2[1 + d(x, y)][1 + d(y, z)]]$$

which concludes the proof.

2.

It is enough to show that one of the properties defining a metric does not hold.

It can be $d(f, g) = 0$ and $f \neq g$. Take

$$f(x) = 0, \forall x \in [0, 1],$$

and

$$g(x) = -2x + 1$$

Then,

$$\int_0^1 (-2x + 1) dx = 0.$$

It can be $d(f, g) < 0$. Consider the null function and the function that take value 1 for all x in $[0, 1]$. Then $d(0, 1) = -\int_0^1 1 dx$. by linearity of the Riemann integral, which is equal to -1 . Then, $d(0, 1) < 0$.

3.

Define $S = (a_1, b_1) \times (a_2, b_2)$ and take $x^0 := (x_1^0, x_2^0) \in S$. Then, for $i \in \{1, 2\}$, there exist $\varepsilon_i > 0$ such that $x_i^0 \in B(x_i^0, \varepsilon_i) \subseteq (a_i, b_i)$. Take $\varepsilon = \min\{\varepsilon_1, \varepsilon_2\}$. Then, for $i \in \{1, 2\}$, $x_i^0 \in B(x_i^0, \varepsilon) \subseteq (a_i, b_i)$ and, defined $B = B(x_1^0, \varepsilon) \times B(x_2^0, \varepsilon)$, we have that $x^0 \in B \subseteq S$. It then suffices to show that $B(x^0, \varepsilon) \subseteq B$. Observe that

$$x \in B(x^0, \varepsilon) \Leftrightarrow d(x, x^0) < \varepsilon,$$

$$d((x_1^0, 0), (x_1, 0)) = \sqrt{(x_1^0 - x_1)^2} = |x_1^0 - x_1|,$$

and

$$d((x_1^0, 0), (x_1, 0)) = \sqrt{(x_1^0 - x_1)^2} \leq \sqrt{(x_1^0 - x_1)^2 + (x_1^0 - x_1)^2} = d(x, x^0).$$

4.

Show the second equality in Remark 388:

$$\cap_{n=1}^{+\infty} \left(-\frac{1}{n}, \frac{1}{n}\right) = \{0\}$$

5.

$$S = \left\{-1, +\frac{1}{2}, -\frac{1}{3}, \frac{1}{4}, -\frac{1}{5}, \frac{1}{6}, \dots\right\}$$

The set is not open: it suffices to find $x \in S$ and such that $x \notin \text{Int } S$; take for example -1 . We want to show that it false that

$$\exists \varepsilon > 0 \text{ such that } (-1 - \varepsilon, -1 + \varepsilon) \subseteq S.$$

In fact, $\forall \varepsilon > 0$, $-1 - \frac{\varepsilon}{2} \in (-1 - \varepsilon, -1 + \varepsilon)$, but $-1 - \frac{\varepsilon}{2} \notin S$. The set is not closed. It suffices to show that $\mathcal{F}(S)$ is not contained in S , in fact that $0 \notin S$ (obvious) and $0 \in \mathcal{F}(S)$. We want to show that $\forall \varepsilon > 0$, $B(0, \varepsilon) \cap S \neq \emptyset$. In fact, $(-1)^n \frac{1}{n} \in B(0, \varepsilon)$ if n is even and $(-1)^n \frac{1}{n} = \frac{1}{n} < \varepsilon$. It is then enough to take n even and $n > \frac{1}{\varepsilon}$.

6.

$$A = (0, 10)$$

The set is (\mathbb{R}, d_2) open, as a union of infinite collection of open sets. The set is not closed, because A^c is not open. 10 or 0 do not belongs to $\text{Int}(A^c)$

7.

The solution immediately follow from Definition of boundary of a set: Let a metric space (X, d) and a set $S \subseteq X$ be given. x is an boundary point of S if

any open ball centered in x intersects both S and its complement in X , i.e., $\forall r \in \mathbb{R}_{++}$, $B(x, r) \cap S \neq \emptyset \quad \wedge \quad B(x, r) \cap S^c \neq \emptyset$.

As you can see nothing changes in definition above if you replace the set with its complement.

8.

$$\begin{aligned} x \in (\mathcal{F}(S))^C &\Leftrightarrow x \notin (\mathcal{F}(S)) \\ &\Leftrightarrow \neg (\forall r \in \mathbb{R}_{++}, B(x, r) \cap S \neq \emptyset \wedge B(x, r) \cap S^c \neq \emptyset) \\ &\Leftrightarrow \exists r \in \mathbb{R}_{++} \text{ such that } B(x, r) \cap S = \emptyset \vee B(x, r) \cap S^c \neq \emptyset \\ &\Leftrightarrow \exists r \in \mathbb{R}_{++} \text{ such that } B(x, r) \subseteq S^c \vee B(x, r) \subseteq S \\ &\Leftrightarrow x \in \text{Int } S^c \vee x \in \text{Int } S \\ &\stackrel{(1)}{\Leftrightarrow} \exists r_x^* \in \mathbb{R}_{++} \text{ such that either a. } B(x, r_x^*) \subseteq \text{Int } S^c \text{ or b. } B(x, r_x^*) \subseteq \text{Int } S. \end{aligned} \tag{21.9}$$

where (1) follows from the fact that the Interior of a set is an open set.

If case a. in (21.9) holds true, then, using Lemma 482, $B(x, r_x^*) \subseteq (\mathcal{F}(S))^C$ and similarly for case b., as desired.

9.

	$Int\ S$	$Cl(S)$	$\mathcal{F}(S)$	$D(S)$	$Is(S)$	open or closed
$S = \mathbb{Q}$	\emptyset	\mathbb{R}	\mathbb{R}	\mathbb{R}	\emptyset	neither open nor closed
$S = (0, 1)$	$(0, 1)$	$[0, 1]$	$\{0, 1\}$	$[0, 1]$	\emptyset	open
$S = \{\frac{1}{n}\}_{n \in \mathbb{N}_+}$	\emptyset	$S \cup \{0\}$	$S \cup \{0\}$	$\{0\}$	S	neither open nor closed

10.

a.

Take $S = \mathbb{N}$. Then, $Int\ S = \emptyset$, $Cl(\emptyset) = \emptyset$, and $Cl(Int\ S) = \emptyset \neq \mathbb{N} = S$.

b.

Take $S = \mathbb{N}$. Then, $Cl(S) = \mathbb{N}$, $Int\ \mathbb{N} = \emptyset$, and $Int\ Cl(S) = \emptyset \neq \mathbb{N} = S$.

11.

a.

True. If S is an open bounded interval, then $\exists a, b \in \mathbb{R}$, $a < b$ such that $S = (a, b)$. Take $x \in S$ and $\delta = \min\{|x - a|, |x - b|\}$. Then $I(x, \delta) \subseteq (a, b)$.

b.

False. $(0, 1) \cup (2, 3)$ is an open set, but it is not an open interval.

c.

False. Take $S := \{0, 1\}$. $0 \in \mathcal{F}(S)$, but $0 \notin D(S)$

d. .

False. Take $S = (0, 1)$. $\frac{1}{2} \in D(S)$, but $\frac{1}{2} \notin \mathcal{F}(S)$.

12.

Recall that: A sequence $(x_n)_{n \in \mathbb{N}} \in X^\infty$ is said to be (X, d) convergent to $x_0 \in X$ (or convergent with respect to the metric space (X, d)) if $\forall \epsilon > 0$, $\exists n_0 \in \mathbb{N}$ such that $\forall n > n_0$, $d(x_n, x_0) < \epsilon$.

a.

$(x_n)_{n \in \mathbb{N}} \in \mathbb{R}^\infty$ such that $\forall n \in \mathbb{N}$, $x_n = 1$

Let $\epsilon > 0$ then by definition of $(x_n)_{n \in \mathbb{N}}$, $\forall n > 0$, $d(x_n, 1) = 0 < \epsilon$. So that

$$\lim_{n \rightarrow \infty} x_n = 1$$

b.

$(x_n)_{n \in \mathbb{N}} \in \mathbb{R}^\infty$ such that $\forall n \in \mathbb{N}$, $x_n = \frac{1}{n}$

Let $\epsilon > 0$. Because \mathbb{N} is unbounded, $\exists n_0 \in \mathbb{N}$, such that $n_0 > \frac{1}{\epsilon}$. Then $\forall n > n_0$, $d(x_n, 0) = \frac{1}{n} < \frac{1}{n_0} < \epsilon$. Then, by definition of a limit, we proved that

$$\lim_{n \rightarrow \infty} x_n = 0$$

13.

Take $(x_n)_{n \in \mathbb{N}} \in [0, 1]^\infty$ such that $x_n \rightarrow x_0$; we want to show that $x_0 \in [0, 1]$. Suppose otherwise, i.e., $x_0 \notin [0, 1]$.

Case 1. $x_0 < 0$. By definition of convergence, chosen $\epsilon = -\frac{x_0}{2} > 0$, there exists $n_\epsilon \in \mathbb{N}$ such that $\forall n > n_\epsilon$, $d(x_n, x_0) < \epsilon$, i.e., $|x_n - x_0| < \epsilon = -\frac{x_0}{2}$, i.e., $x_0 + \frac{x_0}{2} < x_n < x_0 - \frac{x_0}{2} = \frac{x_0}{2} < 0$. Summarizing, $\forall n > n_\epsilon$, $x_n \notin [0, 1]$, contradicting the assumption that $(x_n)_{n \in \mathbb{N}} \in [0, 1]^\infty$.

Case 2. $x_0 > 1$. Similar to case 1.

14.

This is Example 7.15, page 150, Morris (2007):

1. In fact, we have the following result: Let (X, d) be a metric space and $A = \{x_1, \dots, x_n\}$ any finite subset of X . Then A is compact, as shown below.

Let O_i , $i \in I$ be any family of open sets such that $A \subseteq \cup_{i \in I} O_i$. Then for each $x_j \in A$, there exists O_{i_j} such that $x_j \in O_{i_j}$. Then $A \subseteq O_{i_1} \cup O_{i_2} \cup \dots \cup O_{i_n}$. Therefore A is compact.

2. Conversely, let A be compact. Then the family of singleton sets $O_x = \{x\}$, $x \in A$ is such that each O_x is open and $A \subseteq \cup_{x \in A} O_x$. Since A is compact, there exists $O_{x_1}, O_{x_2}, \dots, O_{x_n}$ such that $A \subseteq O_{x_1} \cup O_{x_2} \cup \dots \cup O_{x_n}$, that is, $A \subseteq \{x_1, \dots, x_n\}$. Hence, A is finite.

15.

In general it is false. For example in a discrete metric space: see previous exercise.

16.

Take an open ball $B(x, r)$. Consider $\mathcal{S} = \{B(x, r(1 - \frac{1}{n}))\}_{n \in \mathbb{N} \setminus \{0,1\}}$. Observe that \mathcal{S} is an open cover of $B(x, r)$; in fact $\cup_{n \in \mathbb{N} \setminus \{0,1\}} B(x, r(1 - \frac{1}{n})) = B(x, r)$, as shown below.

$$[\subseteq] \quad x' \in \cup_{n \in \mathbb{N} \setminus \{0,1\}} B(x, r(1 - \frac{1}{n})) \Leftrightarrow \exists n_{x'} \in \mathbb{N} \setminus \{0,1\} \text{ such that } x' \in B(x, r(1 - \frac{1}{n_{x'}})) \subseteq B(x, r).$$

$[\supseteq]$ Take $x' \in B(x, r)$. Then, $d(x, x') < r$. Take n such that $d(x', x) < r(1 - \frac{1}{n_{x'}})$, i.e., $n > \frac{r}{r - d(x', x)}$ (and $n > 1$), then $x' \in B(x, r(1 - \frac{1}{n}))$.

Consider an arbitrary subcover of \mathcal{S} , i.e.,

$$\mathcal{S}' = \left\{ B\left(x, r\left(1 - \frac{1}{n}\right)\right) \right\}_{n \in \mathcal{N}}$$

with $\#\mathcal{N} = N \in \mathbb{N}$. Define $n^* = \min\{n \in \mathcal{N}\}$. Then $\cup_{n \in \mathcal{N}} B(x, r(1 - \frac{1}{n})) = B(x, r(1 - \frac{1}{n^*}))$, and if $d(x', x) \in (r(1 - \frac{1}{n^*}), r)$, then $x' \in B(x, r)$ and $x' \notin \cup_{n \in \mathcal{N}} B(x, r(1 - \frac{1}{n}))$.

17.

1st proof.

We have to show that $f(A \cup B) \subseteq f(A) \cup f(B)$ and $f(A) \cup f(B) \subseteq f(A \cup B)$.

To prove the first inclusion, take $y \in f(A \cup B)$; then $\exists x \in A \cup B$ such that $f(x) = y$. Then either $x \in A$ or $x \in B$ that implies $f(x) = y \in A$ or $f(x) = y \in B$. In both case $y \in f(A) \cup f(B)$.

We now show the opposite inclusion. Let $y \in f(A) \cup f(B)$, then $y \in f(A)$ or $y \in f(B)$, but $y \in f(A)$ implies that $\exists x \in A$ such that $f(x) = y$. The same implication for $y \in f(B)$. As results, $y = f(x)$ in either case with $x \in A \cup B$ i.e. $y \in f(A \cup B)$.

2nd proof.

$$\begin{aligned} y \in f(A \cup B) &\Leftrightarrow \\ &\Leftrightarrow \exists x \in A \cup B \text{ such that } f(x) = y \\ &\Leftrightarrow (\exists x \in A \text{ such that } f(x) = y) \vee (\exists x \in B \text{ such that } f(x) = y) \\ &\Leftrightarrow (y \in f(A)) \vee (y \in f(B)) \\ &\Leftrightarrow y \in f(A) \cup f(B) \end{aligned}$$

18.:

First proof. Take $f = \sin$, $A = [-2\pi, 0]$, $B = [0, 2\pi]$.

Second proof. Consider

$$f : \{0, 1\} \rightarrow \mathbb{R}, \quad x \mapsto 1$$

Then take $A = \{0\}$ and $B = \{1\}$. Then $A \cap B = \emptyset$, so $f(A \cap B) = \emptyset$. But as $f(A) = f(B) = \{1\}$, we have that $f(A) \cap f(B) = \{1\} \neq \emptyset$.

19.

Take $c \in \mathbb{R}$ and define the following function

$$f : X \rightarrow Y, \quad f(x) = c.$$

It suffices to show that the preimage of every open subset of the domain is open in the codomain. The inverse image of any open set K is either X (if $c \in K$) or \emptyset (if $c \notin K$), which are both open sets.

20.

a.

i. \mathbb{R}_+^n is not bounded, then by Proposition 444 it is not compact.

ii. $Cl B(x, r)$ is compact.

From Proposition 444, it suffices to show that the set is closed and bounded.

$Cl B(x, r)$ is closed from Proposition 401.

$Cl B(x, r)$ is bounded because $Cl B(x, r) = \{y \in \mathbb{R}^n : d(x, y) \leq r\} \subseteq B(x, 2r)$.

Let's show in detail the equality.

i. $Cl B(x, r) \subseteq C$.

The function $d_x : \mathbb{R}^n \rightarrow \mathbb{R}$, $d_x(y) = d(x, y) := \left(\sum_{i=1}^n (x_i - y_i)^2\right)^{\frac{1}{2}}$ is continuous. Therefore, $C = d_x^{-1}([0, r])$ is closed. Since $B(x, r) \subseteq C$, by definition of closure, the desired result follows.

ii. $Cl B(x, r) \supseteq C$.

From Corollary 485, it suffices to show that $Ad B(x, r) \supseteq C$. If $d(y, x) < r$, we are done. Suppose that $d(y, x) = r$. We want to show that for every $\varepsilon > 0$, we have that $B(x, r) \cap B(y, \varepsilon) \neq \emptyset$. If $\varepsilon > r$, then $x \in B(x, r) \cap B(y, \varepsilon)$. Now take, $\varepsilon \leq r$. It is enough to take a point "very close to y inside $B(x, r)$ ". For example, we can verify that $z \in B(x, r) \cap B(y, \varepsilon)$, where $z = x + (1 - \frac{\varepsilon}{2r})(y - x)$. Indeed,

$$d(x, z) = \left(1 - \frac{\varepsilon}{2r}\right)d(y, x) = \left(1 - \frac{\varepsilon}{2r}\right)r = r - \frac{\varepsilon}{2} < r,$$

and

$$d(y, z) = \frac{\varepsilon}{2r}d(y, x) = \frac{\varepsilon}{2r}r = \frac{\varepsilon}{2} < \varepsilon.$$

c.

See solution to Exercise 5, where it was shown that S is not closed and therefore using Proposition 444, we can conclude S is not compact.

21.

Observe that given for any $j \in \{1, \dots, m\}$, the continuous functions $g_j : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g = (g_j)_{j=1}^m$, we can define

$$C := \{x \in \mathbb{R}^n : g(x) \geq 0\}.$$

Then C is closed, because of the following argument:

$C = \cap_{j=1}^m g_j^{-1}([0, +\infty))$; since g_j is continuous, and $[0, +\infty)$ is closed, then $g_j^{-1}([0, +\infty))$ is closed in \mathbb{R}^n ; then C is closed because intersection of closed sets.

22.

The set is closed, because $X = f^{-1}(\{0\})$.

The set is not compact: take f as the constant function.

23.

Let $V = B_Y(1, \varepsilon)$, be an open ball around the value 1 of the codomain, with $\varepsilon < 1$. $f^{-1}(V) = \{0\} \cup B_X(1, \varepsilon)$ is the union of an open set and a closed set, so is neither open nor closed.

24.

To apply the Extreme Value Theorem, we first have to check if the function to be maximized is continuous. Clearly, the function $\sum_{i=1}^n x_i$ is continuous as is the sum of affine functions. Therefore, to check for the existence of solutions for the problems we only have to check for the compactness of the restrictions.

The first set is closed, because it is the inverse image of the closed set $[0, 1]$ via the continuous function $|||$. The first set is bounded as well by definition. Therefore the set is compact and the function is continuous, we can apply Extreme Value theorem. The second set is not closed, therefore it is not compact and Extreme Value theorem can not be applied. The third set is unbounded, and therefore it is not compact and the Extreme Value theorem can not be applied.

25.

$[\Leftarrow]$

Obvious.

$[\Rightarrow]$

We want to show that $l \neq 0 \Rightarrow l$ is not bounded, i.e., $\forall M \in \mathbb{R}_{++}, \exists x \in E$ such that $\|l(x)\|_F > M$.

Since $l \neq 0$, $\exists y \in E \setminus \{0\}$ such that $l(y) \neq 0$. Define $x = \frac{2M}{\|l(x)\|_F} y$. Then

$$\|l(y)\|_F = \left\| l \left(\frac{2M}{\|l(x)\|_F} x \right) \right\|_F = \frac{2M}{\|l(x)\|_F} \|l(x)\|_F = 2M > M,$$

as desired.

26.

$a \Rightarrow b$.

Take an arbitrary $\alpha \in \mathbb{R}$; if $\{x \in X : f(x) < \alpha\} = f^{-1}((-\infty, \alpha)) = \emptyset$, we are done. Otherwise, take $x_0 \in f^{-1}((-\infty, \alpha))$. Then, $\alpha - f(x_0) := \varepsilon > 0$ and by definition of upper semicontinuity, we have

$$\exists \delta > 0 \text{ such that } d(x - x_0) < \delta \Rightarrow f(x) < f(x_0) + \varepsilon = f(x_0) + \alpha - f(x_0) = \alpha,$$

i.e., $B(x_0, \delta) \subseteq f^{-1}((-\infty, \alpha))$, i.e., the desired result.

$b \Leftrightarrow c$.

$$\{x \in X : f(x) < \alpha\} = X \setminus \{x \in X : f(x) \geq \alpha\}.$$

$b \Rightarrow a$.

We want to show that

$$\forall x_0 \in X, \forall \varepsilon > 0, \exists \delta > 0 \text{ such that } B(x_0, \delta) \subseteq f^{-1}((-\infty, f(x_0) + \varepsilon)).$$

But by assumption $f^{-1}((-\infty, f(x_0) + \varepsilon))$ is an open set and contains x_0 and therefore the desired result follows.

27.

Take $y \in \{x\} + A$. Then there exists $a \in A$ such that $y = x + a$ and since A is open there exists $\varepsilon > 0$ such that

$$a \in B(a, \varepsilon) \subseteq A. \quad (21.10)$$

We want to show that

i. $\{x\} + B(a, \varepsilon)$ is an open ball centered at $y = x + a$, i.e., $\{x\} + B(a, \varepsilon) = B(x + a, \varepsilon)$, and

ii. $B(x + a, \varepsilon) \subseteq \{x\} + A$.

i.

$[\subseteq]$ $y \in \{x\} + B(a, \varepsilon) \Leftrightarrow \exists z \in X$ such that $d(z, a) < \varepsilon$ and $y = x + z \Rightarrow d(y, x + a) = d(x + z, x + a) \stackrel{Hint}{=} d(z, a) < \varepsilon \Rightarrow y \in B(x + a, \varepsilon)$.

$[\supseteq]$ $y \in B(x + a, \varepsilon) \Leftrightarrow d(y, x + a) < \varepsilon$. Now since $y = x + (y - x)$ and $d(y - x, a) = \|y - x - a\| = \|y - (x + a)\| = d(y, x + a) < \varepsilon$, we get the desired conclusion.

ii.

$y \in B(x + a, \varepsilon) \Leftrightarrow \|y - (x + a)\| < \varepsilon$. Since $y = x + (y - x)$ and $\|(y - x) - a\| < \varepsilon$, i.e., $y - x \in B(a, \varepsilon) \stackrel{(21.10)}{\subseteq} A$, we get the desired result.

28.

By assumption, for any $i \in \{1, 2\}$ and for any $\{x_i^n\}_n \subseteq K_i$, there exists $x_i \in K_i$ such that, up to a subsequence $x_i^n \rightarrow x_i$. Take $\{y^n\} \subseteq K_1 + K_2 = K$. Then $\forall n$, $y^n = x_1^n + x_2^n$ with $x_i^n \in K_i$, $i = 1, 2$. Thus taking converging subsequences of $(x_i^n)_n$, $i \in \{1, 2\}$, we get $y^n \rightarrow x_1 + x_2 \in K$ as desired.

29.

a. We want to show that $\forall x_1, x_2 \in E$, $f(x_1) = f(x_2) \Rightarrow x_1 = x_2$. Indeed

$$0 = d(f(x_1), f(x_2)) = d(x_1, x_2) \Rightarrow x_1 = x_2.$$

b. It follows from a.

c. We want to show that $\forall x_0 \in E, \forall \varepsilon > 0, \exists \delta > 0$ such that

$$d_1(x, x_0) < \delta \Rightarrow d_2(f(x), f(x_0)) < \varepsilon.$$

Take $\delta = \varepsilon$. Then,

$$d_1(x, x_0) < \varepsilon \Rightarrow d_2(f(x), f(x_0)) < \varepsilon.$$

21.2.2 Correspondences

1.

Since u is a continuous function, from the Extreme Value Theorem, we are left with showing that for every (p, w) , $\beta(p, w)$ is non empty and compact, i.e., β is non empty valued and compact valued.

$$x = \left(\frac{w}{Cp^c} \right)_{c=1}^C \in \beta(p, w).$$

$\beta(p, w)$ is closed because it is the intersection of the inverse image of two closed sets via continuous functions.

$\beta(p, w)$ is bounded below by zero.

$\beta(p, w)$ is bounded above because for every c , $x^c \leq \frac{w - \sum_{c' \neq c} p^{c'} x^{c'}}{p^c} \leq \frac{w}{p^c}$, where the first inequality comes from the fact that $px \leq w$, and the second inequality from the fact that $p \in \mathbb{R}_{++}^C$ and $x \in \mathbb{R}_+^C$.

2.

(a) Consider $x', x'' \in \xi(p, w)$. We want to show that $\forall \lambda \in [0, 1]$, $x^\lambda := (1 - \lambda)x' + \lambda x'' \in \xi(p, w)$. Observe that $u(x') = u(x'') := u^*$. From the quasiconcavity of u , we have $u(x^\lambda) \geq u^*$. We are therefore left with showing that $x^\lambda \in \beta(p, w)$, i.e., β is convex valued. To see that, simply, observe that $px^\lambda = (1 - \lambda)px' + \lambda px'' \leq (1 - \lambda)w + \lambda w = w$.

(b) Assume otherwise. Following exactly the same argument as above we have $x', x'' \in \xi(p, w)$, and $px^\lambda \leq w$. Since u is strictly quasiconcave, we also have that $u(x^\lambda) > u(x') = u(x'') := u^*$, which contradicts the fact that $x', x'' \in \xi(p, w)$.

3.

We want to show that for every (p, w) the following is true. For every sequence $\{(p_n, w_n)\}_n \subseteq \mathbb{R}_{++}^C \times \mathbb{R}_{++}$ such that

$$(p_n, w_n) \rightarrow (p, w), \quad x_n \in \beta(p_n, w_n), \quad x_n \rightarrow x,$$

it is the case that $x \in \beta(p, w)$.

Since $x_n \in \beta(p_n, w_n)$, we have that $p_n x_n \leq w_n$. Taking limits of both sides, we get $px \leq w$, i.e., $x \in \beta(p, w)$.

4.

(a) We want to show that $\forall y', y'' \in y(p)$, $\forall \lambda \in [0, 1]$, it is the case that $y^\lambda := (1 - \lambda)y' + \lambda y'' \in y(p)$, i.e., $y^\lambda \in Y$ and $\forall y \in Y$, $py^\lambda \geq py$.

$y^\lambda \in Y$ simply because Y is convex.

$$py^\lambda := (1 - \lambda)py' + \lambda py'' \stackrel{y', y'' \in y(p)}{\geq} (1 - \lambda)py + \lambda py = py.$$


(b) Suppose not; then $\exists y', y'' \in Y$ such that $y' \neq y''$ and such that

$$\forall y \in Y, \quad py' = py'' > py \quad (1).$$

Since Y is strictly convex, $\forall \lambda \in (0, 1)$, $y^\lambda := (1 - \lambda)y' + \lambda y'' \in \text{Int } Y$. Then, $\exists \varepsilon > 0$ such that $B(y^\lambda, \varepsilon) \subseteq Y$. Consider $y^* := y^\lambda + \frac{\varepsilon}{2C} \mathbf{1}$, where $\mathbf{1} := (1, \dots, 1) \in \mathbb{R}^C$. $d(y^*, y^\lambda) = \sqrt{\sum_{c=1}^C \left(\frac{\varepsilon}{2C}\right)^2} = \frac{\varepsilon}{2\sqrt{C}}$. Then, $y^* \in B(y^\lambda, \varepsilon) \subseteq Y$ and, since $p \gg 0$, we have that $py^* > py^\lambda = py' = py''$, contradicting (1).

5.


This exercise is taken from Beavis and Dobbs (1990), pages 74-78.



For every $x \in [0, 2]$, both $\phi_1(x)$ and $\phi_2(x)$ are closed, bounded intervals and therefore convex and compact sets. Clearly ϕ_1 is closed and ϕ_2 is not closed.

ϕ_1 and ϕ_2 are clearly UHC and LHC for $x \neq 1$. Using the definitions, it is easy to see that for $x = 1$, ϕ_1 is UHC, and not LHC and ϕ_2 is LHC and not UHC.

6.



For every $x > 0$, ϕ is a continuous function. Therefore, for those values of x , ϕ is both UHC and LHC.

ϕ is UHC in 0. For every neighborhood of $[-1, 1]$ and for any neighborhood of $\{0\}$ in \mathbb{R}^+ , $\phi(x) \subseteq [-1, 1]$.

ϕ is not LHC in 0. Take the open set $V = (\frac{1}{2}, \frac{3}{2})$; we want to show that $\forall \varepsilon > 0 \exists z^* \in (0, \varepsilon)$ such that $\phi(z^*) \not\subseteq (\frac{1}{2}, \frac{3}{2})$. Take $n \in \mathbb{N}$ such that $\frac{1}{n} < \varepsilon$ and $z^* = \frac{1}{n\pi}$. Then $0 < z^* < \varepsilon$ and $\sin z^* = \sin n\pi = 0 \notin (\frac{1}{2}, \frac{3}{2})$.

Since ϕ is UHC and closed valued, from Proposition 16 is closed.

7.

ϕ is not closed. Take $x_n = \frac{\sqrt{2}}{2n} \in [0, 1]$ for every $n \in \mathbb{N}$. Observe that $x_n \rightarrow 0$. For every $n \in \mathbb{N}$, $y_n = -1 \in \phi(x_n)$ and $y_n \rightarrow -1$. But $-1 \in \phi(0) = [0, 1]$.


ϕ is not UHC. Take $x = 0$ and a neighborhood $V = (-\frac{1}{2}, \frac{3}{2})$ of $\phi(0) = [0, 1]$. Then $\forall \varepsilon > 0, \exists x^* \in (0, \varepsilon) \setminus \mathbb{Q}$. Therefore, $\phi(x^*) = [-1, 0] \not\subseteq V$.

ϕ is not LHC. Take $x = 0$ and the open set $V = (\frac{1}{2}, \frac{3}{2})$. Then $\phi(0) \cap (\frac{1}{2}, \frac{3}{2}) = [0, 1] \cap (\frac{1}{2}, \frac{3}{2}) = (\frac{1}{2}, 1] \neq \emptyset$. But, as above, $\forall \varepsilon > 0, \exists x^* \in (0, \varepsilon) \setminus \mathbb{Q}$. Then $\phi(x^*) \cap V = [-1, 0] \cap (\frac{1}{2}, \frac{3}{2}) = \emptyset$.

8.

(This exercise is taken from Klein, E. (1973), *Mathematical Methods in Theoretical Economics*, Academic Press, New York, NY, page 119).

Observe that $\phi_3(x) = [x^2 - 2, x^2 - 1]$.



$\phi_1([0, 3]) = \{(x, y) \in \mathbb{R}^2 : x \geq 0, x \leq 3, y \geq x^2 - 2, y \leq x^2\}$. $\phi_1([0, 3])$ is defined in terms of weak inequalities and continuous functions and it is closed and therefore ϕ_1 is closed. Similar argument applies to ϕ_2 and ϕ_3 .

Since $[-10, 10]$ is a compact set such that $\phi_1([0, 3]) \subseteq [-10, 10]$, from Proposition 17, ϕ_1 is UHC. Similar argument applies to ϕ_2 and ϕ_3 .

ϕ_1 is LHC. Take an arbitrary $\bar{x} \in [0, 3]$ and an open set V with non-empty intersection with $\phi_1(\bar{x}) = [\bar{x}^2 - 2, \bar{x}^2]$. To fix ideas, take $V = [\varepsilon, \bar{x}^2 + \varepsilon]$, with $\varepsilon \in (0, \bar{x}^2)$. Then, take $U = (\sqrt{\varepsilon}, \sqrt{\bar{x}^2 + \varepsilon})$. Then for every $x \in U$, $\{x^2\} \subseteq V \cap \phi_1(x)$.

Similar argument applies to ϕ_2 and ϕ_3 .

21.3 Differential Calculus in Euclidean Spaces

1 .

The partial derivative of f with respect to the first coordinate at the point (x_0, y_0) , is - if it exists and is finite -

$$\begin{aligned}\lim_{x \rightarrow x_0} \frac{f(x, y_0) - f(x_0, y_0)}{x - x_0} &= \frac{f(x, y_0) - f(x_0, y_0)}{x - x_0} \\ &= \frac{2x^2 - xy_0 + y_0^2 - (2x_0^2 - x_0y_0 + y_0^2)}{x - x_0} \\ &= \frac{2x^2 - 2x_0^2 - (xy_0 - x_0y_0)}{x - x_0} \\ &= 2(x + x_0) - y_0\end{aligned}$$

Then

$$D_1f(x_0, y_0) = 4x_0 - y_0$$

The partial derivative of f with respect to the second coordinate at the point (x_0, y_0) , is - if it exists and is finite -

$$\begin{aligned}\lim_{y \rightarrow y_0} \frac{f(x_0, y) - f(x_0, y_0)}{y - y_0} &= \frac{f(x_0, y) - f(x_0, y_0)}{y - y_0} \\ &= \frac{2x_0^2 - x_0y + y^2 - (2x_0^2 - x_0y_0 + y_0^2)}{y - y_0} \\ &= \frac{-x_0(y - y_0) + y^2 - y_0^2}{y - y_0} \\ &= -x_0 + (y + y_0) \\ D_2f(x_0, y_0) &= -x_0 + 2y_0.\end{aligned}$$

2 .

a.

The domain of f is $\mathbb{R} \setminus \{0\} \times \mathbb{R}$. As arctan is differentiable over the whole domain, we may compute the partial derivative over the whole domain of f at the point (x, y) - we omit from now on the superscript 0

$$\begin{aligned}D_1f(x, y) &= \arctan \frac{y}{x} + x \left(-\frac{y}{x^2}\right) \frac{1}{1 + \left(\frac{y}{x}\right)^2} \\ &= \arctan \frac{y}{x} - y \frac{1}{1 + \left(\frac{y}{x}\right)^2} \\ D_2f(x, y) &= x \frac{1}{x} \frac{1}{1 + \left(\frac{y}{x}\right)^2} = \frac{1}{1 + \left(\frac{y}{x}\right)^2}\end{aligned}$$

b.

The function is defined on $\mathbb{R}_{++} \times \mathbb{R}$. and

$$\forall (x, y) \in \mathbb{R}_{++} \times \mathbb{R}, \quad f(x, y) = e^{y \ln x}$$

Thus as exp and \ln are differentiable over their whole respective domain, we may compute the partial derivatives :

$$\begin{aligned}D_1f(x, y) &= \frac{y}{x} e^{y \ln x} = yx^{y-1} \\ D_2f(x, y) &= \ln(x) e^{y \ln x} = \ln(x) x^y\end{aligned}$$

c.

$$f(x, y) = (\sin(x + y))^{\sqrt{x+y}} = e^{\sqrt{x+y} \ln[\sin(x+y)]} \text{ in } (0, 3).$$

We check that $\sin(0 + 3) > 0$ so that the point belongs to the domain of the function. Both partial derivatives in (x, y) have the same expression since f is symmetric with respect to x and y .

$$D_1 f(x, y) = D_2 f(x, y) = \left[\frac{1}{2\sqrt{x+y}} \ln[\sin(x+y)] + \sqrt{x+y} \cot(x+y) \right] (\sin(x+y))^{\sqrt{x+y}},$$

and

$$D_1 f(0, 3) = D_2 f(0, 3) = \left[\frac{1}{2\sqrt{3}} \ln[\sin(3)] + \sqrt{3} \tan(3) \right] (\sin(3))^{\sqrt{3}}$$

3.

a.

$$D_x f(0, 0) = \lim_{h \rightarrow 0} \frac{f(h, 0) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{0}{h} = 0;$$

b.

$$D_y f(0, 0) = \lim_{k \rightarrow 0} \frac{f(0, k) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{0}{h} = 0;$$

c.

Consider the sequences $(x_n)_{n \in \mathbb{N}}$ and $(y_n)_{n \in \mathbb{N}}$ such that $\forall n \in \mathbb{N}, : x_n = y_n = \frac{1}{n}$. We have that $\lim_{n \rightarrow 0} (x_n, y_n) = 0_{\mathbb{R}^2}$, but

$$f(x_n, y_n) = \frac{\frac{1}{n^2}}{\frac{1}{n^2} + \frac{1}{n^2}}.$$

Then

$$\lim_{n \rightarrow 0} f(x_n, y_n) = \frac{1}{2} \neq f(0, 0) = 0$$

Thus, f is not continuous in $(0, 0)$.

4 .

$$\begin{aligned} f'((1, 1); (\alpha_1, \alpha_2)) &= \lim_{h \rightarrow 0} \frac{f(1 + h\alpha_1, 1 + h\alpha_2) - f(1, 1)}{h} = \\ \lim_{h \rightarrow 0} \frac{1}{h} \left[\frac{2 + h(\alpha_1 + \alpha_2)}{(1 + h\alpha_1)^2 + (1 + h\alpha_2)^2 + 1} - \frac{2}{3} \right] &= \dots \\ &= -\frac{\alpha_1 + \alpha_2}{9} \end{aligned}$$

5 .

1st answer.

We will show the existence of a linear function $T_{(x_0, y_0)}(x, y) = a(x - x_0) + b(y - y_0)$ such that the definition of differential is satisfied. After substituting, we want to show that

$$\lim_{(x, y) \rightarrow (x_0, y_0)} \frac{|x^2 - y^2 + xy - x_0^2 + y_0^2 - x_0 y_0 - a(x - x_0) - b(y - y_0)|}{\sqrt{(x - x_0)^2 + (y - y_0)^2}} = 0.$$

Manipulate the numerator of the above to get $NUM = |(x - x_0)(x + x_0 - a + y) - (y - y_0)(y + y_0 + b - x_0)|$. Now the ratio R whose limit we are interested to obtain satisfies

$$\begin{aligned} 0 \leq R &\leq \frac{|x - x_0||x + x_0 - a + y|}{\sqrt{(x - x_0)^2 + (y - y_0)^2}} + \frac{|y - y_0||y + y_0 + b - x_0|}{\sqrt{(x - x_0)^2 + (y - y_0)^2}} \\ &\leq |x - x_0||x + x_0 - a + y| + |y - y_0||y + y_0 + b - x_0|. \end{aligned}$$

For $a = 2x_0 + y_0$ and $b = x_0 - 2y_0$ we get the limit of R equal zero as required.

2nd answer.

As a polynomial function, we know that f is $C^1(\mathbb{R}^2)$ so we can “guess” that the derivative at the point (x, y) is the following linear application:

$$\begin{aligned} df_{(x,y)} : \mathbb{R}^2 &\rightarrow \mathbb{R} \\ u &\mapsto (2x + y, -2y + x) \cdot \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \end{aligned}$$

In order to comply with the definition, we have to prove that

$$\lim_{u \rightarrow 0} \frac{f((x, y) + (u_1, u_2)) - f((x, y)) - df_{(x,y)}(u)}{\|u\|} = 0$$

$$\begin{aligned} f((x, y) + (u_1, u_2)) - f((x, y)) - df_{(x,y)}(u) &= (x + u_1)^2 - (y + u_2)^2 + (x + u_1)(y + u_2) \\ &\quad - x^2 + y^2 - xy - (2x + y, -2y + x) \cdot \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} \\ &= 2xu_1 + u_1^2 - 2yu_2 - u_2^2 + xu_2 + yu_1 + u_1u_2 \\ &\quad - (2x + y)u_1 - (-2y + x)u_2 \\ &= u_1^2 - u_2^2 + u_1u_2 \\ &= O(\|u\|^2) \end{aligned}$$

Or, we know that $\lim_{u \rightarrow 0} \frac{O(\|u\|^2)}{\|u\|} = 0$. Then we have proven that the candidate was indeed the derivative of f at point (x, y) .

6 .

a) given $x_0 \in \mathbb{R}^n$ we need to find $T_{x_0} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ linear and E_{x_0} with $\lim_{v \rightarrow 0} E_{x_0}(v) = 0$. Take $T_{x_0} = l$ and $E_{x_0} \equiv 0$. Then, the desired result follows.

b) projection is linear, so by a) is differentiable.

7.

From the definition of continuity, we want to show that $\forall x_0 \in \mathbb{R}^n, \forall \varepsilon > 0 \exists \delta > 0$ such that $\|x - x_0\| < \delta \Rightarrow \|l(x) - l(x_0)\| < \varepsilon$. Defined $[l] = A$, we have that

$$\begin{aligned} \|l(x) - l(x_0)\| &= \|A \cdot x - x_0\| = \\ &= \|R^1(A) \cdot (x - x_0), \dots, R^m(A) \cdot (x - x_0)\| \stackrel{(1)}{\leq} \sum_{i=1}^m |R^i(A) \cdot (x - x_0)| \stackrel{(2)}{\leq} \\ &\leq \sum_{i=1}^m \|R^i(A)\| \cdot \|x - x_0\| \leq m \cdot (\max_{i \in \{1, \dots, m\}} \{\|R^i(A)\|\}) \cdot \|x - x_0\|, \end{aligned} \quad (21.11)$$

where (1) follows from Remark 56 and (2) from Proposition 53.4, i.e., Cauchy-Schwarz inequality. Take

$$\delta = \frac{\varepsilon}{m \cdot (\max_{i \in \{1, \dots, m\}} \{\|R^i(A)\|\})}.$$

Then we have that $\|x - x_0\| < \delta$ implies that $\|x - x_0\| \cdot m \cdot (\max_{i \in \{1, \dots, m\}} \{\|R^i(A)\|\}) < \varepsilon$, and from (21.11), $\|l(x) - l(x_0)\| < \varepsilon$, as desired.

8 .

$$Df(x, y) = \begin{bmatrix} \cos x \cos y & -\sin x \sin y \\ \cos x \sin y & \sin x \cos y \\ -\sin x \cos y & -\cos x \sin y \end{bmatrix}.$$

9 .

$$Df(x, y, z) = \begin{bmatrix} g'(x)h(z) & 0 & g(x)h'(z) \\ g'(h(x))h'(x)/y & -g(h(x))/y^2 & 0 \\ \exp(xg(h(x))((g(h(x)) + g'(h(x))h'(x)x)) & 0 & 0 \end{bmatrix}.$$

10 .

a.

f is differentiable over its domain since $x \mapsto \log x$ is differentiable over \mathbb{R}_{++} . Then from Proposition 610, we know that

$$[df_x] = Df(x) = \left[\frac{1}{3x_1}, \quad \frac{1}{6x_2}, \quad \frac{1}{2x_3} \right]$$

Then

$$[df_{x_0}] = \left[\frac{1}{3} \quad \frac{1}{6} \quad \frac{1}{4} \right]$$

By application of Remark 611, we have that

$$f'(x_0, u) = Df(x_0).u = \frac{1}{\sqrt{3}} \begin{bmatrix} \frac{1}{3} & \frac{1}{6} & \frac{1}{4} \end{bmatrix} \cdot \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix} = \frac{\sqrt{3}}{4}$$

b.

As a polynomial expression, f is differentiable over its domain.

$$[df_x] = Df(x) = [2x_1 - 2x_2, \quad 4x_2 - 2x_1 - 6x_3, \quad -2x_3 - 6x_2]$$

Then

$$[df_{x_0}] = [2 \quad 4 \quad 2]$$

and

$$f'(x_0, u) = Df(x_0).u = [2 \quad 4 \quad 2] \cdot \begin{pmatrix} -\frac{1}{\sqrt{2}} \\ 0 \\ \frac{1}{\sqrt{2}} \end{pmatrix} = 0$$

c.

$$[df_x] = Df(x) = [e^{x_1x_2} + x_2x_1e^{x_1x_2}, \quad x_1^2e^{x_1x_2}]$$

Then

$$[df_{x_0}] = [1 \quad 0]$$

and

$$f'(x_0, u) = Df(x_0).u = [1, \quad 0] \cdot \begin{pmatrix} 2 \\ 3 \end{pmatrix} = 2$$

11 .

Then since f is in $C^\infty(\mathbb{R} \setminus \{0\})$, we know that f admits partial derivative *functions* and that these functions admit themselves partial derivatives in (x, y, z) . Since, the function is symmetric in its arguments, it is enough to compute explicitly $\frac{\partial^2 f(x, y, z)}{\partial x^2}$.

$$\frac{\partial f}{\partial x}(x, y, z) = -x(x^2 + y^2 + z^2)^{-\frac{3}{2}}$$

Then

$$\frac{\partial^2 f}{\partial x^2}(x, y, z) = -(x^2 + y^2 + z^2)^{-\frac{3}{2}} + 3x^2(x^2 + y^2 + z^2)^{-\frac{5}{2}}$$

Then $\forall (x, y, z) \in \mathbb{R}^3 \setminus \{0\}$,

$$\begin{aligned} \frac{\partial^2 f}{\partial x^2}(x, y, z) + \frac{\partial^2 f}{\partial y^2}(x, y, z) + \frac{\partial^2 f}{\partial z^2}(x, y, z) &= -3(x^2 + y^2 + z^2)^{-\frac{3}{2}} + (3x^2 + 3y^2 + 3z^2)(x^2 + y^2 + z^2)^{-\frac{5}{2}} \\ &= -3(x^2 + y^2 + z^2)^{-\frac{3}{2}} + 3(x^2 + y^2 + z^2)^{-\frac{3}{2}} \\ &= 0 \end{aligned}$$

12 .

$g, h \in C^2(\mathbb{R}, \mathbb{R}_{++})^2$

$$f : \mathbb{R}^3 \rightarrow \mathbb{R}^3, \quad f(x, y, z) = \left(\frac{g(x)}{h(z)}, \quad g(h(x)) + xy, \quad \ln(g(x) + h(x)) \right)$$

Since $Im(g) \subseteq \mathbb{R}_{++}$, $(x, y, z) \mapsto \frac{g(x)}{h(z)}$ is differentiable as the ratio of differentiable functions. And since $Im(g+h) \subseteq \mathbb{R}_{++}$ and that \ln is differentiable over \mathbb{R}_{++} , $x \mapsto \ln(g(x) + h(x))$ is differentiable by proposition 619. Then

$$Df(x, y, z) = \begin{bmatrix} \frac{g'(x)}{h(z)} & 0 & -h'(z) \frac{g(x)}{h(z)^2} \\ h'(x)g'(h(x)) + y & x & 0 \\ \frac{g'(x)+h'(x)}{g(x)+h(x)} & 0 & 0 \end{bmatrix}$$

13 .

a.

Since

$$h(x) = \begin{pmatrix} e^x + g(x) \\ e^{g(x)} + x \end{pmatrix},$$

then

$$[dh_x] = Dh(x) = \begin{pmatrix} e^x + g'(x) \\ g'(x)e^{g(x)} + 1 \end{pmatrix},$$

$$[dh_0] = Dh(0) = \begin{pmatrix} 1 + g'(0) \\ g'(0)e^{g(0)} + 1 \end{pmatrix},$$

b.

Let us define $l: \mathbb{R} \rightarrow \mathbb{R}^2$, $l(x) = (x, g(x))$. Then $h = f \circ l$. As l is differentiable on \mathbb{R} and f is differentiable on \mathbb{R}^2 we may apply the "chain rule".

$$dh_0 = df_{l(0)} \circ dl_0$$

$$[dl_x] = Dl(x) = \begin{bmatrix} 1 \\ g'(x) \end{bmatrix}$$

$$[df_{(x,y)}] = Df(x, y) = \begin{bmatrix} e^x & 1 \\ 1 & e^y \end{bmatrix}$$

Then

$$[dh_0] = \begin{bmatrix} e^0 & 1 \\ 1 & e^{g(0)} \end{bmatrix} \begin{bmatrix} 1 \\ g'(0) \end{bmatrix} = \begin{bmatrix} 1 + g'(0) \\ 1 + g'(0)e^{g(0)} \end{bmatrix}$$

14 .

$$D_x(b \circ a)(x) = D_y b(y)|_{y=a(x)} \cdot D_x a(x).$$

$$D_y b(y) = \begin{bmatrix} D_{y_1} g(y) & D_{y_2} g(y) & D_{y_3} g(y) \\ D_{y_1} f(y) & D_{y_2} f(y) & D_{y_3} f(y) \end{bmatrix}_{|y=a(x)}$$

$$D_x a(x) = \begin{bmatrix} D_{x_1} f(x) & D_{x_2} f(x) & D_{x_3} f(x) \\ D_{x_1} g(x) & D_{x_2} g(x) & D_{x_3} g(x) \\ 1 & 0 & 0 \end{bmatrix}$$

$$D_x(b \circ a)(x) =$$

$$\begin{aligned} & \begin{bmatrix} D_{y_1} g(f(x), g(x), x_1) & D_{y_2} g(f(x), g(x), x_1) & D_{y_3} g(f(x), g(x), x_1) \\ D_{y_1} f(f(x), g(x), x_1) & D_{y_2} f(f(x), g(x), x_1) & D_{y_3} f(f(x), g(x), x_1) \end{bmatrix} \begin{bmatrix} D_{x_1} f(x) & D_{x_2} f(x) & D_{x_3} f(x) \\ D_{x_1} g(x) & D_{x_2} g(x) & D_{x_3} g(x) \\ 1 & 0 & 0 \end{bmatrix} = \\ & = \begin{bmatrix} D_{y_1} g & D_{y_2} g & D_{y_3} g \\ D_{y_1} f & D_{y_2} f & D_{y_3} f \end{bmatrix} \begin{bmatrix} D_{x_1} f & D_{x_2} f & D_{x_3} f \\ D_{x_1} g & D_{x_2} g & D_{x_3} g \\ 1 & 0 & 0 \end{bmatrix} = \end{aligned}$$

$$= \begin{bmatrix} D_{y_1}g \cdot D_{x_1}f + D_{y_2}gD_{x_1}g + D_{y_3}g & D_{y_1}g \cdot D_{x_2}f + D_{y_2}g \cdot D_{x_2}g & D_{y_1}g \cdot D_{x_3}f + D_{y_2}g \cdot D_{x_3}g \\ D_{y_1}f \cdot D_{x_1}f + D_{y_2}f \cdot D_{x_1}g + D_{y_3}f & D_{y_1}f \cdot D_{x_2}f + D_{y_2}f \cdot D_{x_2}g & D_{y_1}f \cdot D_{x_3}f + D_{y_2}f \cdot D_{x_3}g \end{bmatrix}.$$

15 .

By the sufficient condition of differentiability, it is enough to show that the function $f \in C^1$. Partial derivatives are $\frac{\partial f}{\partial x} = 2x + y$ and $\frac{\partial f}{\partial y} = -2y + x$ – both are indeed continuous, so f is differentiable.

16 .

i) $f \in C^1$ as $Df(x, y, z) = (1 + 4xy^2 + 3yz, 3y^2 + 4x^2y + 3z, 1 + 3xy + 3z^2)$ has continuous entries (everywhere, in particular around (x_0, y_0, z_0)).

ii) $f(x_0, y_0, z_0) = 0$ by direct calculation.

iii) $f'_z = \frac{\partial f}{\partial z}|_{(x_0, y_0, z_0)} = 7 \neq 0$, $f'_y = \frac{\partial f}{\partial y}|_{(x_0, y_0, z_0)} = 10 \neq 0$ and $f'_x = \frac{\partial f}{\partial x}|_{(x_0, y_0, z_0)} = 8 \neq 0$.

Therefore we can apply Implicit Function Theorem around $(x_0, y_0, z_0) = (1, 1, 1)$ to get

$$\frac{\partial x}{\partial z} = -\frac{f'_z}{f'_x} = -7/8,$$

$$\frac{\partial y}{\partial z} = -\frac{f'_z}{f'_y} = -7/10.$$

17 .

a)

$$Df = \begin{bmatrix} 2x_1 & -2x_2 & 2 & 3 \\ x_2 & x_1 & 1 & -1 \end{bmatrix}$$

and each entry of Df is continuous; then f is C^1 . $\det D_x f(x, t) = 2x_1^2 + 2x_2^2 \neq 0$ except for $x_1 = x_2 = 0$. Finally,

$$Dg(t) = -\begin{bmatrix} 2x_1 & -2x_2 \\ x_2 & x_1 \end{bmatrix}^{-1} \begin{bmatrix} 2 & 3 \\ 1 & -1 \end{bmatrix} = -\frac{1}{2x_1^2 + 2x_2^2} \begin{bmatrix} 2x_1 + 2x_2 & 3x_1 - 2x_2 \\ -2x_2 + 2x_1 & -2x_1 - 3x_2 \end{bmatrix}.$$

b)

$$Df = \begin{bmatrix} 2x_2 & 2x_1 & 1 & 2t_2 \\ 2x_1 & 2x_2 & 2t_1 - 2t_2 & -2t_1 + 2t_2 \end{bmatrix}$$

continuous, $\det D_x f(x, t) = 4x_2^2 - 4x_1^2 \neq 0$ except for $|x_1| = |x_2|$. Finally

$$\begin{aligned} Dg(t) &= -\begin{bmatrix} 2x_2 & 2x_1 \\ 2x_1 & 2x_2 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 2t_2 \\ 2t_1 - 2t_2 & -2t_1 + 2t_2 \end{bmatrix} = \\ &= -\frac{1}{4x_2^2 - 4x_1^2} \begin{bmatrix} -4x_1t_1 + 4x_1t_2 + 2x_2 & 4x_1t_1 - 4x_1t_2 + 4x_2t_2 \\ -2x_1 + 4x_2t_1 - 4x_2t_2 & -4x_1t_2 - 4x_2t_1 + 4x_2t_2 \end{bmatrix}. \end{aligned}$$

c)

$$Df = \begin{bmatrix} 2 & 3 & 2t_1 & -2t_2 \\ 1 & -1 & t_2 & t_1 \end{bmatrix}$$

continuous, $\det D_x f(x, t) = -5 \neq 0$ always. Finally

$$Dg(t) = -\begin{bmatrix} 2 & 3 \\ 1 & -1 \end{bmatrix}^{-1} \begin{bmatrix} 2t_1 & -2t_2 \\ t_2 & t_1 \end{bmatrix} = \frac{1}{5} \begin{bmatrix} -2t_1 - 3t_2 & 2t_2 - 3t_1 \\ -2t_1 + 2t_2 & 2t_2 + 2t_1 \end{bmatrix}.$$

18.

As an application of the Implicit Function Theorem, we have that

$$\frac{\partial z}{\partial x} = -\frac{\frac{\partial(z^3 - xz - y)}{\partial x}}{\frac{\partial(z^3 - xz - y)}{\partial z}} = -\frac{-z}{3z^2 - x}$$

if $3z^2 - x \neq 0$. Then,

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{\partial \left(\frac{z(x,y)}{3(z(x,y))^2 - x} \right)}{\partial y} = \frac{\frac{\partial z}{\partial y} (3z^2 - x) - 6 \frac{\partial z}{\partial y} \cdot z^2}{(3z^2 - x)^2}$$

Since,

$$\frac{\partial z}{\partial y} = - \frac{\frac{\partial(z^3 - xz - y)}{\partial y}}{\frac{\partial(z^3 - xz - y)}{\partial z}} = - \frac{-1}{3z^2 - x},$$

we get

$$\frac{\partial^2 z}{\partial x \partial y} = \frac{\frac{1}{3z^2 - x} (3z^2 - x) - 6 \frac{1}{3z^2 - x} \cdot z^2}{(3z^2 - x)^2} = \frac{-3z^2 - x}{(3z^2 - x)^3}$$

19.

As an application of the Implicit Function Theorem, we have that the Marginal Rate of Substitution in (x_0, y_0) is

$$\frac{dy}{dx} \big|_{(x,y)=(x_0,y_0)} = - \frac{\frac{\partial(u(x,y)-k)}{\partial x}}{\frac{\partial(u(x,y)-k)}{\partial y}} \big|_{(x,y)=(x_0,y_0)} < 0$$

$$\frac{d^2 y}{dx^2} = - \frac{\partial \left(\frac{D_x u(x,y(x))}{D_y u(x,y(x))} \right)}{\partial x} = - \frac{\left(\begin{smallmatrix} (-) \\ D_{xx} u \end{smallmatrix} + \begin{smallmatrix} (+) \\ D_{xy} u \end{smallmatrix} \begin{smallmatrix} (-) \\ \frac{dy}{dx} \end{smallmatrix} \right) D_y u - \left(\begin{smallmatrix} (+) \\ D_{xy} u \end{smallmatrix} + \begin{smallmatrix} (-) \\ D_{yy} u \end{smallmatrix} \begin{smallmatrix} (-) \\ \frac{dy}{dx} \end{smallmatrix} \right) D_x u}{(D_y u)^2} \bigg|_{(x,y)=(x_0,y_0)} > 0$$

and therefore the function $y(x)$ describing indifference curves is convex.

20.

Adapt the proof for the case of the derivative of the product of functions from \mathbb{R} to \mathbb{R} .

21.

Differentiate both sides of

$$f(ax_1, ax_2) = a^n f(x_1, x_2)$$

with respect to a and then replace a with 1.

21.4 Nonlinear Programming

1.

(a)

If $\beta = 0$, then $f(x) = \alpha$. The constant function is concave and therefore pseudo-concave, quasi-concave, not strictly concave.

If $\beta > 0$, $f'(x) = \alpha\beta x^{\beta-1}$, $f''(x) = \alpha\beta(\beta-1)x^{\beta-2}$.

$f''(x) \leq 0 \Leftrightarrow \alpha\beta(\beta-1) \geq 0 \stackrel{\alpha \geq 0, \beta \geq 0}{\Leftrightarrow} 0 \leq \beta \leq 1 \Leftrightarrow f \text{ concave} \Rightarrow f \text{ quasi-concave.}$

$f''(x) < 0 \Leftrightarrow (\alpha > 0 \text{ and } \beta \in (0, 1)) \Rightarrow f \text{ strictly concave.}$

(b)

The Hessian matrix of f is

$$D^2 f(x) = \begin{bmatrix} \alpha_1 \beta_1 (\beta_1 - 1) x^{\beta_1-2} & & 0 \\ & \ddots & \\ 0 & & \alpha_n \beta_n (\beta_n - 1) x^{\beta_n-2} \end{bmatrix}$$

$D^2 f(x)$ is negative semidefinite $\Leftrightarrow (\forall i, \beta_i \in [0, 1]) \Rightarrow f$ is concave.

$D^2 f(x)$ is negative definite $\Leftrightarrow (\forall i, \alpha_i > 0 \text{ and } \beta_i \in (0, 1)) \Rightarrow f$ is strictly concave.

The border Hessian matrix is

$$B(f(x)) = \begin{bmatrix} 0 & \alpha_1 \beta_1 x^{\beta_1-1} & - & \alpha_n \beta_n x^{\beta_n-1} \\ \alpha_1 \beta_1 x^{\beta_1-1} & \alpha_1 \beta_1 (\beta_1 - 1) x^{\beta_1-2} & & 0 \\ | & & \ddots & \\ \alpha_n \beta_n x^{\beta_n-1} & 0 & & \alpha_n \beta_n (\beta_n - 1) x^{\beta_n-2} \end{bmatrix}$$

The determinant of the significant leading principal minors are

$$\begin{aligned} \det \begin{bmatrix} 0 & \alpha_1 \beta_1 x^{\beta_1-1} \\ \alpha_1 \beta_1 x^{\beta_1-1} & \alpha_1 \beta_1 (\beta_1 - 1) x^{\beta_1-2} \end{bmatrix} &= -\alpha_1^2 \beta_1^2 (x^{\beta_1-1})^2 < 0 \\ \det \begin{bmatrix} 0 & \alpha_1 \beta_1 x^{\beta_1-1} & \alpha_2 \beta_2 x^{\beta_2-1} \\ \alpha_1 \beta_1 x^{\beta_1-1} & \alpha_1 \beta_1 (\beta_1 - 1) x^{\beta_1-2} & 0 \\ \alpha_2 \beta_2 x^{\beta_2-1} & 0 & \alpha_2 \beta_2 (\beta_2 - 1) x^{\beta_2-2} \end{bmatrix} &= \\ &= -[\alpha_1 \beta_1 x^{\beta_1-1} \alpha_1 \beta_1 x^{\beta_1-1} \alpha_2 \beta_2 (\beta_2 - 1) x^{\beta_2-2}] - [\alpha_2 \beta_2 x^{\beta_2-1} \alpha_2 \beta_2 x^{\beta_2-1} \alpha_1 \beta_1 (\beta_1 - 1) x^{\beta_1-2}] = \\ &= -(\alpha_1 \beta_1 \alpha_2 \beta_2 x^{\beta_1+\beta_2-4}) [\alpha_1 \beta_1 x^{\beta_2} (\beta_2 - 1) + \alpha_2 \beta_2 x^{\beta_1} (\beta_1 - 1)] = \\ &= -\alpha_1 \beta_1 \alpha_2 \beta_2 x^{\beta_1+\beta_2-4} [\alpha_1 \beta_1 x^{\beta_2} (\beta_2 - 1) + \alpha_2 \beta_2 x^{\beta_1} (\beta_1 - 1)] > 0 \end{aligned}$$

iff for $i = 1, 2$, $\alpha_i > 0$ and $\beta_i \in (0, 1)$.

(c)

If $\beta = 0$, then $f(x) = \min\{\alpha, -\gamma\} = 0$.

If $\beta > 0$, we have

The intersection of the two line has coordinates $x^* : \left(= \frac{\alpha+\gamma}{\beta}, \alpha \right)$.

f is clearly not strictly concave, because it is constant in a subset of its domain. Let's show it is concave and therefore pseudo-concave and quasi-concave.

Given $x', x'' \in X$, 3 cases are possible.

Case 1. $x', x'' \leq x^*$.

Case 2. $x', x'' \geq x^*$.

Case 3. $x' \leq x^*$ and $x'' \geq x^*$.

The most difficult case is case 3: we want to show that $(1 - \lambda) f(x') + \lambda f(x'') \leq f((1 - \lambda)x' + \lambda x'')$.

Then, we have

$$(1 - \lambda) f(x') + \lambda f(x'') = (1 - \lambda) \min\{\alpha, \beta x' - \gamma\} + \lambda \min\{\alpha, \beta x'' - \gamma\} = (1 - \lambda) (\beta x' - \gamma) + \lambda \alpha.$$

Since, by construction $\alpha \geq \beta x' - \gamma$,

$$(1 - \lambda) (\beta x' - \gamma) + \lambda \alpha \leq \alpha;$$

since, by construction $\alpha \leq \beta x'' - \gamma$,

$$(1 - \lambda) (\beta x' - \gamma) + \lambda \alpha \leq (1 - \lambda) (\beta x' - \gamma) + \lambda (\beta x'' - \gamma) = \beta [(1 - \lambda)x' + \lambda x''] - \gamma.$$

Then

$$(1 - \lambda) f(x') + \lambda f(x'') \leq \min\{\alpha, \beta [(1 - \lambda)x' + \lambda x''] - \gamma\} = f((1 - \lambda)x' + \lambda x''),$$

as desired.

2.


a.

i. Canonical form.

For given $\pi \in (0, 1)$, $a \in (0, +\infty)$,

$$\begin{array}{llll} \max_{(x,y) \in \mathbb{R}^2} & \pi \cdot u(x) + (1 - \pi) u(y) & \text{s.t.} & \begin{array}{ll} a - \frac{1}{2}x - y \geq 0 & \lambda_1 \\ 2a - 2x - y \geq 0 & \lambda_2 \\ x \geq 0 & \lambda_3 \\ y \geq 0 & \lambda_4 \end{array} \end{array}$$

$$\begin{cases} y = a - \frac{1}{2}x \\ y = 2a - 2x \end{cases}, \text{ solution is } \left[x = \frac{2}{3}a, y = \frac{2}{3}a \right]$$



ii. The set X and the functions f and g (X open and convex; f and g at least differentiable).

a. The domain of all function is \mathbb{R}^2 . Take $X = \mathbb{R}^2$ which is open and convex.

b. $Df(x, y) = (\pi \cdot u'(x), (1 - \pi) u'(y))$. The Hessian matrix is

$$\begin{bmatrix} \pi \cdot u''(x) & 0 \\ 0 & (1 - \pi) u''(y) \end{bmatrix}$$

Therefore, f and g are C^2 functions and f is strictly concave and the functions g^j are affine.

iii. Existence.

C is closed and bounded below by $(0, 0)$ and above by (a, a) :

$$y \leq a - \frac{1}{2}x \leq a$$

$$2x \leq 2a - y \leq 2a.$$

iv. Number of solutions.

The solution is unique because f is strictly concave and the functions g^j are affine and therefore concave.

v. Necessity of K-T conditions.

The functions g^j are affine and therefore concave.

$$x^{++} = \left(\frac{1}{2}a, \frac{1}{2}a \right)$$

$$\begin{aligned} a - \frac{1}{2} \cdot \frac{1}{2}a - \frac{1}{2}a &= \frac{1}{4}a > 0 \\ 2a - 2 \cdot \frac{1}{2}a - \frac{1}{2}a &= \frac{1}{2}a > 0 \\ \frac{1}{2}a &> 0 \\ \frac{1}{2}a &> 0 \end{aligned}$$

vi. Sufficiency of K-T conditions.

The objective function is strictly concave and the functions g^j are affine
vii. K-T conditions.

$$\mathcal{L}(x, y, \lambda_1, \dots, \lambda_4; \pi, a) = \pi \cdot u(x) + (1 - \pi) u(y) + \lambda_1 \left(a - \frac{1}{2}x - y \right) + \lambda_2 (2a - 2x - y) + \lambda_3 x + \lambda_4 y.$$

$$\begin{cases} \pi \cdot u'(x) - \frac{1}{2}\lambda_1 - 2\lambda_2 + \lambda_3 & = 0 \\ (1 - \pi) u'(y) - \lambda_1 - \lambda_2 + \lambda_4 & = 0 \\ \min \left\{ \lambda_1, a - \frac{1}{2}x - y \right\} & = 0 \\ \min \left\{ \lambda_2, 2a - 2x - y \right\} & = 0 \\ \min \left\{ \lambda_3, x \right\} & = 0 \\ \min \left\{ \lambda_4, y \right\} & = 0 \end{cases}$$

b. Inserting $(x, y, \lambda_1, \lambda_2, \lambda_3, \lambda_4) = (\frac{2}{3}a, \frac{2}{3}a, \lambda_1, 0, 0, 0)$, with $\lambda_1 > 0$, in the Kuhn-Tucker conditions we get:

$$\begin{cases} \pi \cdot u'(\frac{2}{3}a) - \frac{1}{2}\lambda_1 & = 0 \\ (1 - \pi) u'(\frac{2}{3}a) - \lambda_1 & = 0 \\ a - \frac{1}{2}\frac{2}{3}a - \frac{2}{3}a & = 0 \\ \min \left\{ 0, 2a - 2\frac{2}{3}a - \frac{2}{3}a \right\} & = 0 \\ \min \left\{ 0, \frac{2}{3}a \right\} & = 0 \\ \min \left\{ 0, \frac{2}{3}a \right\} & = 0 \end{cases}$$

and

$$\begin{cases} \lambda_1 & = 2\pi \cdot u'(\frac{2}{3}a) > 0 \\ \lambda_1 & = (1 - \pi) u'(\frac{2}{3}a) \geq 0 \\ a - \frac{1}{2}\frac{2}{3}a - \frac{2}{3}a & = 0 \\ \min \{0, 0\} & = 0 \\ \min \left\{ 0, \frac{2}{3}a \right\} & = 0 \\ \min \left\{ 0, \frac{2}{3}a \right\} & = 0 \end{cases}$$

Therefore, the proposed vector is a solution if

$$2\pi \cdot u'(\frac{2}{3}a) = (1 - \pi) u'(\frac{2}{3}a) > 0,$$

i.e.,

$$2\pi = 1 - \pi \quad \text{or} \quad \pi = \frac{1}{3} \quad \text{and for any } a \in \mathbb{R}_{++}.$$

c. If the first, third and fourth constraint hold with a strict inequality, and the multiplier associated with the second constraint is strictly positive, Kuhn-Tucker conditions become:

$$\begin{cases} \pi \cdot u'(x) - 2\lambda_2 & = 0 \\ (1 - \pi) u'(y) - \lambda_2 & = 0 \\ a - \frac{1}{2}x - y & > 0 \\ 2a - 2x - y & = 0 \\ x & > 0 \\ y & > 0 \end{cases}$$

$$\begin{cases} \pi \cdot u'(x) - 2\lambda_2 & = 0 \\ (1 - \pi) u'(y) - \lambda_2 & = 0 \\ 2a - 2x - y & = 0 \end{cases}$$

	x	y	λ_2	π	a
$\pi \cdot u'(x) - 2\lambda_2$	$\pi \cdot u''(x)$	0	-2	$u'(x)$	0
$(1 - \pi) u'(y) - \lambda_2$	0	$(1 - \pi) u''(y)$	-1	$-u'(y)$	0
$2a - 2x - y$	-2	-1	0	0	2

$$\begin{aligned}
& \det \begin{bmatrix} \pi \cdot u''(x) & 0 & -2 \\ 0 & (1-\pi)u''(y) & -1 \\ -2 & -1 & 0 \end{bmatrix} = \\
& = \pi u''(x) \det \begin{bmatrix} (1-\pi)u''(y) & -1 \\ -1 & 0 \end{bmatrix} - 2 \det \begin{bmatrix} 0 & -2 \\ (1-\pi)u''(y) & -1 \end{bmatrix} = \\
& = -\pi u''(x) - 4(1-\pi)u''(y) > 0 \\
D_{(\pi,a)}(x,y,\lambda_2) &= - \begin{bmatrix} \pi \cdot u''(x) & 0 & -2 \\ 0 & (1-\pi)u''(y) & -1 \\ -2 & -1 & 0 \end{bmatrix}^{-1} \begin{bmatrix} u'(x) & 0 \\ -u'(y) & 0 \\ 0 & 2 \end{bmatrix}
\end{aligned}$$

Using maple:

$$\begin{aligned}
& \begin{bmatrix} \pi \cdot u''(x) & 0 & -2 \\ 0 & (1-\pi)u''(y) & -1 \\ -2 & -1 & 0 \end{bmatrix}^{-1} = \\
& = \frac{1}{-\pi u''(x) - 4(1-\pi)u''(y)} \begin{bmatrix} -1 & 2 & 2u''(y) - 2\pi u''(y) \\ 2 & -4 & \pi u''(x) \\ 2u''(y) - 2\pi u''(y) & \pi u''(x) & \pi u''(x)u''(y) - \pi^2 u''(x)u''(y) \end{bmatrix} \\
D_{(\pi,a)}(x,y,\lambda_2) &= \\
& = -\frac{1}{-\pi u''(x) - 4(1-\pi)u''(y)} \begin{bmatrix} -1 & 2 & 2u''(y)(1-\pi) \\ 2 & -4 & \pi u''(x) \\ 2u''(y)(1-\pi) & \pi u''(x) & \pi u''(x)u''(y)(1-\pi) \end{bmatrix} \begin{bmatrix} u'(x) & 0 \\ -u'(y) & 0 \\ 0 & 2 \end{bmatrix} = \\
& = \frac{1}{\pi u''(x) + 4(1-\pi)u''(y)} \begin{bmatrix} -u'(x) - 2u'(y) & 4u''(y)(1-\pi) \\ 2u'(x) - 4u'(y) & 2\pi u''(x) \\ 2u''(y)(1-\pi) \cdot u'(x) + \pi u''(x)(-u'(y)) & 2\pi u''(x)u''(y)(1-\pi) \end{bmatrix}
\end{aligned}$$

3.

i. Canonical form.

For given $\pi \in (0,1)$, $w_1, w_2 \in \mathbb{R}_{++}$,

$$\begin{aligned}
& \max_{(x,y,m) \in \mathbb{R}_{++}^2 \times \mathbb{R}} \pi \log x + (1-\pi) \log y \quad s.t \\
& \quad w_1 - m - x \geq 0 \quad \lambda_x \\
& \quad w_2 + m - y \geq 0 \quad \lambda_y
\end{aligned}$$

where λ_x and λ_y are the multipliers associated with the first and the second constraint respectively.

ii. The set X and the functions f and g (X open and convex; f and g at least differentiable)

The set $X = \mathbb{R}_{++}^2 \times \mathbb{R}$ is open and convex. The constraint functions are affine and therefore \mathcal{C}^2 . The gradient and the hessian matrix of the objective function are computed below:

	x	y	m
$\pi \log x + (1-\pi) \log y$	$\frac{\pi}{x}$	$\frac{1-\pi}{y}$	0
	x	y	m
$\frac{\pi}{x}$	$-\frac{\pi}{x^2}$	0	0
$\frac{1-\pi}{y}$	0	$-\frac{1-\pi}{y^2}$	0
0	0	0	0

Therefore, the objective function is \mathcal{C}^2 and concave, but not strictly concave.

iii. Existence.

The problem has the same solution set as the following problem:

$$\begin{aligned} \max_{(x,y,m) \in \mathbb{R}_{++}^2 \times \mathbb{R}} \quad & \pi \log x + (1 - \pi) \log y \quad s.t. \\ & w_1 - m - x \geq 0 \quad \lambda_x \\ & w_2 + m - y \geq 0 \quad \lambda_y \\ & \pi \log x + (1 - \pi) \log y \geq \pi \log w_1 + (1 - \pi) \log w_2 \end{aligned}$$

whose constraint set is compact (details left to the reader).

iv. Number of solutions.

The objective function is concave and the constraint functions are affine; uniqueness is not insured on the basis of the sufficient conditions presented in the notes.

v. Necessity of K-T conditions.

Constraint functions are affine and therefore pseudo-concave. Choose $(x, y, m)^{++} = (\frac{w_1}{2}, \frac{w_2}{2}, 0)$.

vi. Sufficiency of Kuhn-Tucker conditions.

f is concave and therefore pseudo-concave and constraint functions are affine and therefore quasi-concave..

vii. K-T conditions.

$$\begin{aligned} D_x L = 0 &\Rightarrow \frac{\pi}{x} - \lambda_x = 0 \\ D_y L = 0 &\Rightarrow \frac{1-\pi}{y} - \lambda_y = 0 \\ D_m L = 0 &\Rightarrow -\lambda_x + \lambda_y = 0 \\ &\min \{w_1 - m - x, \lambda_x\} = 0 \\ &\min \{w_2 + m - y, \lambda_y\} = 0 \end{aligned}$$

viii. Solve the K-T conditions.

Constraints are binding: $\lambda_x = \frac{\pi}{x} > 0$ and $\lambda_y = \frac{1-\pi}{y} > 0$. Then, we get

$$\begin{aligned} \lambda_x &= \frac{\pi}{x} \text{ and } x = \frac{\pi}{\lambda_x} \\ \lambda_y &= \frac{1-\pi}{y} \text{ and } y = \frac{1-\pi}{\lambda_y} \\ \lambda_x &= \lambda_y := \lambda \\ w_1 - m - x &= 0 \\ w_2 + m - y &= 0 \end{aligned}$$

$$\begin{aligned} \lambda_x &= \frac{\pi}{x} \text{ and } x = \frac{\pi}{\lambda_x} \\ \lambda_y &= \frac{1-\pi}{y} \text{ and } y = \frac{1-\pi}{\lambda_y} \\ \lambda_x &= \lambda_y \\ w_1 - m - \frac{\pi}{\lambda} &= 0 \\ w_2 + m - \frac{1-\pi}{\lambda} &= 0 \end{aligned}$$

Then $w_1 - \frac{\pi}{\lambda} = -w_2 + \frac{1-\pi}{\lambda}$ and $\lambda = \frac{1}{w_1 + w_2}$. Therefore

$$\begin{aligned} \lambda_x &= \frac{1}{w_1 + w_2} \\ \lambda_y &= \frac{1}{w_1 + w_2} \\ x &= \pi (w_1 + w_2) \\ y &= (1 - \pi) (w_1 + w_2) \end{aligned}$$

b., c.

Computations of the desired derivatives are straightforward.

4.

i. Canonical form.

$$\begin{aligned} \max_{(x,y) \in \mathbb{R}^2} \quad & -x^2 - y^2 + 4x + 6y \quad s.t. \quad \begin{aligned} -x - y + 6 &\geq 0 & \lambda_1 \\ 2 - y &\geq 0 & \lambda_2 \\ x &\geq 0 & \mu_x \\ y &\geq 0 & \mu_y \end{aligned} \end{aligned}$$

ii. The set X and the functions f and g (X open and convex; f and g at least differentiable)

$X = \mathbb{R}^2$ is open and convex. The constraint functions are affine and therefore C^2 . The gradient and hessian matrix of the objective function are computed below.

$$-x^2 - y^2 + 4x + 6y \quad \begin{array}{cc} x & y \\ -2x + 4 & -2y + 6 \end{array}$$

,

$$\begin{array}{cc} x & y \\ -2x + 4 & -2 \\ -2y + 6 & 0 \end{array} \quad \begin{array}{c} 0 \\ -2 \end{array}$$

Therefore the objective function is C^2 and strictly concave.

iii. Existence.

The constraint set C is nonempty (0 belongs to it) and closed. It is bounded below by 0. y is bounded above by 2. x is bounded above because of the first constraint: $x \leq 6 - y \stackrel{y \geq 0}{\leq} 6$. Therefore C is compact.

iv. Number of solutions.

Since the objective function is strictly concave (and therefore strictly quasi-concave) and the constraint function are affine and therefore quasi-concave, the solution is unique.

v. Necessity of K-T conditions.

Constraints are affine and therefore pseudo-concave. Take $(x^{++}, y^{++}) = (1, 1)$.

vi. Sufficiency of K-T conditions.

The objective function is strictly concave and therefore pseudo-concave. Constraints are affine and therefore quasi-concave.

vii. K-T conditions.

$$\mathcal{L}(x, y, \lambda_1, \lambda_2, \mu_x, \mu_y) = -x^2 - y^2 + 4x + 6y + \lambda_1 \cdot (-x - y + 6) + \lambda_2 \cdot (2 - y) + \mu_x x + \mu_y y.$$

$$\begin{aligned} -2x + 4 - \lambda_1 + \mu_x &= 0 \\ -2y + 6 - \lambda_1 - \lambda_2 + \mu_y &= 0 \\ \min \{-x - y + 6, \lambda_1\} &= 0 \\ \min \{2 - y, \lambda_2\} &= 0 \\ \min \{x, \mu_x\} &= 0 \\ \min \{y, \mu_y\} &= 0 \end{aligned}$$

b.

$$\begin{aligned} +4 - \lambda_1 + \mu_x &= 0 \\ -2y + 6 - \lambda_2 + \mu_y &= 0 \\ \min \{-y + 6, \lambda_1\} &= 0 \\ \min \{2 - y, \lambda_2\} &= 0 \\ \min \{0, \mu_x\} &= 0 \\ \min \{y, \mu_y\} &= 0 \end{aligned}$$

Since $y \leq 2$, we get $-y + 6 > 0$ and therefore $\lambda_1 = 0$. But then $\mu_x = -4$, which contradicts the Kuhn-Tucker conditions above.

c.

$$\begin{aligned} -4 + 4 - \lambda_1 + \mu_x &= 0 \\ -4 + 6 - \lambda_2 + \mu_y &= 0 \\ \min \{-4 + 6, \lambda_1\} &= 0 \\ \min \{2 - 2, \lambda_2\} &= 0 \\ \min \{2, \mu_x\} &= 0 \\ \min \{2, \mu_y\} &= 0 \\ -\lambda_1 + \mu_x &= 0 \\ +2 - \lambda_2 + \mu_y &= 0 \\ \lambda_1 &= 0 \\ \min \{0, \lambda_2\} &= 0 \\ \mu_x &= 0 \\ \mu_y &= 0 \end{aligned}$$

$$\begin{aligned}
\mu_x &= 0 \\
+2 - \lambda_2 &= 0 \\
\lambda_1 &= 0 \\
\min \{0, \lambda_2\} &= 0 \\
\mu_x &= 0 \\
\mu_y &= 0 \\
\mu_x &= 0 \\
\lambda_2 &= 2 \\
\lambda_1 &= 0 \\
\mu_x &= 0 \\
\mu_y &= 0
\end{aligned}$$

Therefore $(x^*, y^*, \lambda_1^*, \lambda_2^*, \mu_x^*, \mu_y^*) = (2, 2, 0, 2, 0)$ is a solution to the Kuhn-Tucker conditions

5.²

5. a.

i. Canonical form.

For given $\delta \in (0, 1)$, $e \in \mathbb{R}_{++}$,

$$\begin{aligned}
\max_{(c_1, c_2, k) \in \mathbb{R}^3} \quad & u(c_1) + \delta u(c_2) \\
s.t. \quad & \\
& e - c_1 - k \geq 0 \quad \lambda_1 \\
& f(k) - c_2 \geq 0 \quad \lambda_2 \\
& c_1 \geq 0 \quad \mu_1 \\
& c_2 \geq 0 \quad \mu_2 \\
& k \geq 0. \quad \mu_3
\end{aligned}$$

ii. The set X and the functions f and g (X open and convex; f and g at least differentiable).

$X = \mathbb{R}^3$ is open and convex. Let's compute the gradient and the Hessian matrix of the second constraint:

$$\begin{array}{cccc}
& c_1 & c_2 & k \\
f(k) - c_2 & 0 & -1 & f'(k) \\
c_1 & c_2 & k & \\
0 & 0 & 0 & 0 \\
-1 & 0 & 0 & 0 \\
f'(k) & 0 & 0 & f''(k) < 0
\end{array}$$

Therefore the second constraint function is C^2 and concave; the other constraint functions are affine.

Let's compute the gradient and the Hessian matrix of the objective functions:

$$\begin{array}{cccc}
& c_1 & c_2 & k \\
u(c_1) + \delta u(c_2) & u'(c_1) & \delta u'(c_2) & 0 \\
c_1 & c_2 & k & \\
u'(c_1) & u''(c_1) & 0 & 0 \\
\delta u'(c_2) & 0 & \delta u''(c_2) & 0 \\
0 & 0 & 0 & 0
\end{array}$$

Therefore the objective function is C^2 and concave.

²Exercise 7 and 8 are taken from David Cass' problem sets for his Microeconomics course at the University of Pennsylvania.

iii. Existence.

The objective function is continuous on \mathbb{R}^3 .

The constraint set is closed because inverse image of closed sets via continuous functions. It is bounded below by 0. It is bounded above: suppose not then

if $c_1 \rightarrow +\infty$, then from the first constraint it must be $k \rightarrow -\infty$, which is impossible;

if $c_2 \rightarrow +\infty$, then from the second constraint and the fact that $f' > 0$, it must be $k \rightarrow +\infty$, violating the first constraint;

if $k \rightarrow +\infty$, then the first constraint is violated.

Therefore, as an application of the Extreme Value Theorem, a solution exists.

iv. Number of solutions.

Since the objective function is concave and the constraint functions are either concave or affine, uniqueness is not insured on the basis of the sufficient conditions presented in the notes.

v. Necessity of K-T conditions.

The constraints are affine or concave. Take $(c_1^{++}, c_2^{++}, k^{++}) = (\frac{e}{4}, \frac{1}{2}f(\frac{e}{4}), \frac{e}{4})$. Then the constraints are verified with strict inequality.

vi. Sufficiency of K-T conditions.

The objective function is concave and therefore pseudo-concave. The constraint functions are either concave or affine and therefore quasi-concave.

vii. K-T conditions.

$\mathcal{L}(c_1, c_2, k, \lambda_1, \lambda_2, \mu_1, \mu_2, \mu_3) := u(c_1) + \delta u(c_2) + \lambda_1(e - c_1 - k) + \lambda_2(f(k) - c_2) + \mu_1 c_1 + \mu_2 c_2 + \mu_3 k$.

$$\begin{cases} u'(c_1) - \lambda_1 + \mu_1 = 0 \\ \delta u'(c_2) - \lambda_2 + \mu_2 = 0 \\ -\lambda_1 + \lambda_2 f'(k) + \mu_3 = 0 \\ \min\{e - c_1 - k, \lambda_1\} = 0 \\ \min\{f(k) - c_2, \lambda_2\} = 0 \\ \min\{c_1, \mu_1\} = 0 \\ \min\{c_2, \mu_2\} = 0 \\ \min\{k, \mu_3\} = 0 \end{cases}$$

viii. Solve the K-T conditions.

Since we are looking for positive solution we get

$$\begin{cases} u'(c_1) - \lambda_1 = 0 \\ \delta u'(c_2) - \lambda_2 = 0 \\ -\lambda_1 + \lambda_2 f'(k) = 0 \\ \min\{e - c_1 - k, \lambda_1\} = 0 \\ \min\{f(k) - c_2, \lambda_2\} = 0 \\ \mu_1 = 0 \\ \mu_2 = 0 \\ \mu_3 = 0 \end{cases}$$

$$\begin{cases} u'(c_1) - \lambda_1 = 0 \\ \delta u'(c_2) - \lambda_2 = 0 \\ -\lambda_1 + \lambda_2 f'(k) = 0 \\ e - c_1 - k = 0 \\ f(k) - c_2 = 0 \end{cases}$$

Observe that from the first two equations of the above system, $\lambda_1, \lambda_2 > 0$.

5. b.

i. Canonical form.

For given $p > 0$, $w > 0$ and $\bar{l} > 0$,

$$\begin{aligned} \max_{(x,l) \in \mathbb{R}^2} \quad & u(x, l) \\ \text{s.t.} \quad & -px - wl + w\bar{l} \geq 0 \\ & \bar{l} - l \geq 0 \\ & x \geq 0 \\ & l \geq 0 \end{aligned}$$

- ii. The set X and the functions f and g (X open and convex; f and g at least differentiable). $X = \mathbb{R}^2$ is open and convex.

The constraint functions are affine and therefore C^2 . The objective function is C^2 and differentiable strictly quasi concave by assumption.

- iii. Existence.

The objective function is continuous on \mathbb{R}^3 .

The constraint set is closed because inverse image of closed sets via continuous functions. It is bounded below by 0. It is bounded above: suppose not then

if $x \rightarrow +\infty$, then from the first constraint ($px + wl = w\bar{l}$), it must be $l \rightarrow -\infty$, which is impossible. Similar case is obtained, if $l \rightarrow +\infty$.

Therefore, as an application of the Extreme Value Theorem, a solution exists.

- iv. Number of solutions.

The budget set is convex. The function is differentiable strictly quasi concave and therefore strictly quasi-concave and the solution is unique.

- v. Necessity of K-T conditions.

The constraints are pseudo-concave $(x^{++}, l^{++}) = (\frac{w\bar{l}}{3p}, \frac{\bar{l}}{3})$ satisfies the constraints with strict inequalities.

- vi. Sufficiency of K-T conditions.

The objective function is differentiable strictly quasi-concave and therefore pseudo-concave. The constraint functions are quasi-concave

- vii. K-T conditions.

$$\mathcal{L}(c_1, c_2, k, \lambda_1, \lambda_2, \lambda_3, \lambda_4; p, w, \bar{l}) := u(x, l) + \lambda_1(-px - wl + w\bar{l}) + \lambda_2(\bar{l} - l) + \lambda_3x + \lambda_4l.$$

$$\begin{aligned} D_x u - \lambda_1 p + \lambda_3 &= 0 \\ D_l u - \lambda_1 w - \lambda_2 + \lambda_4 &= 0 \\ \min \{-px - wl + w\bar{l}, \lambda_1\} &= 0 \\ \min \{\bar{l} - l, \lambda_2\} &= 0 \\ \min \{x, \lambda_3\} &= 0 \\ \min \{l, \lambda_4\} &= 0 \end{aligned}$$

- viii. Solve the K-T conditions.

Since we are looking for solutions at which $x > 0$ and $0 < l < \bar{l}$, we get

$$\begin{aligned} D_x u - \lambda_1 p + \lambda_3 &= 0 \\ D_l u - \lambda_1 w - \lambda_2 + \lambda_4 &= 0 \\ \min \{-px - wl + w\bar{l}, \lambda_1\} &= 0 \\ \lambda_2 &= 0 \\ \lambda_3 &= 0 \\ \lambda_4 &= 0 \end{aligned}$$

$$\begin{aligned} D_x u - \lambda_1 p &= 0 \\ D_l u - \lambda_1 w &= 0 \\ \min \{-px - wl + w\bar{l}, \lambda_1\} &= 0 \end{aligned}$$

and then $\lambda_1 > 0$ and

$$\begin{cases} D_x u - \lambda_1 p = 0 \\ D_l u - \lambda_1 w = 0 \\ -px - wl + w\bar{l} = 0 \end{cases}.$$

7.

- a.

Let's apply the Implicit Function Theorem ($:=$ IFT) to the conditions found in Exercise 7.(a). Writing them in the usual informal way we have:

	c_1	c_2	k	λ_1	λ_2	e	a
$u'(c_1) - \lambda_1 = 0$		$u''(c_1)$			-1		
$\delta u'(c_2) - \lambda_2 = 0$			$\delta u''(c_2)$			-1	
$-\lambda_1 + \lambda_2 f'(k) = 0$			$\lambda_2 f''(k)$	-1	$f'(k)$		$\lambda_2 \alpha k^{\alpha-1}$
$e - c_1 - k = 0$	-1		-1			1	
$f(k) - c_2 = 0$		-1	$f'(k)$				k^α

To apply the IFT, we need to check that the following matrix has full rank

$$M := \begin{bmatrix} u''(c_1) & & & -1 & & \\ & \delta u''(c_2) & & & -1 & \\ & & \lambda_2 f''(k) & -1 & f'(k) & \\ -1 & & -1 & & & \\ & -1 & f'(k) & & & \end{bmatrix}$$

Suppose not then there exists $\Delta := (\Delta c_1, \Delta c_2, \Delta k, \Delta \lambda_1, \Delta \lambda_2) \neq 0$ such that $M\Delta = 0$, i.e.,

$$\begin{cases} u''(c_1) \Delta c_1 + & & -\Delta \lambda_1 & = 0 \\ & \delta u''(c_2) \Delta c_2 + & & -\Delta \lambda_2 & = 0 \\ & & \lambda_2 f''(k) \Delta k + & -\Delta \lambda_1 + f'(k) \Delta \lambda_2 & = 0 \\ -\Delta c_1 + & & -\Delta k & = 0 \\ & -\Delta c_2 + & f'(k) \Delta k + & & = 0 \end{cases}$$

Recall that

$$[M\Delta = 0 \Rightarrow \Delta = 0] \quad \text{iff} \quad M \quad \text{has full rank.}$$

The idea of the proof is either you prove directly $[M\Delta = 0 \Rightarrow \Delta = 0]$, or you 1. assume $M\Delta = 0$ and $\Delta \neq 0$ and you get a contradiction.

If we define $\Delta c := (\Delta c_1, \Delta c_2)$, $\Delta \lambda := (\Delta \lambda_1, \Delta \lambda_2)$, $D^2 := \begin{bmatrix} u''(c_1) \Delta c_1 & \\ & \delta u''(c_2) \Delta c_2 \end{bmatrix}$, the above system can be rewritten as

$$\begin{cases} D^2 \Delta c + & & -\Delta \lambda & = 0 \\ & \lambda_2 f''(k) \Delta k + & [-1, f'(k)] \Delta \lambda & = 0 \\ -\Delta c + & \begin{bmatrix} -1 \\ f'(k) \end{bmatrix} \Delta k & & = 0 \end{cases}.$$

$$\begin{cases} \Delta c^T D^2 \Delta c + & & -\Delta c^T \Delta \lambda & = 0 & (1) \\ & \Delta k \lambda_2 f''(k) \Delta k + & \Delta k [-1, f'(k)] \Delta \lambda & = 0 & (2) \\ -\Delta \lambda^T \Delta c + & \Delta \lambda^T \begin{bmatrix} -1 \\ f'(k) \end{bmatrix} \Delta k & & = 0 & (3) \end{cases}$$

$$\Delta c^T D^2 \Delta c \stackrel{(1)}{=} \Delta c^T \Delta \lambda = -\Delta \lambda^T \Delta c \stackrel{(3)}{=} \Delta \lambda^T \begin{bmatrix} -1 \\ f'(k) \end{bmatrix} \Delta k \stackrel{(2)}{=} \Delta k [-1, f'(k)] \Delta \lambda =$$

$$= \Delta k \lambda_2 f''(k) \Delta k > 0,$$

while $\Delta c^T D^2 \Delta c = (\Delta c_1)^2 u''(c_1) + (\Delta c_2)^2 \delta u''(c_1) < 0$. since we got a contradiction, M has full rank.

Therefore, in a neighborhood of the solution we have

$$D_{(e,a)}(c_1, c_2, k, \lambda_1, \lambda_2) = - \begin{bmatrix} u''(c_1) & & & -1 & & \\ & \delta u''(c_2) & & & -1 & \\ & & \lambda_2 f''(k) & -1 & f'(k) & \\ -1 & & -1 & & & \\ & -1 & f'(k) & & & \end{bmatrix}^{-1} \begin{bmatrix} \\ \\ \lambda_2 \alpha k^{\alpha-1} \\ 1 \\ k^\alpha \end{bmatrix}.$$

To compute the inverse of the above matrix, we can use the following fact about the inverse of partitioned matrix (see Goldberger, (1964), page 27:

Let A be an $n \times n$ nonsingular matrix partitioned as

Defined $D^2 := \begin{bmatrix} D_x^2 & D_{xl}^2 \\ D_{xl}^2 & D_l^2 \end{bmatrix}$, $q := \begin{bmatrix} -p \\ -w \end{bmatrix}$, we have $M := \begin{bmatrix} D^2 & -q \\ -q^T & \end{bmatrix}$.

Suppose not then there exists $\Delta := (\Delta y, \Delta \lambda) \in (\mathbb{R}^2 \times \mathbb{R}) \setminus \{0\}$ such that $M\Delta = 0$, i.e.,

$$\begin{cases} D^2 \Delta y & -q \Delta \lambda & = 0 & (1) \\ -q^T \Delta y & & = 0 & (2) \end{cases}$$

We are going to show

Step 1. $\Delta y \neq 0$; Step 2. $Du \cdot \Delta y = 0$; Step 3. It is not the case that $\Delta y^T D^2 \Delta y < 0$.

These results contradict the assumption about u .

Step 1.

Suppose $\Delta y = 0$. Since $q \gg 0$, from (1), we get $\Delta \lambda = 0$, and therefore $\Delta = 0$, a contradiction.

Step 2.

From the First Order Conditions, we have

$$Du - \lambda_1 q = 0 \quad (3).$$

$$Du \Delta y \stackrel{(3)}{=} \lambda_1 q \Delta y \stackrel{(2)}{=} 0.$$

Step 3.

$$\Delta y^T D^2 \Delta y \stackrel{(1)}{=} \Delta y^T q \Delta \lambda \stackrel{(2)}{=} 0.$$

Therefore, in a neighborhood of the solution we have

$$D_{(p,w,\bar{l})}(x, l, \lambda_1) = - \begin{bmatrix} D_x^2 & D_{xl}^2 & -p \\ D_{xl}^2 & D_l^2 & -w \\ -p & -w & \end{bmatrix}^{-1} \begin{bmatrix} -\lambda_1 \\ -\lambda_1 \\ -1 \quad \bar{l} - l \quad w \end{bmatrix}.$$

Unfortunately, here we cannot use the formula seen in the Exercise 4 (a) because the Hessian of the utility function is not necessarily nonsingular. We can invert the matrix using the definition of inverse. (For the inverse of a partitioned matrix with this characteristics see also Dhrymes, P. J., (1978), *Mathematics for Econometrics*, 2nd edition, Springer-Verlag, New York, NY, Addendum pages 142-144.

With obvious notation and using Maple, we get

$$\begin{bmatrix} d_x & d & -p \\ d & d_l & -w \\ -p & -w & 0 \end{bmatrix}^{-1} = \begin{bmatrix} \frac{w^2}{d_x w^2 - 2dpw + p^2 d_l} & -p \frac{w}{d_x w^2 - 2dpw + p^2 d_l} & -\frac{-dw + pd_l}{d_x w^2 - 2dpw + p^2 d_l} \\ -p \frac{w}{d_x w^2 - 2dpw + p^2 d_l} & \frac{p^2}{d_x w^2 - 2dpw + p^2 d_l} & \frac{-d_x w + dp}{d_x w^2 - 2dpw + p^2 d_l} \\ -\frac{-dw + pd_l}{d_x w^2 - 2dpw + p^2 d_l} & \frac{-d_x w + dp}{d_x w^2 - 2dpw + p^2 d_l} & \frac{-d_x d_l + d^2}{d_x w^2 - 2dpw + p^2 d_l} \end{bmatrix}$$

Therefore,

$$D_{(p,w,\bar{l})}(x, l, \lambda_1) =$$

$$- \begin{bmatrix} \frac{w^2}{d_x w^2 - 2dpw + p^2 d_l} & -p \frac{w}{d_x w^2 - 2dpw + p^2 d_l} & -\frac{-dw + pd_l}{d_x w^2 - 2dpw + p^2 d_l} \\ -p \frac{w}{d_x w^2 - 2dpw + p^2 d_l} & \frac{p^2}{d_x w^2 - 2dpw + p^2 d_l} & \frac{-d_x w + dp}{d_x w^2 - 2dpw + p^2 d_l} \\ -\frac{-dw + pd_l}{d_x w^2 - 2dpw + p^2 d_l} & \frac{-d_x w + dp}{d_x w^2 - 2dpw + p^2 d_l} & \frac{-d_x d_l + d^2}{d_x w^2 - 2dpw + p^2 d_l} \end{bmatrix} \begin{bmatrix} -\lambda_1 & 0 & 0 \\ 0 & -\lambda_1 & 0 \\ -1 & \bar{l} - l & w \end{bmatrix} =$$

$$= \begin{bmatrix} -\frac{-w^2 \lambda_1 - dw + pd_l}{d_x w^2 - 2dpw + p^2 d_l} & -\frac{pw \lambda_1 - ldw + lpd_l}{d_x w^2 - 2dpw + p^2 d_l} & \frac{-dw + pd_l}{d_x w^2 - 2dpw + p^2 d_l} w \\ \frac{-pw \lambda_1 - d_x w + dp}{d_x w^2 - 2dpw + p^2 d_l} & \frac{p^2 \lambda_1 - ld_x w + ldp}{d_x w^2 - 2dpw + p^2 d_l} & -\frac{-d_x w + dp}{d_x w^2 - 2dpw + p^2 d_l} w \\ -\frac{-\lambda_1 dw + \lambda_1 pd_l + d_x d_l - d^2}{d_x w^2 - 2dpw + p^2 d_l} & -\frac{-\lambda_1 d_x w + \lambda_1 dp - ld_x d_l + ld^2}{d_x w^2 - 2dpw + p^2 d_l} & -\frac{-d_x d_l + d^2}{d_x w^2 - 2dpw + p^2 d_l} w \end{bmatrix}$$

$$D_p l = \frac{-pw \lambda_1 - d_x w + dp}{d_x w^2 - 2dpw + p^2 d_l}$$

$$D_w l = \frac{p^2 \lambda_1 - ld_x w + ldp}{d_x w^2 - 2dpw + p^2 d_l}.$$

The sign of these expressions is ambiguous, unless other assumptions are made.

7.

\Rightarrow

Since f is concave, then

$$f\left(\frac{1}{2}x + \frac{1}{2}y\right) \geq \frac{1}{2}f(x) + \frac{1}{2}f(y).$$

Since f is homogenous of degree 1, then

$$f\left(\frac{1}{2}(x+y)\right) = \frac{1}{2}f(x+y).$$

Therefore,

$$f(x+y) \geq f(x) + f(y).$$

[\Leftarrow]

Since f is homogenous of degree 1, then for any $z \in \mathbb{R}^2$ and any $a \in \mathbb{R}_+$, we have

$$f(az) = af(z). \quad (21.12)$$

By assumption, we have that

$$\text{for any } x, y \in \mathbb{R}^2, \quad f(x+y) \geq f(x) + f(y). \quad (21.13)$$

Then, for any $\lambda \in [0, 1]$,

$$f((1-\lambda)x + \lambda y) \stackrel{(21.13)}{\geq} f((1-\lambda)x) + f(\lambda y) \stackrel{(21.12)}{=} (1-\lambda)f(x) + \lambda f(y),$$

as desired.

References

- Aliprantis, C. D. and O. Burkinshaw, (1990), *Principles of Real Analysis*, 2nd edition, Academic Press, St Louis, Missouri.
- Apostol, T. M., (1967), *Calculus, Volume 1*, 2nd edition, John Wiley & Sons, New York, NY.
- Apostol, T. M., (1974), *Mathematical Analysis*, 2nd edition, Addison-Wesley Publishing Company, Reading, MA.
- Azariadis, C., (2000), *Intertemporal Macroeconomics*, Blackwell Publisher Ltd, Oxford, UK.
- Balasko, Y., (1988), *Foundations of the Theory of General Equilibrium*, Academic Press, Boston.
- Bazaraa, M.S. and C. M. Shetty, (1976), *Foundations of Optimization*, Springer-Verlag, Berlin.
- Bartle, R. G., (1964), *The Elements of Real Analysis*, John Wiley & Sons, New York, NY.
- Beavis, B. and I. Dobbs, (1990), *Optimization and stability theory for economic analysis*, Cambridge University Press, Cambridge, UK.
- Cass D., (1991), *Nonlinear Programming for Economists*, mimeo, University of Pennsylvania.
- de la Fuente, A., (2000), *Mathematical Methods and Models for Economists*, Cambridge University Press, Cambridge, UK.
- Dhrymes, P. J., (1978), *Mathematics for Econometrics*, 2nd edition Springer-Verlag, New York, NY.
- El-Hodiri, M. A., (1991), *Extrema of smooth functions*, Springer-Verlag, Berlin.
- Francini, E., (2004), *Esercitazioni del corso di Analisi Superiore*, mimeo, Università degli Studi di Firenze.
- Girsanov, I.V., (1972), *Lectures on Mathematical Theory of Extremum Problems*, Springer-Verlag, Berlin and New York.
- Goldberger, A.S, (1964), *Econometric Theory*, John Wiley & Sons, New York, NY.
- Gori, F., (2014), *Notes for the course "Mathematical Economics"*, mimeo, Università degli Studi di Firenze.
- Hildebrand, W., (1974), *Core and equilibria of a large economy*, Princeton University Press, Princeton, NJ.
- Jahn, J., (1996), *Introduction to the Theory of Nonlinear Optimization, 2nd revised edition*, Springer-Verlag, Berlin.
- Herstein, I. N., (1975), *Topics in Algebra*, 2nd edition, John Wiley & Sons, New York, NY.
- Hirsch, M. W. and S. Smale, (1974), *Differential Equations, Dynamical Systems and Linear Algebra*, Academic Press, New York, NY.
- Hoffman, K. and R. Kunze, (1971), *Linear Algebra*, 2nd edition, Prentice-Hall, Inc., Englewood Cliffs, New Jersey.
- Lang S. (1971), *Linear Algebra*, second edition, Addison Wesley, Reading.
- Lang, S., (1986), *Introduction to Linear Algebra*, 2nd edition, Springer-Verlag, New York, NY.
- Lipschutz, S., (1989), *3000 Solved Problems in Linear Algebra*, McGraw-Hill, New York, NY.
- Lipschutz, S., (1991), *Theory and Problems in Linear Algebra*, second edition, Schaum's Outline Series, McGraw-Hill, New York, NY.
- Lipschutz, S., (1965), *General Topology*, McGraw-Hill, New York, NY.

- Mangasarian O. L. (1994), *Nonlinear Programming*, SIAM, Philadelphia.
- McLean, R., (1985), Class notes for the course of Mathematical Economics (708), University of Pennsylvania, Philadelphia, PA, mimeo.
- Montrucchio, L., (1998), *Introduzione alle teoria delle scelte*, (in Italian), Carocci Editore, Rome, Italy.
- Moore, J. C., (1999), *Mathematical Methods for Economic Theory 1*, Springer-Verlag, Berlin.
- Moore, J. C., (1999), *Mathematical Methods for Economic Theory 2*, Springer-Verlag, Berlin.
- Ok, E. A., (2007), *Real Analysis with Economic Applications*, Princeton University Press, Princeton NJ.
- Ross, K. A., (1980), *Elementary analysis: The theory of calculus*, Springer-Verlag, New York, NY.
- Rudin, W., (1976), *Principles of Mathematical Analysis*, 3rd edition, McGraw-Hill, New York, NY.
- Simon, C. S. (1986), Scalar and vector maximization: calculus techniques with economic applications, in Reiter S, *Studies in mathematical economics*, The Mathematical Association of America, p. 62-159
- Simon, C. P. and L. Blume, (1994), *Mathematics for Economists*, Norton, New York
- Simmons, G. F., (1963), *Introduction to Topology and Modern Analysis*, McGraw-Hill, New York.
- Smith, L., (1992), *Linear Algebra*, 2nd edition, Springer-Verlag, New York, NY.
- Stokey, N. C., Lucas, E. C. (with Prescott, E. C.), (1989), *Recursive methods in Economic Dynamics*, Harvard University Press, Cambridge, MA.
- Sydsaeter, K., (1981), *Topics in Mathematical Analysis for Economists*, Academic Press, London, UK.
- Taylor, A. E. and W. R. Mann, (1984), *Advanced Calculus*, 3rd ed., John Wiley & Sons, New York, NY.
- Villanacci, A., (in progress), Basic Convex Analysis, mimeo, Università degli Studi di Firenze.
- Villanacci, A., (in progress), Notes on Mathematics 3: Linear Algebra, Linear and Nonlinear Systems of Differential Equations, mimeo, Università degli Studi di Firenze.
- Villanacci, A., (in progress), Notes on Mathematics 4, Topology, Banach Spaces, Differential Calculus and Measure, mimeo, Università degli Studi di Firenze.
- Wrede, R. C. and M. Spiegel (2002), *Theory and Problems of Advanced Calculus*, 2nd edition, Schaum's outlines, McGraw Hill, New York.